

Quantum Lorentz group having Gauss decomposition property

S.L. Woronowicz^{*†}

ETH Zürich

and

S. Zakrzewski^{†‡}

TU Clausthal

April 13, 2002

Abstract

A new deformation of $SL(2, \mathbf{C})$ (considered as a real Lie group) is constructed and shown to have a Gauss type decomposition. The groups entering this decomposition are identified as $E_\mu(2)$ and its Pontryagin dual $\widehat{E}_\mu(2)$. The whole group is the double group built over $E_\mu(2)$.

0 Introduction

The first quantum deformation of Lorentz group was described in [5]. The theory presented in that paper based on the following four assumptions:

1. The tensor square of the (two-dimensional) spinor representation splits into a direct sum of two components, one of which is a one-dimensional trivial representation.
2. The tensor product of the spinor representation by the complex conjugate one is irreducible and does not depend on the order of factors.
3. The quantum Lorentz group contains $S_\mu U(2)$ as a subgroup (Throughout the paper, $\mu \in]0, 1[$ denotes the deformation parameter).
4. The quantum Lorentz group is not a proper subgroup of a quantum group satisfying the above three conditions.

^{*}Supported in equal parts by the grant of the Ministry of Education of Poland and by Schweizerischer Nationalfonds

[†]Supported in equal parts by the grant of the Ministry of Education of Poland and by Alexander von Humboldt Stiftung

[‡]Permanent address: Department of Mathematical Methods in Physics, Faculty of Physics, University of Warsaw, Hoża 74, 00-682 Warszawa, Polska

Communicated by H. Araki, August 12, 1991.

1991 Mathematical Subject Classification : 20N99, 57T05.

The role of the last assumption (which was used but not explicitly formulated in [5]) is explained in [9]. It was shown that the above requirements determine the quantum Lorentz group completely (in fact there are two solutions, the choice consists in declaring which of the two spinor representation is the fundamental one).

It turns out that deleting the third assumption we still have a limited number of possibilities. We refer to our paper [9] where we classify (on the Hopf *-algebra level) all quantum group of 2×2 matrices satisfying the assumptions 1, 2 and 4. In the present paper we investigate in details one of the possibilities distinguished by the fact that the corresponding quantum Lorentz group exhibits Gauss-type decomposition.

The classical Gauss decomposition [2] says that except a closed set of measure zero any element g of $SL(2, \mathbf{C})$ has a unique decomposition of the form

$$g = n_+ h n_-$$

where n_+ (n_- resp.) belongs to the subgroup N_+ (N_- resp.) of all upper (lower resp.) triangular matrices with one on the diagonal and h belongs to the subgroup H of all diagonal matrices. In short

$$G' = N_+ H N_- \tag{1}$$

where G' is an open full measure subset of $SL(2, \mathbf{C})$.

In the quantum case however the subgroups analogous to N_{\pm} do not exist (because of the nontrivial commutation relations between diagonal and off-diagonal elements). To deal with this difficulty one may include the diagonal part of the decomposition into the both factors N_{\pm} considering the decomposition of the form

$$G' = M_+ M_-$$

where M_+ (M_- resp.) is the subgroup of all upper (lower) triangular matrices. This form of Gauss decomposition was proposed by Faddeev Reshetikhin and Takhtajan [4]. One should notice however that in this case the decomposition is not unique.

In this paper we shall prove the quantum version of the decomposition of the form

$$G' = G_1 G_2$$

where G_1 (G_2 resp) is the subgroup of upper (lower resp.) triangular matrices having unitary (positive resp.) elements on the diagonal. Comparing with (1) we see that H has been split into H_{unitary} included into N_+ and H_{positive} included into N_- . We would like to point out that in certain sense the Gauss decomposition in the quantum case is more satisfactory than the classical one: due to the commutation relations the diagonal elements are invertible so the decomposition has the global character with no exceptions.

G_1 is the (double covering) of the group of motions of the euclidean plane, G_2 is isomorphic to the group of transformations of the plane generated by dilations and translations. In the quantum case the second group is the Pontriagin dual of the first one. Due to this fact we may use the double group construction (cf [3], [5]). As a result we get the quantum Lorentz group on the C^* -level. In what follows this group will be called G_{μ} .

The paper is composed in the following way. In Section 1 we introduce the Hopf *-algebra being a quantum deformation of the polynomial algebra on $SL(2, \mathbf{C})$. It is generated by four

elements α, β, γ and δ satisfying certain commutation relations. The relations completed by a spectral condition formulated at the beginning of Section 4 are the main input of our theory. The anatomy of the commutation relations is investigated in Section 2. The main aim of this Section is to assign the strong meaning to the relations: since the involved elements are unbounded one has to define carefully the domains where the relations are supposed to hold. In Section 3 we consider the Hilbert space representations of our relations. The main result shows that α and δ are invertible and that $\text{Sp } \beta\gamma$ exhibits a radius quantization. Section 4 is devoted to the Gauss decomposition. In Section 5 we repeat the double group construction of [5] replacing the Iwasawa decomposition by that of Gauss to achieve our main goal. Namely we define the universal C^* -algebra A “generated” by four elements α, β, γ and δ affiliated with it, satisfying the commutation relations and the spectral condition. It plays the role of the algebra of “all continuous functions on G_μ vanishing at infinity”. The algebra A is equipped with the comultiplication $\Phi \in \text{Mor}(A, A \otimes A)$ acting on the generators in the standard way.

The notation used in this paper is explained in [7] and [8]. In particular

$$\overline{\mathbf{C}}^\mu = \{t \in \mathbf{C} : t = 0 \text{ or } |t| \in \mu^{\mathbf{Z}}\},$$

$F_\mu(\cdot)$ is the continuous function on $\overline{\mathbf{C}}^\mu$ introduced by

$$F_\mu(t) = \prod_{k=0}^{\infty} \frac{1 + \mu^{2k\bar{t}}}{1 + \mu^{2k}t}.$$

All C^* -algebras are concrete (i.e: acting on Hilbert spaces). For any Hilbert space H , $C^*(H)$ denotes the set of all C^* -algebras contained in $B(H)$. Let $A \in C^*(H_A)$, $B \in C^*(H_B)$ and ϕ be a representation of A acting on H_B . We remind that $\phi \in \text{Mor}(A, B)$ if and only if $\phi(A)B$ is contained and dense in B . The symbol “ η ” denotes the C^* -affiliation relation ([1], [6]): for any $A \in C^*(H)$ and any closed operator T acting on H , $T\eta A$ if and only if

$$\begin{aligned} [T(I + T^*T)^{-\frac{1}{2}}]A &\subset A, \\ A[T(I + T^*T)^{-\frac{1}{2}}] &\subset A, \\ [(I + T^*T)^{-\frac{1}{2}}]A &\text{ is dense in } A. \end{aligned}$$

The actions of morphisms admit the natural extension to the affiliated elements: if $\phi \in \text{Mor}(A, B)$ and $T\eta A$ then $\phi(T)$ is a well defined element affiliated with B .

The closure of the sum of two operators R and S will be denoted by $R\dot{+}S$

1 Hopf $*$ -algebra level

We fix deformation parameter $\mu \in]0, 1]$. Let \mathcal{A} be the unital $*$ -algebra generated by four elements α, β, γ and δ satisfying the following 17 relations:

$$\left. \begin{aligned} \alpha\beta &= \mu\beta\alpha, & \beta\gamma &= \gamma\beta, & \alpha\gamma &= \mu\gamma\alpha, \\ \beta\delta &= \mu\delta\beta, & \gamma\delta &= \mu\delta\gamma, \\ \alpha\delta - \mu\beta\gamma &= I, & \delta\alpha - \mu^{-1}\beta\gamma &= I \end{aligned} \right\} \quad (2)$$

$$\left. \begin{aligned} \alpha\beta^* &= \mu\beta^*\alpha, & \beta\gamma^* &= \gamma^*\beta, & \alpha\gamma^* &= \mu^{-1}\gamma^*\alpha, \\ \beta\delta^* &= \mu^{-1}\delta^*\beta, & \alpha\delta^* &= \delta^*\alpha, & \gamma\delta^* &= \mu\delta^*\gamma, \\ \alpha\alpha^* &= \alpha^*\alpha, & \beta\beta^* &= \beta^*\beta, & \gamma\gamma^* &= \gamma^*\gamma, \\ & & \delta\delta^* &= \delta^*\delta. \end{aligned} \right\} \quad (3)$$

The relations may be rewritten in a compact matrix form. Let $u = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ and $\bar{u} = \begin{pmatrix} \alpha^* & \beta^* \\ \gamma^* & \delta^* \end{pmatrix}$. Then

$$u \oplus u = \begin{pmatrix} \alpha\alpha & \alpha\beta & \beta\alpha & \beta\beta \\ \alpha\gamma & \alpha\delta & \beta\gamma & \beta\delta \\ \gamma\alpha & \gamma\beta & \delta\alpha & \delta\beta \\ \gamma\gamma & \gamma\delta & \delta\gamma & \delta\delta \end{pmatrix},$$

and one can easily verify that relations (2) are equivalent to

$$(u \oplus u)E = EI \quad \text{and} \quad IE^* = E^*(u \oplus u)$$

where

$$E = \begin{pmatrix} 0 \\ 1 \\ -\mu \\ 0 \end{pmatrix} \quad \text{and} \quad E^* = (0, 1, -\mu, 0).$$

Similarly

$$u \oplus \bar{u} = \begin{pmatrix} \alpha\alpha^* & \alpha\beta^* & \beta\alpha^* & \beta\beta^* \\ \alpha\gamma^* & \alpha\delta^* & \beta\gamma^* & \beta\delta^* \\ \gamma\alpha^* & \gamma\beta^* & \delta\alpha^* & \delta\beta^* \\ \gamma\gamma^* & \gamma\delta^* & \delta\gamma^* & \delta\delta^* \end{pmatrix}, \quad \bar{u} \oplus u = \begin{pmatrix} \alpha^*\alpha & \alpha^*\beta & \beta^*\alpha & \beta^*\beta \\ \alpha^*\gamma & \alpha^*\delta & \beta^*\gamma & \beta^*\delta \\ \gamma^*\alpha & \gamma^*\beta & \delta^*\alpha & \delta^*\beta \\ \gamma^*\gamma & \gamma^*\delta & \delta^*\gamma & \delta^*\delta \end{pmatrix}.$$

and relations (3) are equivalent to

$$X(\bar{u} \oplus u) = (u \oplus \bar{u})X,$$

where

$$X = \begin{pmatrix} \mu & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \mu \end{pmatrix}$$

Using the matrix form of the commutation relations one may prove

Theorem 1.1 1. *There exists unique *-algebra homomorphism*

$$\Phi : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$$

such that

$$\begin{pmatrix} \Phi(\alpha) & \Phi(\beta) \\ \Phi(\gamma) & \Phi(\delta) \end{pmatrix} = u \oplus u = \begin{pmatrix} \alpha \otimes \alpha + \beta \otimes \gamma & \alpha \otimes \beta + \beta \otimes \delta \\ \gamma \otimes \alpha + \delta \otimes \gamma & \gamma \otimes \beta + \delta \otimes \delta \end{pmatrix} \quad (4)$$

2. There exists unique $*$ -character

$$e : \mathcal{A} \rightarrow \mathbf{C}$$

such that

$$\begin{pmatrix} e(\alpha), e(\beta) \\ e(\gamma), e(\delta) \end{pmatrix} = \begin{pmatrix} 1, 0 \\ 0, 1 \end{pmatrix}$$

3. There exists unique linear antimultiplicative mapping

$$\kappa : \mathcal{A} \rightarrow \mathcal{A}$$

such that

$$\begin{pmatrix} \kappa(\alpha), \kappa(\beta) \\ \kappa(\gamma), \kappa(\delta) \end{pmatrix} = u^{-1} = \begin{pmatrix} \delta & , & -\mu^{-1}\beta \\ -\mu\gamma & , & \alpha \end{pmatrix}$$

$$\begin{pmatrix} \kappa(\alpha^*), \kappa(\beta^*) \\ \kappa(\gamma^*), \kappa(\delta^*) \end{pmatrix} = (\bar{u})^{-1} = \begin{pmatrix} \delta^* & , & -\mu\beta^* \\ -\mu^{-1}\gamma^* & , & \alpha^* \end{pmatrix}$$

4. $(\mathcal{A}, \Phi, e, \kappa)$ is a Hopf $*$ -algebra

In the rest of the paper we shall construct a C^* -algebra “generated” by four distinguished elements α, β, γ and δ affiliated with it such that the relations (2) and (3) are satisfied in a certain strong sense. We shall also prove that there exists unique C^* -algebra category morphism $\Phi \in \text{Mor}(A, A \otimes A)$ acting on distinguished elements α, β, γ and δ in the way described by (4).

2 (p, q) -commuting normal operators

Throughout this section, p and q will denote two fixed strictly positive real numbers. The purpose of this section is to give a meaning to the following relations

$$\begin{aligned} RS &= pSR, \\ RS^* &= qS^*R, \end{aligned} \tag{5}$$

where R and S are normal (unbounded) operators acting on a Hilbert space H . We shall use the following

Definition 2.1 *Let R and S be normal operators acting on a Hilbert space H . We say that (R, S) is a (p, q) -commuting pair if*

1. $|R|$ and $|S|$ strongly commute
2. $(\text{Phase } R)(\text{Phase } S) = (\text{Phase } S)(\text{Phase } R)$
3. On $(\ker R)^\perp$ we have

$$(\text{Phase } R) |S| (\text{Phase } R)^* = \sqrt{pq} |S|$$

4. On $(\ker S)^\perp$ we have

$$(\text{Phase } S) |R| (\text{Phase } S)^* = \sqrt{q/p} |R|$$

The set of all (p, q) -commuting pairs of normal operators acting on a Hilbert space H will be denoted by $\mathcal{C}_H(p, q)$ or simply by $\mathcal{C}(p, q)$ when it makes no confusion.

If $(R, S) \in \mathcal{C}(p, q)$ then the composition $R \circ S$ is closeable and its closure will be denoted by RS . One can easily verify that

$$\begin{aligned} \text{Phase}(RS) &= (\text{Phase } R)(\text{Phase } S) \\ |RS| &= \sqrt{p/q} |R||S|. \end{aligned} \quad (6)$$

The operator RS is normal if and only if $q = 1$. Similarly one may consider products SR, R^*S, S^*R , etc.

Let $(R, S) \in \mathcal{C}(p, q)$. Then one can easily verify that

$$(R, S^*) \in \mathcal{C}(q, p) \quad (7)$$

$$(R, \text{Phase } S) \in \mathcal{C}(\sqrt{p/q}, \sqrt{q/p}) \quad (8)$$

$$(R, |S|) \in \mathcal{C}(\sqrt{pq}, \sqrt{pq}) \quad (9)$$

$$(S, R) \in \mathcal{C}(p^{-1}, q) \quad (10)$$

Moreover if $\ker S = \{0\}$ then

$$(R, S^{-1}) \in \mathcal{C}(p^{-1}, q^{-1}) \quad (11)$$

If $(R, S_1) \in \mathcal{C}(p_1, q_1)$, $(R, S_2) \in \mathcal{C}(p_2, q_2)$ and $(S_1, S_2) \in \mathcal{C}(p_3, q_3)$ then using (6) to compute $\text{Phase}(S_1 S_2)$ and $|S_1 S_2|$ we see that

$$(R, S_1 S_2) \in \mathcal{C}(p_1 p_2, q_1 q_2). \quad (12)$$

Similarly

$$(R_1 R_2, S) \in \mathcal{C}(p_1 p_2, q_1 q_2). \quad (13)$$

provided $(R_1, S) \in \mathcal{C}(p_1, q_1)$, $(R_2, S) \in \mathcal{C}(p_2, q_2)$ and $(R_1, R_2) \in \mathcal{C}(p_3, q_3)$.

Two normal operators R and S strongly commute (have joint spectral decomposition) if and only if $(R, S) \in \mathcal{C}(1, 1)$.

We shall use the following version of Theorems 2.1 and 2.2 of [7].

Theorem 2.2 *Let $(R, S) \in \mathcal{C}(\mu^{-2}, 1)$. Assume that $\ker R = \{0\}$. Then the following three conditions are equivalent:*

- (i) $R + S$ admits a normal extension,
- (ii) $R \dot{+} S$ is normal,
- (iii) $\text{Sp}(R^{-1}S) \subset \overline{\mathbf{C}}^\mu$.

Moreover, if one of the above conditions holds then $R \dot{+} S$ is unitarily equivalent to R :

$$R \dot{+} S = F_\mu(R^{-1}S) R F_\mu(R^{-1}S)^* \quad (14)$$

Proof: If $S = 0$ then $R^{-1}S = 0$, $F_\mu(R^{-1}S) = I$. Hence the conditions (i)–(iii) are satisfied and the formula (14) holds. Therefore we may assume that $\ker S = \{0\}$. In this case our theorem coincides with Theorem 2.1 of [7].

Q.E.D.

We also have

Theorem 2.3 Let $(R, S) \in \mathcal{C}(\mu^{-2}, 1)$. Assume that $\ker S = \{0\}$. Then the following three conditions are equivalent:

- (i) $R + S$ admits a normal extension,
- (ii) $R \dot{+} S$ is normal,
- (iii) $\text{Sp}(RS^{-1}) \subset \overline{\mathcal{C}}^\mu$.

Moreover, if one of the above conditions holds then $R \dot{+} S$ is unitarily equivalent to S :

$$R \dot{+} S = F_\mu(RS^{-1})^* S F_\mu(RS^{-1}) \quad (15)$$

Proof: If $R = 0$ then $RS^{-1} = 0$, $F_\mu(RS^{-1}) = I$. Hence the conditions (i)–(iii) are satisfied and the formula (15) holds. Therefore we may assume that $\ker R = \{0\}$. Then, by virtue of Theorem 2.2: (i) \Leftrightarrow (ii) $\Leftrightarrow ((\text{Sp}(R^{-1}S)) \subset \overline{\mathcal{C}}^\mu)$. Clearly the latter condition is equivalent to (iii) $(RS^{-1} = \mu^2(R^{-1}S)^{-1})$, the transformation $t \mapsto \mu^2 t^{-1}$ maps $\overline{\mathcal{C}}^\mu - \{0\}$ onto itself.

Assume that $\text{Sp}(RS^{-1}) \subset \overline{\mathcal{C}}^\mu$. Then $\text{Sp}((RS^{-1})^*) \subset \overline{\mathcal{C}}^\mu$ and inserting in Theorem 2.2 S^* and R^* instead of R and S we get

$$R^* \dot{+} S^* = F_\mu(RS^{-1})^* S^* F_\mu(RS^{-1}). \quad (16)$$

Clearly $(R \dot{+} S)^* \supset (R^* \dot{+} S^*)$. Remembering that normal operators have no proper normal extensions we see that $(R \dot{+} S)^* = (R^* \dot{+} S^*)$. Now (15) follows immediately from (16).

Q.E.D.

Let Q be a normal operator acting on a Hilbert space H , $E_Q(\cdot)$ be the spectral measure associated with Q and $x \in H$. We denote by $\text{supp}_Q(x)$ the support of the measure $(x | dE_Q(\cdot)x)$. We say that x is of compact Q -support if $\text{supp}_Q(x)$ is bounded (hence compact). Clearly the concept of compact Q -support and that of compact $|Q|$ -support coincide.

The link between (5) and Definition 2.1 is given by the following

Theorem 2.4 Let R and S be normal operators acting on a Hilbert space H . Then the following conditions are equivalent:

1. (R, S) is a (p, q) -commuting pair,
2. The set \mathcal{D} of all vectors with compact R and S support is dense, invariant under the action of R, R^*, S, S^* and

$$\begin{aligned} RSx &= pSRx, \\ RS^*x &= qS^*Rx, \end{aligned} \quad (17)$$

for any $x \in \mathcal{D}$.

3. There exists a dense domain \mathcal{D} consisting of vectors with compact R and S -support invariant under the action of R, R^*, S and S^* such that relations (17) hold for any $x \in \mathcal{D}$.

Proof:

1 \Rightarrow 2: The density of \mathcal{D} follows from the strong commutativity of $|R|$ and $|S|$. To show that \mathcal{D} is R, S, R^* and S^* -invariant it is sufficient to notice that due to the Conditions 3 and 4 of Definition 2.1:

$$\begin{aligned} \text{supp}_{|R|}((\text{Phase } S)x) &\subset \sqrt{p/q} \text{supp}_{|R|}(x), & \text{supp}_{|R|}((\text{Phase } S)^*x) &\subset \sqrt{q/p} \text{supp}_{|R|}(x) \\ \text{supp}_{|S|}((\text{Phase } R)x) &\subset \sqrt{1/pq} \text{supp}_{|S|}(x), & \text{supp}_{|S|}((\text{Phase } R)^*x) &\subset \sqrt{pq} \text{supp}_{|S|}(x) \end{aligned}$$

The trivial verification of (17) is left to the reader.

2 \Rightarrow 3: trivial.

3 \Rightarrow 1: It follows easily from (17) that

$$\begin{aligned} R^* S x &= q^{-1} S R^* x, \\ R^* S^* x &= p^{-1} S^* R^* x, \end{aligned} \tag{18}$$

for any $x \in \mathcal{D}$. Using (17) and (18) one can easily verify that for any non-negative integers k, l, r and s we have

$$(R^*)^r R^{k+r} (S^*)^s S^{l+s} x = (a^k S^*)^s (a^k S)^{l+s} (b^l R^*)^r (b^l R)^{k+r} x,$$

where $a = \sqrt{pq}$ and $b = \sqrt{p/q}$. It shows that

$$(\text{Phase } R)^k f(|R|) (\text{Phase } S)^l g(|S|) x = (\text{Phase } S)^l g(a^k |S|) (\text{Phase } R)^k f(b^l |R|) x,$$

where $f(\lambda)$ is a linear combinations of monomials λ^{k+2r} ($r = 0, 1, 2, \dots$) and $g(\lambda)$ is a linear combinations of monomials λ^{l+2s} ($s = 0, 1, 2, \dots$). Remembering that x is of compact $|R|$ - and $|S|$ -support and that the set of polynomials is dense in $L^2(\nu)$ for any measure ν with compact support we see that

$$(\text{Phase } R)^k f(|R|) (\text{Phase } S)^l g(|S|) = (\text{Phase } S)^l g(a^k |S|) (\text{Phase } R)^k f(b^l |R|),$$

holds for any bounded measurable functions f and g on \mathbf{R}^+ such that $f(0) = g(0) = 0$. Considering the following particular cases

1. $k = l = 0$;
2. $k = l = 1$, $f(\lambda) = g(\lambda) = 1$ for $\lambda > 0$;
3. $k = 1, l = 0$, $f(\lambda) = 1$ for $\lambda > 0$;
4. $k = 0, l = 1$, $g(\lambda) = 1$ for $\lambda > 0$;

we see that Conditions 1, 2, 3 and 4 of Definition 2.1 are satisfied.

Q.E.D.

3 The Hilbert space level

Motivated by Theorem 2.4 we introduce

Definition 3.1 Let $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{\gamma}$ and $\tilde{\delta}$ be closed operators acting on a Hilbert space \tilde{H} . We say that $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ satisfies the relations (2) and (3) if

1. $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{\gamma}$ and $\tilde{\delta}$ are normal,
2. $(\tilde{\alpha}, \tilde{\beta})$ and $(\tilde{\gamma}, \tilde{\delta})$ are (μ, μ) -commuting pairs,
3. $(\tilde{\alpha}, \tilde{\gamma})$ and $(\tilde{\beta}, \tilde{\delta})$ are (μ, μ^{-1}) -commuting pairs,
4. $(\tilde{\beta}, \tilde{\gamma})$ is a $(1, 1)$ -commuting pair,
5. For any $x \in \mathcal{D}(\tilde{\delta}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ we have $\tilde{\delta}x \in \mathcal{D}(\tilde{\alpha})$ and

$$\tilde{\alpha}\tilde{\delta}x - \mu\tilde{\beta}\tilde{\gamma}x = x$$

6. For any $x \in \mathcal{D}(\tilde{\alpha}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ we have $\tilde{\alpha}x \in \mathcal{D}(\tilde{\delta})$ and

$$\tilde{\delta}\tilde{\alpha}x - \mu^{-1}\tilde{\beta}\tilde{\gamma}x = x$$

We have

Theorem 3.2 *Let $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ be a quadruple of closed operators satisfying the relations (2) and (3). Then*

1. $\text{Sp}(\tilde{\beta}\tilde{\gamma}) \subset \overline{\mathbf{C}}^\mu$
2. $\ker \tilde{\delta} = \{0\}$ and $\tilde{\delta}^{-1} = F_\mu(\mu\tilde{\beta}\tilde{\gamma})\tilde{\alpha}F_\mu(\mu\tilde{\beta}\tilde{\gamma})^*$
3. $\tilde{\alpha} = \tilde{\delta}^{-1} + \mu\tilde{\beta}\tilde{\gamma}\tilde{\delta}^{-1}$

Proof: Let $y \in \ker \tilde{\delta} \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$. then $\tilde{\delta}^*y = \tilde{\delta}y = 0$. By virtue of Condition 5 of Definition 3.1: $\tilde{\beta}\tilde{\gamma}y = -\mu^{-1}y$. Consequently $(\tilde{\beta}\tilde{\gamma})^*y = -\mu^{-1}y$ and for any $x \in \mathcal{D}(\tilde{\alpha}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ we get (cf Condition 6 of Definition 3.1)

$$\begin{aligned} (y | x) &= (y | \tilde{\delta}\tilde{\alpha}x - \mu^{-1}\tilde{\beta}\tilde{\gamma}x) \\ &= (\tilde{\delta}^*y | \tilde{\alpha}x) - \mu^{-1}((\tilde{\beta}\tilde{\gamma})^*y | x) = \mu^{-2}(y | x) \end{aligned}$$

Therefore $(y | x) = 0$ and remembering that $\mathcal{D}(\tilde{\alpha}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ is dense (cf Theorem 2.4.3; $(\tilde{\alpha}, \tilde{\beta}\tilde{\gamma})$ is a $(\mu^2, 1)$ -commuting pair) we see that $\ker \tilde{\delta} \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma}) = \{0\}$. On the other hand we know (cf Conditions 2, 3 and 4 of Definition 3.1) that $|\tilde{\delta}|$ and $|\tilde{\beta}\tilde{\gamma}|$ strongly commute. Therefore $\ker \tilde{\delta} \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ is dense in $\ker \tilde{\delta}$ and $\ker \tilde{\delta} = \{0\}$.

It means that $\tilde{\delta}$ is invertible: $\tilde{\delta}^{-1}$ is an unbounded normal operator. By virtue of (13), $(\tilde{\beta}\tilde{\gamma}, \tilde{\delta}) \in \mathcal{C}(\mu^2, 1)$. Using (11) and (10) we get $(\tilde{\delta}^{-1}, \tilde{\beta}\tilde{\gamma}) \in \mathcal{C}(\mu^2, 1)$.

Let $y \in \mathcal{D}(\tilde{\delta}^{-1}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$. Then $y = \tilde{\delta}x$ where $x \in \mathcal{D}(\tilde{\delta}) \cap \mathcal{D}(\tilde{\beta}\tilde{\gamma})$ and using Condition 5 of Definition 3.1 we see that $y \in \mathcal{D}(\tilde{\alpha})$ and $\tilde{\alpha}y = x + \mu\tilde{\beta}\tilde{\gamma}x = \tilde{\delta}^{-1}y + \mu\tilde{\beta}\tilde{\gamma}\tilde{\delta}^{-1}y$. It shows that

$$\tilde{\alpha} \supset \tilde{\delta}^{-1} + \mu\tilde{\beta}\tilde{\gamma}\tilde{\delta}^{-1}.$$

Using Theorem 2.3 with R and S replaced by $\mu\tilde{\beta}\tilde{\gamma}\tilde{\delta}^{-1}$ and $\tilde{\delta}^{-1}$ and remembering that normal operators have no proper normal extensions we get

$$\begin{aligned} \tilde{\alpha} &= \tilde{\delta}^{-1} + \mu\tilde{\beta}\tilde{\gamma}\tilde{\delta}^{-1}, \\ \text{Sp } \mu\tilde{\beta}\tilde{\gamma} &\subset \overline{\mathbf{C}}^\mu, \\ \tilde{\alpha} &= F_\mu(\mu\tilde{\beta}\tilde{\gamma})^*\tilde{\delta}^{-1}F_\mu(\mu\tilde{\beta}\tilde{\gamma}). \end{aligned}$$

Now statements 1, 2, and 3 follow immediately.

Q.E.D.

The reader certainly noticed that Definition 3.1 takes into account only 16 out of 17 relations (2) and (3): the relation “ $\alpha\delta^* = \delta^*\alpha$ ” seems to be forgotten. It turns out however that it may be derived from other relations if we know that δ is invertible: Assuming the nine relations

$$\begin{aligned} \beta\delta &= \mu\delta\beta, & \gamma\delta &= \mu\delta\gamma, & \beta\gamma &= \gamma\beta, \\ \beta\delta^* &= \mu^{-1}\delta^*\beta, & \gamma\delta^* &= \mu\delta^*\gamma, & \beta\gamma^* &= \gamma^*\beta, \\ \beta\beta^* &= \beta^*\beta, & \gamma\gamma^* &= \gamma^*\gamma, & \delta\delta^* &= \delta^*\delta \end{aligned}$$

and setting $\alpha = \delta^{-1} + \mu\beta\gamma\delta^{-1}$ one may easily derive the remaining seven relations (2) and (3).

4 Gauss decomposition

According to (4)

$$\Phi(\delta) = \gamma \otimes \beta + \delta \otimes \delta.$$

Therefore we may expect that $\gamma \otimes \beta + \delta \otimes \delta$ is a normal element or at least has a normal extension. However (cf Theorem 2.2) this is the case only if $\text{Sp}(\gamma\delta^{-1} \otimes \beta\delta^{-1}) \subset \overline{\mathbf{C}}^\mu$. One can easily check that (except the trivial case, when either β or γ vanishes) the latter is equivalent to

$$\begin{aligned} \text{Sp} |\gamma\delta^{-1}| &\subset r\mu^{\mathbf{Z}} \cup \{0\}, \\ \text{Sp} |\beta\delta^{-1}| &\subset r^{-1}\mu^{\mathbf{Z}} \cup \{0\}, \end{aligned}$$

where r is a positive real number. Multiplying β and dividing γ by r (this operation preserves all the relations considered so far) we may assume that $r = 1$.

The above considerations show that the satisfactory theory of the Lorentz group investigated in this paper cannot be based on the algebraic relations (2) and (3) only. We are forced to add conditions of the analytic nature. We met this phenomenon for the first time investigating quantum $E(2)$ -group (cf [6], page 424).

Definition 4.1 *Let $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ be a quadruple of closed operators satisfying the relations (2) and (3). We say that $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ satisfies the spectral condition if*

$$\begin{aligned} \text{Sp} |\tilde{\gamma}\tilde{\delta}^{-1}| &\subset \mu^{\mathbf{Z}} \cup \{0\}, \\ \text{Sp} |\tilde{\beta}\tilde{\delta}^{-1}| &\subset \mu^{\mathbf{Z}} \cup \{0\}. \end{aligned} \tag{19}$$

Let $\overline{\Sigma}_\mu$ be the closure (in \mathbf{R}^2) of the set $\Sigma_\mu = \{(s, \mu^r) : s, r - \frac{s}{2} \in \mathbf{Z}\}$. Then we have

Theorem 4.2 *Let $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ be a quadruple of closed operators on a Hilbert space \tilde{H} satisfying the relations (2) and (3) and the spectral condition (19). Then*

$$\left. \begin{aligned} \tilde{\alpha} &= \tilde{v}\mu^{\tilde{N}/2} + \tilde{n}\tilde{b} \\ \tilde{\beta} &= \tilde{n}\mu^{-\tilde{N}/2} \\ \tilde{\gamma} &= \tilde{v}^*\tilde{b} \\ \tilde{\delta} &= \tilde{v}^*\mu^{-\tilde{N}/2} \end{aligned} \right\} \tag{20}$$

where \tilde{v} , \tilde{n} , \tilde{N} and \tilde{b} are closed operators such that

$$\left. \begin{aligned} (i) & \tilde{v} \text{ is unitary} \\ (ii) & \tilde{n} \text{ is normal and } \text{Sp} \tilde{n} \subset \overline{\mathbf{C}}^\mu \\ (iii) & \tilde{v}\tilde{n}\tilde{v}^* = \mu\tilde{n} \end{aligned} \right\} \tag{21}$$

$$\left. \begin{aligned} (i) & \tilde{N} \text{ is selfadjoint} \\ (ii) & \tilde{b} \text{ is normal} \\ (iii) & \tilde{N} \text{ and } |\tilde{b}| \text{ strongly commute} \\ (iv) & \text{The joint spectrum} \\ & \text{Sp}(\tilde{N}, |\tilde{b}|) \subset \overline{\Sigma}_\mu \\ (v) & \text{On } (\ker \tilde{b})^\perp \\ & (\text{Phase } \tilde{b})^* \tilde{N} (\text{Phase } \tilde{b}) = \tilde{N} + 2I \end{aligned} \right\} \tag{22}$$

$$\left. \begin{array}{l} \text{Any element of the pair } (\tilde{v}, \tilde{n}) \text{ strongly com-} \\ \text{mutes with any element of the pair } (\tilde{N}, \tilde{b}) \end{array} \right\} \quad (23)$$

The elements \tilde{v} , \tilde{n} , \tilde{N} and \tilde{b} are uniquely determined by $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{\gamma}$ and $\tilde{\delta}$.

Conversely if \tilde{v} , \tilde{n} , \tilde{N} and \tilde{b} are closed operators satisfying relations (21), (22) and (23) then the quadruple $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ of operators introduced by (20) satisfies the relations (2) and (3) and the spectral condition (19).

Moreover for any non-degenerate C^* -algebra \tilde{A} acting on \tilde{H}

$$(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta} \eta \tilde{A}) \Leftrightarrow (\tilde{v}, \tilde{n}, \tilde{N}, \tilde{b} \eta \tilde{A})$$

Remark: The best way to memorize the relations (20) is to write them in the matrix form:

$$\begin{pmatrix} \tilde{\alpha} & \tilde{\beta} \\ \tilde{\gamma} & \tilde{\delta} \end{pmatrix} = \begin{pmatrix} \tilde{v} & \tilde{n} \\ 0 & \tilde{v}^* \end{pmatrix} \begin{pmatrix} \mu^{\tilde{N}/2} & 0 \\ \tilde{b} & \mu^{-\tilde{N}/2} \end{pmatrix}. \quad (24)$$

The first matrix on the right hand side is of the form of basic representation of $E_\mu(2)$, whereas the second one is in the same way related to $\hat{E}_\mu(2)$ (cf [6] and [8]). We call (24) the Gauss decomposition (cf [2]).

Proof: We know (cf Theorem 3.2.2) that $\tilde{\delta}$ is invertible. Comparing (19) with Theorem 3.2.1 one can easily show that $\text{Sp } \tilde{\delta}^2 \subset \overline{\mathbf{C}}^\mu$. Therefore the polar decomposition of $\tilde{\delta}$ may be written in the following form

$$\tilde{\delta} = \tilde{v}^* \mu^{-\tilde{N}/2} \quad (25)$$

where \tilde{v} is a unitary and \tilde{N} is a selfadjoint operator with integer spectrum and \tilde{v} strongly commutes with \tilde{N} .

We know that $(\tilde{\gamma}, \tilde{\delta}) \in \mathcal{C}(\mu, \mu)$. Therefore (cf Conditions 2 and 4 of Definition 2.1) $\tilde{v}^* = \text{Phase } \tilde{\delta}$ strongly commutes with $\tilde{\gamma}$ and

$$\tilde{b} = \tilde{v} \tilde{\gamma} \quad (26)$$

is a normal operator strongly commuting with \tilde{v} . Moreover $(\tilde{b}, |\tilde{\delta}|) = (\tilde{b}, \mu^{-\tilde{N}/2}) \in \mathcal{C}(\mu, \mu)$ and relations (iii) and (v) of (22) follow immediately. Remembering that $\text{Sp } \tilde{N} \subset \mathbf{Z}$ and using the first inclusion (19) we get relation (iv) of (22).

By virtue of (9) and (11), $(\tilde{\beta}, |\tilde{\delta}|) \in \mathcal{C}(1, 1)$ and $(\tilde{\beta}, |\tilde{\delta}|^{-1}) \in \mathcal{C}(1, 1)$. Therefore

$$\tilde{n} = \tilde{\beta} |\tilde{\delta}|^{-1} = \tilde{\beta} \mu^{\tilde{N}/2} \quad (27)$$

is a normal operator strongly commuting with \tilde{N} . According to the second inclusion (19) $\text{Sp } \tilde{n} \subset \overline{\mathbf{C}}^\mu$. Combining the relation $(\tilde{\beta}, \tilde{\delta}) \in \mathcal{C}(\mu, \mu^{-1})$ with the obvious one $(|\tilde{\delta}|^{-1}, \tilde{\delta}) \in \mathcal{C}(1, 1)$ we get $(\tilde{n}, \tilde{\delta}) \in \mathcal{C}(\mu, \mu^{-1})$ and (cf (8)) $(\tilde{n}, \tilde{v}^*) \in \mathcal{C}(\mu, \mu^{-1})$ and relation (iii) of (21) follows.

Using in turn (8), (7), (12) and (10) we get $(\tilde{\beta}, \tilde{v}^*) \in \mathcal{C}(\mu, \mu^{-1})$, $(\tilde{\beta}, \tilde{v}) \in \mathcal{C}(\mu^{-1}, \mu)$, $(\tilde{\beta}, \tilde{b}) \in \mathcal{C}(\mu^{-1}, \mu)$ and $(\tilde{b}, \tilde{\beta}) \in \mathcal{C}(\mu, \mu)$. Similarly using (9), (11) and (13) we obtain $(\tilde{\gamma}, |\tilde{\delta}|) \in \mathcal{C}(\mu, \mu)$, $(\tilde{\gamma}, |\tilde{\delta}|^{-1}) \in \mathcal{C}(\mu^{-1}, \mu^{-1})$ and $(\tilde{b}, |\tilde{\delta}|^{-1}) \in \mathcal{C}(\mu^{-1}, \mu^{-1})$. Using now (12) we get $(\tilde{b}, \tilde{n}) \in \mathcal{C}(1, 1)$. It shows that \tilde{b} and \tilde{n} strongly commute.

The last three formulae of (20) follow immediately from (25), (26) and (27). The first one follows from Theorem 3.2.3. This way the first part of the theorem is proved.

Assume now the relations (21), (22) and (23) and consider operators $\tilde{\alpha}$, $\tilde{\beta}$, $\tilde{\gamma}$ and $\tilde{\delta}$ introduced by (20). The only nontrivial points of the second part of the theorem are the normality of $\tilde{\alpha}$ and the commutation relations involving Phase $\tilde{\alpha}$ and $|\tilde{\alpha}|$. Let $S = \tilde{v}\mu^{\tilde{N}/2} = \tilde{\delta}^{-1}$ and $R = \tilde{n}\tilde{b} = \mu\tilde{\beta}\tilde{\gamma}S$. Then $(R, S) \in \mathcal{C}(\mu^{-2}, 1)$, $\text{Sp}(RS^{-1}) \subset \overline{\mathbf{C}}^\mu$ and $\tilde{\alpha} = S\dot{+}R$. By virtue of Theorem 2.3, $\tilde{\alpha}$ is normal and

$$\tilde{\alpha} = F_\mu(\mu\tilde{\beta}\tilde{\gamma})^*\tilde{v}\mu^{\tilde{N}/2}F_\mu(\mu\tilde{\beta}\tilde{\gamma}) \quad (28)$$

Using the last formula one can easily check that $(\tilde{\alpha}, \tilde{\beta}) \in \mathcal{C}(\mu, \mu)$ and $(\tilde{\alpha}, \tilde{\gamma}) \in \mathcal{C}(\mu, \mu^{-1})$. This ends the proof of the second part of the theorem.

If $\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta}\eta\tilde{A}$ then (cf Theorem 3.2.2) $\tilde{\delta}^{-1}\eta\tilde{A}$. Therefore $\tilde{v} = (\text{Phase } \tilde{\delta})^*$ and $\tilde{N} = -2\log_\mu(|\tilde{\delta}|)$ are affiliated with \tilde{A} and by virtue of (26) and (27): $\tilde{n}, \tilde{b}\eta\tilde{A}$.

Conversely if $\tilde{v}, \tilde{n}, \tilde{N}, \tilde{b}\eta\tilde{A}$ then according to the last three relations (20) $\tilde{\beta}, \tilde{\gamma}, \tilde{\delta}\eta\tilde{A}$. Using (28) we get $\tilde{\alpha}\eta\tilde{A}$.

Q.E.D.

5 C^* -algebra level

In this Section we construct the universal C^* -algebra A related with (2), (3) and (19). The algebra A plays the role of function algebra of G_μ . The comultiplication $\Phi \in \text{Mor}(A, A \otimes A)$ endowing G_μ with the group structure will be introduced.

Comparing Theorem 4.2 with the content of [8] we discover that the relations (21) and (22) coincide with the relations imposed on “generators” v, n and N, b of the function algebras A and B of $E_\mu(2)$ and $\hat{E}_\mu(2)$ quantum groups. In what follows we write A_1 and A_2 instead of A and B .

The algebra A_1 is described in details in Section 1 of [8]. We know that there exists comultiplication $\Phi_1 \in \text{Mor}(A_1, A_1 \otimes A_1)$ (denoted by Φ in [8]) such that

$$\left. \begin{aligned} \Phi_1(v) &= v \otimes v \\ \Phi_1(n) &= v \otimes n \dot{+} n \otimes v^* \end{aligned} \right\} \quad (29)$$

The algebra A_2 is introduced in Section 3 of [8]. It is known that there exists comultiplication $\Phi_2 \in \text{Mor}(A_2, A_2 \otimes A_2)$ (denoted by $\hat{\Phi}$ in [8]) such that

$$\left. \begin{aligned} \Phi_2(N) &= N \otimes I \dot{+} I \otimes N \\ \Phi_2(b) &= b \otimes \mu^{N/2} \dot{+} \mu^{-N/2} \otimes b. \end{aligned} \right\} \quad (30)$$

Let

$$w = F_\mu(\mu^{N/2}b \otimes vn)(I \otimes v)^{N \otimes I} \quad (31)$$

Then w is a unitary element of $M(A_2 \otimes A_1)$ and (cf Theorems 3.2 and 4.1 of [8])

$$\left. \begin{aligned} (\text{id} \otimes \Phi_1)w &= w_{12}w_{13} \\ (\Phi_2 \otimes \text{id})w &= w_{23}w_{13}. \end{aligned} \right\} \quad (32)$$

Let $A = A_1 \otimes A_2$ and (cf (20))

$$\left. \begin{aligned} \alpha &= v \otimes \mu^{N/2} \dot{+} n \otimes b \\ \beta &= n \otimes \mu^{-N/2} \\ \gamma &= v^* \otimes \tilde{b} \\ \delta &= v^* \otimes \mu^{-N/2} \end{aligned} \right\} \quad (33)$$

Then $\alpha, \beta, \gamma, \delta \in \eta A$. Combining the Theorems 1.1 and 3.1 of [8] with Theorem 4.2 we get

Theorem 5.1 1. Let π be a representation of A in a Hilbert space \widetilde{H} and

$$\left. \begin{aligned} \tilde{\alpha} &= \pi(\alpha), & \tilde{\beta} &= \pi(\beta), \\ \tilde{\gamma} &= \pi(\gamma), & \tilde{\delta} &= \pi(\delta). \end{aligned} \right\} \quad (34)$$

Then the quadruple $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ satisfies the relations (2), (3) and the spectral condition (19).

2. Any quadruple $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$ of closed operators on a Hilbert space \widetilde{H} satisfying the relations (2), (3) and (19) is of the form (34), where π is a representation of A in \widetilde{H} . The representation π is uniquely determined by $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta})$.

3. Let $\pi, \widetilde{H}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta}$ be as in Statement 1 and $\tilde{A} \in C^*(\widetilde{H})$. Then

$$(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta}) \eta \tilde{A} \iff (\pi \in \text{Mor}(A, \tilde{A})).$$

This theorem shows that the algebra A is the one corresponding to the commutation relation (2), (3) and (19). Copying formulae (4.9) and (4.16) of [5] we set:

$$\Phi = (\text{id} \otimes \sigma \otimes \text{id})(\Phi_1 \otimes \Phi_2),$$

where $\sigma \in \text{Mor}(A_1 \otimes A_2, A_2 \otimes A_1)$ is introduced by the formula

$$\sigma(a \otimes x) = w(x \otimes a)w^*$$

for any $a \in A_1$ and $x \in A_2$. Clearly $\Phi \in \text{Mor}(A, A \otimes A)$. Repeating the computations of [5] (pp 406–408) one can easily show that Φ is coassociative. We shall prove

Theorem 5.2 The action of Φ on the distinguished elements $\alpha, \beta, \gamma, \delta \in \eta A$ is described by the standard formulae:

$$\begin{aligned} \Phi(\alpha) &= \alpha \otimes \alpha \dot{+} \beta \otimes \gamma \\ \Phi(\beta) &= \alpha \otimes \beta \dot{+} \beta \otimes \delta \\ \Phi(\gamma) &= \gamma \otimes \alpha \dot{+} \delta \otimes \gamma \\ \Phi(\delta) &= \gamma \otimes \beta \dot{+} \delta \otimes \delta \end{aligned}$$

Proof: By virtue of the second part of Theorem 4.4 of [5] it is sufficient to show that

$$(\text{id} \otimes \sigma) \left[\left(\begin{array}{c} v, n \\ 0, v^* \end{array} \right) \oplus \left(\begin{array}{c} \mu^{N/2}, 0 \\ b, \mu^{-N/2} \end{array} \right) \right] = \left(\begin{array}{c} \mu^{N/2}, 0 \\ b, \mu^{-N/2} \end{array} \right) \oplus \left(\begin{array}{c} v, n \\ 0, v^* \end{array} \right)$$

or in the expanded form

$$\begin{aligned} \sigma(v \otimes \mu^{N/2} \dot{+} n \otimes b) &= \mu^{N/2} \otimes v \\ \sigma(n \otimes \mu^{-N/2}) &= \mu^{N/2} \otimes n \\ \sigma(v^* \otimes b) &= b \otimes v \\ \sigma(v^* \otimes \mu^{-N/2}) &= b \otimes n \dot{+} \mu^{-N/2} \otimes v^* \end{aligned}$$

These relations are equivalent to

$$\mu^{N/2} \otimes v \dot{+} b \otimes n = w^*(\mu^{N/2} \otimes v)w \quad (35)$$

$$w(\mu^{-N/2} \otimes n)w^* = \mu^{N/2} \otimes n \quad (36)$$

$$w(b \otimes v^*)w^* = b \otimes v \quad (37)$$

$$w(\mu^{-N/2} \otimes v^*)w^* = b \otimes n \dot{+} \mu^{-N/2} \otimes v^* \quad (38)$$

To prove (36) and (37) it is sufficient to notice that $(I \otimes v)^{N \otimes I}(\mu^{-N/2} \otimes n)(I \otimes v)^{-N \otimes I} = \mu^{N/2} \otimes n$ and $(I \otimes v)^{N \otimes I}(b \otimes v^*)(I \otimes v)^{-N \otimes I} = b \otimes v$, whereas $\mu^{N/2} b \otimes v n$ strongly commutes with $\mu^{N/2} \otimes n$ and $b \otimes v$.

To prove (38) it is sufficient to notice that $(I \otimes v)^{N \otimes I}$ commutes with $\mu^{-N/2} \otimes v^*$ and then use the formula (14) with R and S replaced by $\mu^{-N/2} \otimes v^*$ and $b \otimes n$ resp.

Similarly to prove (35) it is sufficient to use the formula (15) with R and S replaced by $\mu^{N-2I} b \otimes v^2 n$ and $\mu^{N/2} \otimes v$ and then use the equality: $(I \otimes v)^{-N \otimes I}(\mu^{N-2I} b \otimes v^2 n)(I \otimes v)^{N \otimes I} = b \otimes n$.

Q.E.D.

Acknowledgment

One of the authors (SLW) is grateful to Professor J. Fröhlich for his kind hospitality during the authors visit to the ETH Zürich.

References

- [1] S. Baaj and P. Jungl: Théorie bivariant de Kasparow et opérateur non bornés dans les C^* -modules hilbertiens, C. R. Acad. Sci. Paris, Série I, 296 (1983) 875–878, see also S. Baaj: Multiplicateur non bornés, Thèse 3^{ème} Cycle, Université Paris VI, 11 Decembre 1980
- [2] A.O. Barut and R. Rączka: Theory of group representations and applications. Warszawa PWN – Polish Scientific Publishers 1977

- [3] V.G. Drinfel'd: "Quantum groups", in Proceedings of the International Congress of Mathematicians, Berkeley 1986, American Mathematical Society, 1987,798–820.
- [4] L. Faddeev, N. Reshetikhin and L. Takhtajan: Quantum Groups, in: Braid group, knot theory and statistical mechanics, Ed. C. N. Yang, M. L. Ge, World scientific, 1989, pp. 97-110.
- [5] P. Podleś and S.L. Woronowicz: Quantum deformation of Lorentz group. Commun. Math. Phys. 130, 381–431 (1990)
- [6] S.L. Woronowicz: Unbounded elements affiliated with C^* -algebras and non-compact quantum groups. Commun. Math. Phys. 136, 399–432 (1991)
- [7] S.L. Woronowicz: Operator equalities related to quantum $E(2)$ group, submitted to Commun. Math. Phys.
- [8] S.L. Woronowicz: Quantum $E(2)$ -group and its Pontryagin dual, submitted to Lett. Math. Phys.
- [9] S.L. Woronowicz and S. Zakrzewski: Quantum deformations of Lorentz group. Hopf $*$ -algebra level. (in preparation)