

IPPP/12/10

DCPT/12/20

March 10, 2014

SUSY_FLAVOR v2.1: a computational tool for FCNC and CP-violating processes in the MSSM

A. Crivellin^a, J. Rosiek^b, P. Chankowski^b, A. Dedes^c, S. Jäger^d P. Tanedo^e

^a*Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics,
University of Bern, CH-3012 Bern, Switzerland*

^b*Institute of Theoretical Physics, University of Warsaw, 00-681 Warsaw, Poland*

^c*Division of Theoretical Physics, University of Ioannina, GR 45110, Greece*

^d*Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, UK*

^e*Institute for High Energy Phenomenology, Newman Laboratory of Elementary Particle Physics, Cornell University, Ithaca, NY 14853, USA*

Abstract

We present SUSY_FLAVOR version 2 — a Fortran 77 program that calculates low-energy flavor observables in the general R -parity conserving MSSM. For a set of MSSM parameters as input, the code gives predictions for:

1. Electric dipole moments of the leptons and the neutron.
2. Anomalous magnetic moments (i.e. $g - 2$) of the leptons.
3. Radiative lepton decays ($\mu \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma, e\gamma$).
4. Rare Kaon decays ($K_L^0 \rightarrow \pi^0 \bar{\nu}\nu$ and $K^+ \rightarrow \pi^+ \bar{\nu}\nu$).
5. Leptonic B decays ($B_{s,d} \rightarrow l^+l^-$, $B \rightarrow \tau\nu$ and $B \rightarrow D\tau\nu$).
6. Radiative B decays ($B \rightarrow \bar{X}_s\gamma$).
7. $\Delta F = 2$ processes ($\bar{K}^0 - K^0$, $\bar{D} - D$, $\bar{B}_d - B_d$ and $\bar{B}_s - B_s$ mixing).

Comparing to SUSY_FLAVOR v1, where the matching conditions were calculated at one-loop level, SUSY_FLAVOR v2 performs the resummation of all chirally enhanced corrections, i.e. takes into account the enhanced effects from $\tan\beta$ and/or large trilinear soft mixing terms to all orders in perturbation theory. Also, in SUSY_FLAVOR v2 new routines calculating of $B \rightarrow (D)\tau\nu$, $g - 2$, radiative lepton decays and $Br(l \rightarrow l'\gamma)$ were added. All calculations are done using exact diagonalization of the sfermion mass matrices. In SUSY_FLAVOR v2.1 parameter initialization in SLHA2 formats has been generalized and simplified. The program can be obtained from http://www.fuw.edu.pl/susy_flavor.

1 Introduction

Flavor physics was in the recent years one of the most active and fastest developing fields in the high energy physics. Numerous new experiments, spanning a wide energy range from neutrino mass measurements to hard proton scattering at LHC collider, helped to improve significantly the accuracy of various measurements related to flavor-observables. Almost all such experiments reported result which are in agreement with the Standard Model (SM) predictions, with a few exception where small observed deviations still require further confirmation (like e.g. $g - 2$ muon magnetic moment anomaly [1, 2]).

The extensive set of measurements available for rare decays puts strong constraints on the flavor structure of physics beyond the Standard Model. In particular, it imposes stringent limits on the flavor- and CP- violating parameters of the Minimal Supersymmetric Standard Model (MSSM) [3], where the flavor changing neutral currents (FCNCs) originate, in addition to the CKM induced FCNCs, from the fact that one cannot (in general) simultaneously diagonalize the mass matrices of fermions and sfermions. Such a misalignment leads to FCNCs which can involve the strong coupling constant and which do not necessarily respect the hierarchy of the CKM matrix. Moreover, many of the MSSM parameters can take complex values and are potential sources of CP violation. Thus supersymmetric contributions to flavor and/or CP-violating processes can, in principle, exceed the SM predictions by orders of magnitude. The apparent absence of such big effects leads to the constraint that MSSM couplings which may generate FCNCs and CP violation are actually strongly suppressed. The difficulty to explain this suppression is known as the “SUSY flavor problem” and the “SUSY CP problem”. Even if one assumes the so-called Minimal Flavor Violation (MFV) hypothesis [4] which requires that *all* FCNC effects originate from the Yukawa couplings of the superpotential, supersymmetric contributions to various flavor and CP-violating amplitudes can still be of comparable (or sometimes even much larger, like in the case of the electron and neutron EDMs or $B_s \rightarrow \mu^+ \mu^-$) size as the corresponding SM contributions.

As the accuracy of the flavor experiments constantly improves, it is important to have an universal computational tool which helps to compare new data with the predictions of the MSSM. Developing such a tool is a non-trivial task requiring extensive and often tedious calculations. Numerous analyses have been published in the literature, but because of the complexity of the problem, they usually consider only a few rare decays simultaneously. Furthermore, many analyses done for general flavor violation in the MSSM use the mass insertion approximation for the soft terms (MIA) (see e.g. [5, 6]) which simplifies the calculations but does not produce correct results if flavor violation (and/or chirality violation) in the sfermion sector becomes large.

In a series of papers published since 1997 [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17], many supersymmetric FCNC and CP-violating observables were analyzed within the setup of the most general R -parity conserving MSSM using exact diagonalization of the sfermion

mass matrices. A FORTRAN computer programs based on the common set of Feynman rules of Ref. [18] were developed for each process (using also parts of code written for earlier papers on MSSM Higgs physics [19]) and, after collecting them together, published as **SUSY_FLAVOR v1** [20].

SUSY_FLAVOR v1 was able to calculate only the 1-loop supersymmetric virtual corrections, whereas, as widely discussed in the literature [21, 22, 23, 24, 25, 26, 27, 28, 13, 29, 30, 31, 32, 33, 34, 35], in the regime of large $\tan\beta$ or large trilinear A -terms it is important to take into account the chirally enhanced corrections to all orders of perturbation theory.

Chiral enhancement is always related to fermion-Higgs couplings. Because these couplings have mass-dimension 4, the corresponding corrections have the property that they do not vanish in the decoupling limit ($m_{\text{SUSY}} \rightarrow \infty$) but rather converge to a constant. This in turn also means that the flavor-changing neutral Higgs couplings, which are induced by chirally-enhanced SUSY corrections, are still relevant for heavy SUSY particles. Thus, especially for observables which are sensitive to Higgs contributions (like for example $B_{s,d} \rightarrow \mu^+\mu^-$ or $B_{d,s}$ mixing), the consistent resummation and inclusion of all chirally enhanced corrections is very important.

In Ref. [17] such resummation was performed in the most general MSSM taking into account all possible sources of chiral enhancement; in the decoupling limit ($m_{\text{SUSY}} \gg v_1, v_2$ analytical formulae has been given. Results of ref. [17] has been implemented in **SUSY_FLAVOR v2**. The consistent treatment of all chirally enhanced effects in the general MSSM (and the corresponding threshold corrections), including correct calculation of neutral Higgs penguins in such scenario, is a unique feature of **SUSY_FLAVOR v2** not shared at the moment by other publicly available programs calculating rare decays in the supersymmetric models. In addition we added in new version several new routines calculating additional physical observables (mostly in lepton sector). All processes which can be calculated by **SUSY_FLAVOR v2** are listed in Table 1.

Several other programs allowing to analyze various aspects of the MSSM flavor phenomenology have been published. The most relevant to **SUSY_FLAVOR** are: **CPsuperH** [36], **SusyBSG** [37], **SPheno** [38], **SuperIso** [39] and **SUSEFLAV** [40]. **SusyBSG** is dedicated to high-precision predictions for $B \rightarrow X_s\gamma$ while **CPsuperH** and **SuperIso** calculate processes similar to the ones computed by **SUSY_FLAVOR**. However, these existing codes are restricted to the Minimal Flavor Violation scenario, whereas **SUSY_FLAVOR** can simultaneously calculate the set of rare decays listed in Table 1 without any (apart from the R -parity conservation) restrictions on the choice of MSSM parameters. Other publicly available codes that are relevant to **SUSY_FLAVOR** (which can e.g. calculate the MSSM soft parameters used as input to **SUSY_FLAVOR**, or for the same set of input parameters calculate non-FCNC related observables) are **FeynHiggs**[41], **SoftSUSY**[42], **SuSpect**[43], **MicrOMEGAs**[44], **DarkSUSY**[45] and **NMHDECAY**[46].

In summary, the basic features of **SUSY_FLAVOR v2** are:

- The program utilizes the most general R -parity conserving Lagrangian for the MSSM. In addition to the standard soft breaking terms, it can accommodate for additional non-holomorphic trilinear soft-SUSY breaking terms,

$$A_l'^{IJ} H_i^{2\star} L_i^I R^J + A_d'^{IJ} H_i^{2\star} Q_i^I D^J + A_u'^{IJ} H_i^{1\star} Q_i^I U^J + \text{H.c.} , \quad (1)$$

that do not appear in the minimal supergravity scenario but are present in the most general softly broken supersymmetric effective Lagrangian [47]. These non-holomorphic terms can give rise to sizable effects in Higgs-fermion couplings.

- There is, in principle, no limit on the size of flavor violating parameters because the calculation does not rely on the MIA expansion. However, if the off-diagonal elements are larger than the diagonal ones, imaginary sfermion masses would be induced. Complex “mass insertions” of the form

$$\delta_{QXY}^{IJ} = \frac{(M_Q^2)^{IJ}_{XY}}{\sqrt{(M_Q^2)^{II}_{XX}(M_Q^2)^{JJ}_{YY}}} , \quad (2)$$

(I, J denote quark flavors, X, Y denote superfield chirality, and Q indicates either the up or down quark superfield sector, similarly for slepton superfields) are taken as inputs, but they only serve to conveniently parametrize the sfermion mass matrices. **SUSY_FLAVOR** numerically calculates the exact tree-level spectrum and mixing matrices, which are later used in loop calculations.

- After calculating SUSY spectrum, **SUSY_FLAVOR** performs the resummation of the chirally enhanced corrections (following the systematic approach of ref. [17]), arising in the regime of large $\tan\beta$ and/or large trilinear soft sfermion mixing. The values of the Yukawa couplings and the CKM matrix elements of the superpotential are calculated (by taking into account the threshold corrections) and are then used for the calculations of the SUSY loop contributions to flavor observables. These chirally enhanced corrections also lead to flavor-changing neutral Higgs couplings and corrections to charged Higgs vertices which are implemented as well in the calculation of the amplitudes.
- As an intermediate step parton-level form factors for quark and lepton 2-, 3- and 4-point Green functions are calculated. They are later dressed in hadronic matrix elements (see Table 3 in Sec. 3) to obtain predictions for the physical quantities listed in Table 1. The set of Green’s functions computed by **SUSY_FLAVOR** as intermediate “building blocks” is quite universal and can be used for a calculation of various processes not yet implemented in **SUSY_FLAVOR**.

This article is organized as follows. In Sec. 2 we discuss the conventions used for the MSSM parameters (more explicit description can be found in the manual of **SUSY_FLAVOR** v1 [20]). Sec. 3 describes the internal structure of **SUSY_FLAVOR**, the most important steps

Observable	Experiment
$\Delta F = 0$	
$\frac{1}{2}(g - 2)_e$	$(1159652188.4 \pm 4.3) \times 10^{-12}$ [48]
$\frac{1}{2}(g - 2)_\mu$	$(11659208.7 \pm 8.7) \times 10^{-10}$ [2]
$\frac{1}{2}(g - 2)_\tau$	$< 1.1 \times 10^{-3}$ [49]
$ d_e (\text{ecm})$	$< 1.6 \times 10^{-27}$ [50]
$ d_\mu (\text{ecm})$	$< 2.8 \times 10^{-19}$ [51]
$ d_\tau (\text{ecm})$	$< 1.1 \times 10^{-17}$ [52]
$ d_n (\text{ecm})$	$< 2.9 \times 10^{-26}$ [53]
$\Delta F = 1$	
$\text{Br}(\mu \rightarrow e\gamma)$	$< 2.8 \times 10^{-11}$ [54]
$\text{Br}(\tau \rightarrow e\gamma)$	$< 3.3 \times 10^{-8}$ [55]
$\text{Br}(\tau \rightarrow \mu\gamma)$	$< 4.4 \times 10^{-8}$ [55]
$\text{Br}(K_L \rightarrow \pi^0\nu\nu)$	$< 6.7 \times 10^{-8}$ [56]
$\text{Br}(K^+ \rightarrow \pi^+\nu\nu)$	$17.3_{-10.5}^{+11.5} \times 10^{-11}$ [57]
$\text{Br}(B_d \rightarrow ee)$	$< 1.13 \times 10^{-7}$ [58]
$\text{Br}(B_d \rightarrow \mu\mu)$	$< 1.8 \times 10^{-8}$ [59]
$\text{Br}(B_d \rightarrow \tau\tau)$	$< 4.1 \times 10^{-3}$ [60]
$\text{Br}(B_s \rightarrow ee)$	$< 7.0 \times 10^{-5}$ [61]
$\text{Br}(B_s \rightarrow \mu\mu)$	$< 1.08 \times 10^{-8}$ [59]
$\text{Br}(B_s \rightarrow \tau\tau)$	---
$\text{Br}(B_s \rightarrow \mu e)$	$< 2.0 \times 10^{-7}$ [52]
$\text{Br}(B_s \rightarrow \tau e)$	$< 2.8 \times 10^{-5}$ [52]
$\text{Br}(B_s \rightarrow \mu\tau)$	$< 2.2 \times 10^{-5}$ [52]
$\text{Br}(B^+ \rightarrow \tau^+\nu)$	$(1.65 \pm 0.34) \times 10^{-4}$ [52]
$\text{Br}(B_d \rightarrow D\tau\nu)/\text{Br}(B_d \rightarrow Dl\nu)$	$(0.407 \pm 0.12 \pm 0.049)$ [62]
$\text{Br}(B \rightarrow X_s\gamma)$	$(3.52 \pm 0.25) \times 10^{-4}$ [63]
$\Delta F = 2$	
$ \epsilon_K $	$(2.229 \pm 0.010) \times 10^{-3}$ [52]
ΔM_K	$(5.292 \pm 0.009) \times 10^{-3} \text{ ps}^{-1}$ [52]
ΔM_D	$(2.37_{-0.71}^{+0.66}) \times 10^{-2} \text{ ps}^{-1}$ [52]
ΔM_{B_d}	$(0.507 \pm 0.005) \text{ ps}^{-1}$ [63]
ΔM_{B_s}	$(17.77 \pm 0.12) \text{ ps}^{-1}$ [64]

Table 1: List of observables calculated by **SUSY_FLAVOR v2** and their currently measured values or 95% CL. bounds (except for $\text{Br}(B_d \rightarrow e^+e^-)$ and $\text{Br}(B_d \rightarrow \tau^+\tau^-)$ for which the 90% C.L. bounds are given).

of the calculations and the file structure of the library. In Sec. 4 we define the input parameters and present the initialization sequence for **SUSY_FLAVOR**. Sec. 5 discusses how the resummation of the chirally enhanced corrections to all orders of perturbation theory is performed. Routines for calculating the FCNC and CPV observables collected in Table 1 are described in Sec. 6. In Sec. 7 the output format for the quantities calculated by **SUSY_FLAVOR** is presented. We conclude in Sec. 8 with a summary of the presentation. Appendix A contains brief instructions on how to install and run the **SUSY_FLAVOR** package. In appendices B and C we provide templates for initializing **SUSY_FLAVOR** from within the program and using an external file in the SLHA2 format [65], respectively. Both of these templates produce the set of test results listed in Appendix D.

SUSY_FLAVOR can be downloaded from the following address¹:

http://www.fuw.edu.pl/susy_flavor

2 Lagrangian and conventions

SUSY_FLAVOR is capable of calculating physical observables within the most general R -parity conserving MSSM, with one exception: currently it assumes massless neutrinos (and no right neutrino and right sneutrino fields in the Lagrangian [66]), so the PMNS mixing matrix does not appear in any lepton and slepton couplings. Neutrino flavor mixing and the PMNS matrix should be taken into account once new experiments are able to identify the flavor of the neutrinos produced in rare decays, but at present this is not experimentally feasible. Still, over 100 Lagrangian parameters are taken as input to **SUSY_FLAVOR** and can be initialized independently.

SUSY_FLAVOR has been in development since 1996, long before the Les Houches Accord [67] (SLHA), followed in 2008 by SLHA2 [65], for common MSSM conventions was agreed on. By the time SLHA2 became a commonly accepted standard, it was no longer feasible to change the internal **SUSY_FLAVOR** structure as it would require careful checking and rewriting of thousands of lines of a complicated code. Thus, the internal routines of the library follow the conventions for the MSSM Lagrangian and Feynman rules given in the earlier paper [18]. However, by default **SUSY_FLAVOR** can be initialized with a SLHA2 compatible set of parameters and all necessary translations are done in a way invisible for an external user.

Actually, the choice of convention for the input parameters of **SUSY_FLAVOR** is a user-defined option. If required, parameters can be also initialized directly following the [18] conventions. The choice between SLHA2 and ref. [18] can be made by setting the relevant control variable, as described in Sec. 4.1. In Table 2 we summarize the differences between

¹For an additional information, bug reports or any other questions related to the code please contact **SUSY_FLAVOR** maintainer at the address *janusz.rosiek@fuw.edu.pl*

the conventions of the extended SLHA2 [65] and those of [18]. These differences are quite minor and translation can be done by changing few signs and/or transposing matrices in the soft SUSY breaking sector.

SLHA2 [65]	Ref. [18]
$\hat{T}_U, \hat{T}_D, \hat{T}_E$	$-A_u^T, +A_d^T, +A_l^T$
$\hat{m}_Q^2, \hat{m}_{\tilde{L}}^2$	m_Q^2, m_L^2
$\hat{m}_{\tilde{u}}^2, \hat{m}_{\tilde{d}}^2, \hat{m}_{\tilde{l}}^2$	$(m_U^2)^T, (m_D^2)^T, (m_E^2)^T$
$\mathcal{M}_{\tilde{u}}^2, \mathcal{M}_{\tilde{d}}^2$	$(\mathcal{M}_U^2)^T, (\mathcal{M}_D^2)^T$

Table 2: Comparison of SLHA2 [65] and Ref. [18] conventions.

One should note that in **SUSY_FLAVOR** one can also use non-standard trilinear scalar couplings, involving the complex conjugated Higgs fields (sometimes called “non-analytic” or “non-holomorphic” A -terms). In the notation of [18] they read as:

$$A_l'^{IJ} H_i^{2\star} L_i^I E^J + A_d'^{IJ} H_i^{2\star} Q_i^I D^J + A_u'^{IJ} H_i^{1\star} Q_i^I U^J + \text{h.c.} \quad (3)$$

Usually these couplings are not considered as they are not generated in standard SUSY breaking models. However, for completeness they are included in **SUSY_FLAVOR** and by default initialized to zero. Users may decide to set them to some non-trivial values in order to check their impact on rare decay phenomenology (loop corrections non-holomorphic A -terms may lead to large flavor-changing neutral Higgs couplings).

In general, the parameter μ , the soft-SUSY breaking Higgs-mass term m_{12}^2 , the gaugino mass parameters $M_{1,2,3}$, the soft sfermion mass matrices and the trilinear soft couplings may be complex. Global rephasing of all fermion fields of the theory and of one of the Higgs multiplets can render two of these parameters real [7]. We choose them to be the gluino mass M_3 and the Higgs mass term m_{12}^2 . The latter choice keeps the Higgs vacuum expectation values (VEV) and, therefore, the parameter $\tan \beta$ real at tree level.

3 Structure of the code

Calculations in **SUSY_FLAVOR** take the following steps:

- 1. Parameter initialization.** This is described in details in Sec. 4. Users can adjust the basic Standard Model parameters according to latest experimental data and initialize all (or the chosen subset of) supersymmetric soft masses and couplings and Higgs sector parameters. The supersymmetric input parameters for the **SUSY_FLAVOR** must be given at the SUSY scale and program offers no internal routines for evolving them to other scales. At this step also various QCD- and hadronic-related quantities, like e.g. hadronic matrix element values, can be adjusted.

2. Calculation of the physical masses and the mixing angles. After setting the input parameters, **SUSY_FLAVOR** calculates the eigenvalues of the mass matrices of all MSSM particles and their mixing matrices at tree level. Diagonalization is done numerically without any approximations.

3. Resummation of the chirally enhanced effects. In the regime of large $\tan\beta$ and/or large trilinear SUSY breaking terms, large chirally enhanced corrections to Yukawa couplings and CKM matrix elements arise. **SUSY_FLAVOR v2** can perform resummation of these corrections to all orders of perturbation theory. After calculating threshold corrections, the Yukawa couplings and CKM elements of the superpotential (i.e. the “bare” parameters) are determined. Using these quantities the chirally enhanced effects are calculated and absorbed into effective Higgs-fermion and fermion-sfermion-gaugino(higgsino) vertices. Using these vertices in the calculation of flavor observables, all chirally enhanced corrections are automatically taken into account. The level of resummation (no resummation, approximate analytical resummation in the decoupling limit, iterative numerical resummation) is a user defined option.

4. Calculation of the Wilson coefficients at the SUSY scale. The one-loop Wilson coefficients of the effective operators required for a given process are calculated using the sfermion mixing matrices and the physical masses as input. Again, the formulae used in the code do not rely on any approximations, such as the MIA expansion. In the current version, **SUSY_FLAVOR** calculates Wilson coefficients generated by the diagrams listed in Table 3. All Wilson coefficients are calculated at the energy scale assumed to be the average mass of SUSY particles contributing to a given process or the top quark scale.

Box	Penguin	Self energy
$dddd$	$Z\bar{d}d, \gamma\bar{d}d, g\bar{d}d$	d -quark
$uuuu$	$H_i^0\bar{d}d, A_i^0\bar{d}d$	u -quark
$ddll$	$H_i^0\bar{u}u, A_i^0\bar{u}u$	charged lepton l
$dd\nu\nu$	$\gamma\bar{l}l$	

Table 3: One loop parton level diagrams implemented in **SUSY_FLAVOR**.

It is important to stress that routines of **SUSY_FLAVOR** calculating form factors accept fermion generation indices as input parameters. Thus in Table 3 d and u , l and ν denote quarks or leptons of *any* generation. Hence, the actual number of amplitudes which can be calculated using combinations of these form factors is much larger than used by the rare decay rates currently implemented fully in **SUSY_FLAVOR**, opening possibility for further developments of the library.

5. Strong corrections. In the final step **SUSY_FLAVOR** performs (when necessary) the QCD evolution of the Wilson coefficients from the high scale (SUSY or top quark

mass scale) to the low energy scale appropriate for a given decay, calculates the relevant hadronic matrix elements, and returns predictions for physical quantities. The formulae for QCD and hadronic corrections are primarily based on calculations performed in the SM and supplemented, when necessary, with contributions from non-standard operators which usually are neglected in the SM, because they are suppressed by powers of the light quark Yukawa couplings. This part of **SUSY_FLAVOR** is based on analyses published by other authors, whereas points 1-4 are implemented using our own calculations. The accuracy of strong corrections differ from process to process, from negligible or small (leptonic EDM, “gold-plated” decay modes $K \rightarrow \pi\bar{\nu}\nu$ [75]) to order of magnitude uncertainties (unknown long distance contributions to Δm_K or Δm_D). Even in the case of large QCD uncertainties, the result of the calculation performed by **SUSY_FLAVOR** can be of some use. Flavor violation in the sfermion sector can lead to huge modifications of many observables, sometimes by several orders of magnitude, so that comparison with experimental data can help to constrain the soft flavor-violating terms even if the strong corrections are not very well known.

In Table 4 we list the files included in **SUSY_FLAVOR** library with a brief description of their content and purpose. Most of the 2-, 3- and 4-point Green functions are calculated for vanishing external momenta (exception are up-quark self energies and Higgs-up quark 3-point functions where Higgs boson and top quark masses are not small enough to be neglected). As mentioned before, by “*u* quark” and “*d* quark” we mean all generations of quarks. In addition to files listed in Table 4, the library contains the master driver files `susy_flavor_file.f` and `susy_flavor_prog.f` which illustrate the proper initialization sequence for **SUSY_FLAVOR** parameters and produce a sample of results for the implemented observables.

4 Parameter initialization in **SUSY_FLAVOR**

Apart from initialization routines used by **SUSY_FLAVOR** and their arguments we list here the FORTRAN common blocks storing the most important program data (other common blocks serve for the internal purposes and usually do not need to be accessed by users). As mentioned in the previous section, supersymmetric input parameters should be given at the SUSY scale (only for some SM parameters, like running quark masses, the input scale is user defined).

By default, **SUSY_FLAVOR** uses the following implicit type declaration in all routines:

```
implicit double precision (a-h,o-z)
```

so that all variables with the names starting from `a` to `h` and from `o` to `z` are automatically defined as `double precision` and those with names starting from `i` to `n` are of `integer` type. In what follows we explicitly indicate variables that do not obey this rule. Such variables are always listed in explicit type statements inside the procedures. Complex

b_fun.f:	general 2-point loop functions
bsg_nl.f:	formulae for $\text{Br}(B \rightarrow X_s\gamma)$, including QCD corrections
c_fun.f:	general 3-point loop functions
cd_fun.f:	3-, 4- and some 5-point loop functions at vanishing external momenta
cdm_q.f:	u - and d -quark chromoelectric dipole moments
cdm_g.f:	gluon chromoelectric dipole moment
db_fun.f:	derivatives of general 2-point loop functions
dd_gamma.f:	d quark- d quark-photon 1-loop triangle diagram
ddg_fun.f:	general gauge boson-fermion-fermion 1-loop triangle diagram
dd_gluon.f:	d quark- d quark-gluon 1-loop triangle diagram
dd_ll.f:	d quark- d quark-lepton-lepton 1-loop box diagram
dd_mix.f:	4- d quark 1-loop box diagram
dd_vv.f:	d quark- d quark-neutrino-neutrino 1-loop box diagram
d_self0.f:	full d -quark self-energy
edm_q.f:	u - and d -quark electric dipole moments
eisch1.f:	auxiliary numerical routine - hermitian matrix diagonalization
l_self0_dlim.f:	routines for the various decompositions of the lepton self energies
ll_gamma.f:	lepton-lepton-photon 1-loop triangle diagram
mh_diag.f:	diagonalization of tree level mass matrices angles
mh_init.f:	initialization of MSSM parameters
phen_21.f:	formulae for $\text{Br}(\mu \rightarrow e\gamma)$, $\text{Br}(\tau \rightarrow \mu\gamma, e\gamma)$, lepton $g - 2$ anomaly and EDMs
phen_2q.f:	formulae for $\text{Br}(K_L^0 \rightarrow \pi^0\bar{\nu}\nu)$, $\text{Br}(K^+ \rightarrow \pi^+\bar{\nu}\nu)$, $\text{Br}(B_{s(d)} \rightarrow l^+l^-)$, $\text{Br}(B \rightarrow \tau\nu, D\tau\nu)$ and neutron EDM
phen_4q.f:	formulae for the meson mixing observables: Δm_K , ϵ_K , Δm_D , $\Delta m_{B_{d(s)}}$
qcd_fun.f:	auxiliary QCD calculations - running α_s , running quark masses etc.
q_self0_dlim.f:	routines for the various decompositions of the u - and d -quark self energies
rombint.f:	auxiliary numerical routine - Romberg numerical integration
sflav_io.f:	input/output routines for the SLHA2 data format
sflav_main.f:	main routine calculating all physical observables
sff_fun.f:	general scalar-fermion-fermion 1-loop triangle diagram
suu_vert.f:	CP-even neutral Higgs boson- u quark- u quark 1-loop triangle diagram
u_self0.f:	u -quark self-energy
uu_mix.f:	full 4- u quark 1-loop box diagram
vegas.f:	auxiliary numerical routine - Vegas Monte Carlo integration
vf_def.f:	definitions of fermion tree-level vertices
vg_def.f:	definitions of gauge boson tree-level vertices
vh_def.f:	definitions of Higgs boson tree-level vertices
yuk_ren.f:	renormalization of the Yukawa couplings and CKM matrix including the chirally enhanced corrections
zdd_vert0.f:	Z boson- d quark- d quark 1-loop triangle diagram

Table 4: Alphabetical list of files included in **SUSY_FLAVOR** library.

parameters mentioned in this article are declared in `SUSY_FLAVOR` as `double complex` type. Mass parameters are always given in GeV.

`SUSY_FLAVOR` provides two ways of initializing input parameters. Firstly, they can be read from the file `susy_flavor.in`. The structure of this file follows the SLHA2 convention [65], with optional extensions which we describe in Sec. 4.1. Initializing parameters in the input file is done by a call to single subroutine `sflav_input` and does not require detailed knowledge of the program internal structure. This option, as it require disk file access for each parameter set may not be most efficient for scans over the MSSM parameter space. Therefore, `SUSY_FLAVOR` provides also a set of routines designed to initialize parameters defined in the program, which can be used to prepare programs that scan over large parameter sets. As described in Sec. 4.2, these routines require more care in use, as they should be initialized in the proper order, i.e. the gauge sector first, then the fermion sector, Higgs sector, and SUSY sectors at the end (the initialization sequences for the gaugino, slepton and squark sectors are independent).

An examples of an initialization sequence for `SUSY_FLAVOR`, illustrating both options mentioned above, is presented Appendix B. The sample input file `susy_flavor.in` is given in Appendix C. Test output generated for parameters used in Appendices B and C is enclosed in Appendix D.

4.1 Parameter initialization from the input file

Input parameters for `SUSY_FLAVOR` can be set by editing appropriate entries of the file `susy_flavor.in` and subsequently calling the subroutine `sflav_input`, which reads the input file, stores the the MSSM Lagrangian parameters in FORTRAN common blocks and calculates tree-level physical masses and mixing matrices. After calling `sflav_input`, all physical observable described in Sec. 6 can be calculated. The input file `susy_flavor.in` is written in the SLHA2 format, with some extensions which we list below.

The initialization proceeds as follows. Before reading the input file, all parameters are set to some initial values. They are:

- basic SM parameters

$$\begin{aligned}\alpha_{em}(M_Z) &= 1/127.934 \\ \alpha_s(M_Z) &= 0.1172 \\ M_Z &= 91.1876 \text{ GeV} \\ M_W &= 80.398 \text{ GeV} \\ s_W^2(\text{MSBar}) &= 0.232\end{aligned}$$

- running quark masses:

$$m_u(2 \text{ GeV}) = 2.15 \text{ MeV}$$

$$\begin{aligned}
m_d(2 \text{ GeV}) &= 4.7 \text{ MeV} \\
m_s(2 \text{ GeV}) &= 93.5 \text{ MeV} \\
m_c(m_c) &= 1.275 \text{ GeV} \\
m_b(m_b) &= 4.18 \text{ GeV}
\end{aligned}$$

- pole fermion masses:

$$\begin{aligned}
m_t &= 173.5 \text{ GeV} \\
m_e &= 0.5109989 \text{ MeV} \\
m_\mu &= 105.658 \text{ MeV} \\
m_\tau &= 1.77684 \text{ GeV}
\end{aligned}$$

- CKM Wolfenstein parameters $\lambda = 0.2258$, $A = 0.808$, $\bar{\rho} = 0.177$, $\bar{\eta} = 0.36$.
- all MSSM mass parameters (μ , gaugino and sfermion masses, trilinear A terms) are set to 0. $\tan \beta$ and the CP-odd Higgs mass M_A , which we use as the input parameters for the Higgs sector, are also set to 0.
- hadronic-related parameters (QCD scales and effective coefficients, hadronic matrix elements etc.) are set to values described in Sections 6.1–6.11. Their compact list is given in **Block** `SFLAV_HADRON` in Appendix C.

Subsequently, input Blocks are read from the file `susy_flavor.in` in the following order: `SOFTINP`, `SMINPUTS`, `VCKMIN`, `MINPAR` ($\tan \beta$ only, other entries ignored), `EXTPAR`, `IMEXTPAR`, `MSL2IN`, `IMMSL2IN`, `MSE2IN`, `IMMSE2IN`, `TEIN`, `IMTEIN`, `TEINH`, `IMTEINH`, `MSQ2IN`, `IMMSQ2IN`, `MSU2IN`, `IMMSU2IN`, `MSD2IN`, `IMMSD2IN`, `TUIN`, `IMTUIN`, `TUINH`, `IMTUINH`, `TDIN`, `IMTDIN`, `TDINH`, `IMTDINH`, `SFLAV_HADRON`.

In principle presence of *any* Block is optional - if given Block is absent, program falls back to default parameter values listed above. Obviously, at least flavor-diagonal SUSY mass parameters have to be defined in at least one Block, otherwise vanishing default masses cause program crash. If a parameter is multiply defined in several Blocks (for example left slepton mass parameters in **Block** `EXTPAR` and later in **Blocks** `MSL2IN`, `IMMSL2IN`), the value from Block read as latest in the list above overwrites (without warning!) the values from preceding Blocks. Blocks do not need to be complete, i.e. to contain all entries described in SLHA2 specification - it is sufficient to define minimal set of parameters relevant for given problem, others would be filled with default values.

Comparing to standard SLHA2 conventions, `SUSY_FLAVOR` uses following extensions:

1. We define an optional **Block** `SOFTINP` defining choice of input conventions. If such block is not present, program assumes default values of control variables:

Variable value	Sfermion sector parametrization
<code>iconv = 1</code>	default: MSSM parameters defined in SLHA2 conventions.
<code>iconv = 2</code>	MSSM parameters defined in conventions of Ref. [18].
<code>input_type = 1</code>	off-diagonal soft terms are given as dimensionless mass insertions.
<code>input_type = 2</code>	default: sfermion soft terms given as absolute dimensional values.
<code>ilev = 0</code>	no resummation of chirally enhanced corrections performed, all SUSY contributions taken strictly at 1-loop level.
<code>ilev = 1</code>	resummation of chirally enhanced corrections performed with the use of analytical formulae valid in the decoupling limit $M_{SUSY} \gg v_1, v_2$.
<code>ilev = 2</code>	default: resummation of chirally enhanced corrections performed using the numerical iterative solutions for bare Yukawa couplings and CKM matrix elements.

2. `SUSY_FLAVOR` uses two non-standard (comparing to SLHA2) entries of `Block SMINPUTS`. Entry 30 is used to define M_W and entry 31 to define s_W^2 in MSbar renormalization scheme.

3. Following the SLHA2 convention, full sfermion soft mass matrices can be defined in the `MSL2IN`, `MSE2IN`, `MSQ2IN`, `MSD2IN`, `MSU2IN` and `IMMSL2IN`, `IMMSE2IN`, `IMMSQ2IN`, `IMMSD2IN`, `IMMSU2IN` blocks. The `input_type` parameter in the `SOFTINP` block defines the dimension of the off-diagonal terms. If `input_type = 1`, the off-diagonal entries given in `susy_flavor.in` are assumed to be dimensionless mass insertions δ_X^{IJ} and the actual flavor violating sfermion soft mass terms are calculated as

$$(m_X^2)_{IJ} = (m_X^2)_{JI}^* = \delta_X^{IJ} \sqrt{(m_X^2)_{II}(m_X^2)_{JJ}}, \quad (4)$$

where $X = L, E, Q, U, D$ and I, J enumerate superpartners of the mass-eigenstate quarks.

4. The blocks `TEIN`, `TDIN`, `TUIN` and `IMTEIN`, `IMTDIN`, `IMTUIN` define the full trilinear SUSY breaking terms. They are in general not hermitian and one is required to define all entries. Again the parameter `input_type` defines the format and dimension of the off-diagonal terms. If `input_type = 1`, then all relevant `susy_flavor.in` entries are treated as dimensionless numbers and expanded to the full trilinear SUSY breaking terms as:

$$\begin{aligned} A_l^{IJ} &= \delta_{LLR}^{IJ} \left((m_L^2)_{II} (m_E^2)_{JJ} \right)^{\frac{1}{4}}, \\ A_d^{IJ} &= \delta_{DLR}^{IJ} \left((m_Q^2)_{II} (m_D^2)_{JJ} \right)^{\frac{1}{4}}, \\ A_u^{IJ} &= \delta_{ULR}^{IJ} \left((m_Q^2)_{II} (m_U^2)_{JJ} \right)^{\frac{1}{4}}. \end{aligned} \quad (5)$$

Note that A -terms are normalized to diagonal sfermion masses, not to the diagonal trilinear terms, and that in eq. (5) for simplicity we use $(m_Q^2)_{II}$ as the diagonal mass scale for both up and down left squark fields (in general related by the CKM rotation).

5. The “non-holomorphic” LR mixing terms of eq. (3) are not included in the SLHA2 specification of the MSSM parameters. They can be defined if necessary in blocks `TEINH`,

`TDINH`, `TUINH` and `IMTEINH`, `IMTDINH`, `IMTUINH`. If such blocks are not present, all such terms are set to 0. As standard LR mixing terms, non-holomorphic ones are also not hermitian in general. Again depending on the value of `input_type` they can be given as dimensionful or dimensionless. In the second case (`input_type = 1`) the dimensionful non-holomorphic terms are calculated in a way analogous to eq. (5).

4.2 Parameter initialization inside the program

`SUSY_FLAVOR` input parameters can be initialized directly inside the driver program using the set of routines described below. Before the proper initialization sequence, the user can set the `iconv` variable value to choose the input convention:

<code>common/sf_cont/eps,indx(3,3),iconv</code>	
<code>iconv=1</code>	SLHA2 [65] input conventions
<code>iconv=2</code>	[18] input conventions

After choosing the input conventions, one should subsequently initialize the gauge, matter fermion, Higgs, SUSY fermion and sfermion sectors (exactly in this order), using the procedures described in detail in the following sections.

4.2.1 Gauge sector

As input, `SUSY_FLAVOR` takes the gauge boson masses (M_W, M_Z) and the gauge coupling constants (electromagnetic and strong) at the M_Z scale. They are initialized by:

Routine and arguments	Purpose and MSSM parameters
<code>vpar_update(zm,wm,alpha_em,st2)</code>	Sets electromagnetic sector parameters
<code>zm</code>	M_Z , Z boson mass
<code>wm</code>	M_W , W boson mass
<code>alpha_em</code>	$\alpha_{em}(M_Z)$, QED coupling at M_Z scale
<code>st2</code>	s_W^2 in MSBar scheme
<code>lam_fit(alpha_s)</code>	Sets $\alpha_s(M_Z)$ and Λ_{QCD} for 4-6 flavors at the NNLO level
<code>lam_fit_nlo(alpha_s)</code>	Sets $\alpha_s(M_Z)$ and Λ_{QCD} for 4-6 flavors at the NLO level
<code>alpha_s</code>	$\alpha_s(M_Z)$, strong coupling at M_Z scale

4.2.2 Matter fermion sector

`SUSY_FLAVOR` assumes that neutrinos are massless. Pole masses of the charged leptons are initialized in the file `mh_init.f` in block data `init_phys`. They are stored in the `em` array in `common/fmass/em(3),um(3),dm(3)` and can be directly modified there. Their default values are:

Lepton mass	Value
m_e	$\text{em}(1) = 0.0005109989$
m_μ	$\text{em}(2) = 0.105658$
m_τ	$\text{em}(3) = 1.77684$

In the quark sector the most important input parameters are the running top and bottom masses at a given renormalization scale and the CKM matrix angles and phase. They can be set by:

Routine and arguments	Purpose and MSSM parameters
<code>init_fermion_sector(tm,tscale,bm,bscale)</code>	Sets running top and bottom quark mass
<code>tm,tscale</code>	$m_t(\mu_t)$, running $\overline{\text{MS}}$ top quark mass
<code>bm,bscale</code>	$m_b(\mu_b)$, running $\overline{\text{MS}}$ bottom quark mass
<code>ckm_init(s12,s23,s13,delta)</code>	Option 1: initialization of the CKM matrix
<code>s12,s23,s13</code>	$\sin \theta_{12}, \sin \theta_{23}, \sin \theta_{13}$, sines of the CKM angles
<code>delta</code>	δ , the CKM phase in radians
<code>ckm_wolf(alam,a,rhobar,etabar)</code>	Option 2: initialization of the CKM matrix
<code>alam,a,rhobar,etabar</code>	Wolfenstein parameters $\lambda, A, \bar{\rho}, \bar{\eta}$

The light quark masses are also initialized in the `block data init_phys` of the file `mh_init.f` and stored in `common/fmass_high/umu(3),uml(3),amu(3),dmu(3),dml(3),amud(3)`. The arrays `uml(dml)` contain up(down) quark masses at the scale `amu(amu)`, respectively. Their default values are:

Running quark mass	Mass value	Mass scale
$m_d(\mu_d)$	$dml(1) = 0.0047$	$amud(1) = 2$
$m_s(\mu_s)$	$dml(2) = 0.0935$	$amud(2) = 2$
$m_b(\mu_b)$	$dml(3) = 4.18$	$amud(3) = 4.18$
$m_u(\mu_u)$	$uml(1) = 0.00215$	$amu(1) = 2$
$m_c(\mu_c)$	$uml(2) = 1.275$	$amu(2) = 1.275$
$m_t(\mu_t)$	$uml(3) = 163.1$	$amu(3) = 163.1$

The variables of the arrays `uml`, `amu`, `dml`, `amud` can be directly accessed and modified if necessary. However, for consistency, after such modifications the user should call the routine `init_run_qmass` which calculates running quark masses at the high m_t scale (stored in `common/fmass_high/` in the arrays `umu`, `dmu` and in `common/fmass/` in the arrays `um`, `dm`) for later use in the running Yukawa couplings and in SUSY loop calculations.

4.2.3 Higgs sector

Following the common convention, we take the Higgs mixing parameter μ , the CP-odd Higgs boson mass M_A , and the ratio of vacuum expectation values $\tan \beta = v_2/v_1$ as the input parameters (in order to calculate values of Higgs mass terms in the Lagrangian, one

needs to set also the μ parameter already here):

```
subroutine init_higgs_sector(pm,tb,amu,ierr)
```

Argument MSSM parameters

pm CP-odd Higgs mass M_A

tb Ratio of Higgs VEVs, $\tan \beta = \frac{v_2}{v_1}$

amu Higgs mixing parameter μ (complex)

ierr output error code: $ierr \neq 0$ if Higgs sector initialization failed

4.2.4 Sfermion sector

SUSY_FLAVOR uses two subroutines to initialize sfermion parameters, `init_slepton_sector` and `init_squark_sector`. They accept as input diagonal masses and off-diagonal dimensionless mass insertions, expanded later to entries of the soft mass matrices as defined by eqs. (4,5) (this is only a choice of parametrization and does not lead to any loss of generality). The sfermion initialization routines have the following arguments:

```
subroutine init_slepton_sector(sll,slr,slmi_l,slmi_r,slmi_lr,slmi_lrp,ierr)
```

Argument MSSM parameters

sll Array of the diagonal left-handed slepton masses $(m_L^2)_{LL}^{II} = sll(I)^2$, $I = 1 \dots 3$

slr Array of the diagonal right-handed slepton masses $(m_L^2)_{RR}^{II} = slr(I)^2$, $I = 1 \dots 3$

slmi_l Array of the off-diagonal left-handed slepton mass insertions $(\delta_L)_{LL}^{12} = slmi_l(1)$, $(\delta_L)_{LL}^{23} = slmi_l(2)$, $(\delta_L)_{LL}^{13} = slmi_l(3)$ (complex parameters); remaining LL mass insertions are initialized via hermitian conjugation

slmi_r Array of the off-diagonal right-handed slepton mass insertions $(\delta_L)_{RR}^{12} = slmi_r(1)$, $(\delta_L)_{RR}^{23} = slmi_r(2)$, $(\delta_L)_{RR}^{13} = slmi_r(3)$ (complex parameters); the remaining RR mass insertions are initialized via hermitian conjugation

slmi_lr Matrix with the standard (holomorphic) slepton trilinear LR mass insertions $(\delta_L)_{LR}^{IJ} = slmi_lr(I, J)$, $I, J = 1 \dots 3$ (complex parameters)

slmi_lrp Matrix with the non-holomorphic slepton trilinear LR mass insertions $(\delta'_L)_{LR}^{IJ} = slmi_lrp(I, J)$, $I, J = 1 \dots 3$ (complex parameters)

ierr output error code: $ierr \neq 0$ if slepton sector initialization failed (negative physical slepton mass²)

subroutine init_squark_sector(sql,squ,sqd,asu,asd,sqmi_l,sumi_r,sdmi_r, sumi_lr,sdmi_lr,sumi_lrp,sdmi_lrp,ierr)	
Argument	MSSM parameters
sql	Array of the diagonal left-handed down-squark masses $(m_D^2)_{LL}^{II} = \text{sql}(I)^2$, $I = 1 \dots 3$
squ	Array of the diagonal right-handed up-squark masses $(m_U^2)_{RR}^{II} = \text{squ}(I)^2$, $I = 1 \dots 3$
sqd	Array of the diagonal right-handed down-squark masses $(m_D^2)_{RR}^{II} = \text{sqd}(I)^2$, $I = 1 \dots 3$
sqmi_l	Array of the off-diagonal left-handed down squark mass insertions $(\delta_D)_{LL}^{12} = \text{sqmi_l}(1)$, $\delta_D_{LL}^{23} = \text{sqmi_l}(2)$, $(\delta_D)_{LL}^{13} = \text{sqmi_l}(3)$ (complex parameters); remaining down LL mass insertions are initialized via hermitian conjugation; up LL mass matrix obtained via $SU(2)$ relation
sumi_r	Array of the off-diagonal right-handed up-squark mass insertions $(\delta_U)_{RR}^{12} = \text{sumi_r}(1)$, $(\delta_U)_{RR}^{23} = \text{sumi_r}(2)$, $(\delta_U)_{RR}^{13} = \text{sumi_r}(3)$ (complex parameters); remaining up RR mass insertions are initialized via hermitian conjugation
sdmi_r	Array of the off-diagonal right-handed down-squark mass insertions $(\delta_D)_{RR}^{12} = \text{sdmi_r}(1)$, $(\delta_D)_{RR}^{23} = \text{sdmi_r}(2)$, $(\delta_D)_{RR}^{13} = \text{sdmi_r}(3)$ (complex parameters); remaining down RR mass insertions are initialized via hermitian conjugation
sumi_lr	Matrix with the standard (holomorphic) up-squark trilinear LR mass insertions $(\delta_U)_{LR}^{IJ} = \text{sumi_lr}(I, J)$, $I, J = 1 \dots 3$ (complex parameters)
sdmi_lr	Matrix with the standard (holomorphic) down-squark trilinear LR mass insertions $(\delta_D)_{LR}^{IJ} = \text{sdmi_lr}(I, J)$, $I, J = 1 \dots 3$ (complex parameters)
sumi_lrp	Matrix with the non-holomorphic up-squark trilinear LR mass insertions $(\delta'_U)_{LR}^{IJ} = \text{sumi_lp}(I, J)$, $I, J = 1 \dots 3$ (complex parameters)
sdmi_lrp	Matrix with the non-holomorphic down-squark trilinear LR mass insertions $(\delta'_D)_{LR}^{IJ} = \text{sdmi_lp}(I, J)$, $I, J = 1 \dots 3$ (complex parameters)
ierr	output error code: $ierr \neq 0$ if squark sector initialization failed (negative physical squark mass ²)

4.2.5 Supersymmetric fermion sector

Initialization is done by the routine `init_ino_sector`:

subroutine init_ino_sector(gm1, gm2, gm3, amu, tb, ierr)	
Argument	MSSM parameters
gm1, gm2	$U(1), SU(2)$ gaugino masses (complex)
gm3	$SU(3)$ gaugino mass
tb	$\tan \beta = \frac{v_2}{v_1}$, the ratio of Higgs VEVs
amu	the Higgs mixing parameter μ (complex)
ierr	output warning code: $ierr \neq 0$ for chargino or neutralino lighter than $M_Z/2$

If one sets $M_1 = 0$ in the call to `init_ino_sector` then the GUT-derived relation $M_1 = \frac{5}{3} \tan^2 \theta_W M_2$ is used for M_1 .

4.3 Tree-level physical masses and mixing angles

After performing the full initialization sequence in `SUSY_FLAVOR`, all the MSSM Lagrangian parameters, physical tree-level particle masses (with the exception of the running quark masses), and mixing matrices are calculated and stored in common blocks. If necessary, they can be directly accessed and modified. Note, however, that after any modifications of the Lagrangian parameters, relevant procedures calculating physical masses and mixing angles have to be called again. In Table 5 we list the important blocks storing MSSM parameters. Common blocks containing masses and mixing angles are listed in Table 6.

5 Resummation of chirally enhanced corrections

The resummation of the chirally enhanced corrections, including the threshold corrections to Yukawa couplings and CKM matrix elements, is an important new feature added to `SUSY_FLAVOR` in version 2.0. Such corrections arise in the case of large values of $\tan \beta$ or large trilinear SUSY-breaking terms. They formally go beyond the 1-loop approximation, but should be included due to their numerical importance². Implementation of the resummation in `SUSY_FLAVOR` follows the systematic approach of ref. [17] and takes into account all contributions involving sfermions and gauginos (gluino, chargino and neutralino exchanges). The level of resummation is a user selectable option and can be done using the following routine:

Routine:	<code>subroutine set_resummation_level(ilev,ierr)</code>
Input:	<code>ilev=0</code> : no resummation <code>ilev=1</code> : analytical solution used for bare Yukawa couplings and the bare CKM matrix elements (i.e. parameters of the superpotential), valid in the “decoupling limit” $M_{SUSY} \gg v_1, v_2$ <code>ilev=2</code> : exact iterative numerical solution for the bare Yukawa couplings and the bare CKM matrix elements.
Output:	<code>ierr=0</code> : resummation successful <code>ierr<0</code> : exact resummation (<code>ilev=2</code>) requested but failed (no convergence), instead analytical resummation in the decoupling limit performed successfully <code>ierr>0</code> : resummation failed (both for <code>ilev=1,2</code>), only 1-loop expressions will be used in calculations of the physical observables

Details of calculations: Ref. [17]

²It is even possible that the light fermion masses and off-diagonal CKM elements are generated entirely by chirally-enhanced self-energies involving the trilinear A -terms [68]

Common block and variables	Lagrangian parameters
common/vpar/st,ct,st2,ct2,sct,sct2,e,e2,alpha,wm,wm2,zm,zm2,pi,sq2 st,ct,st2,ct2,sct,sct2 e,e2,alpha wm,wm2,zm,zm2 pi,sq2	Weinberg angle functions, respectively $s_W, c_W, s_W^2, c_W^2, s_W c_W, s_W^2 c_W^2$ electric charge powers at M_Z scale: e, e^2, α_{em} gauge boson masses: M_W, M_W^2, M_Z, M_Z^2 numerical constants, π and $\sqrt{2}$
common/hpar/hm1,hm2,hm12,hmu hm1,hm2 hm12 hmu	soft Higgs masses $m_{H_1}^2, m_{H_2}^2$ soft Higgs mixing parameter m_{12}^2 Higgs mixing parameter μ (complex)
common/vev/v1,v2 v1,v2	Higgs vacuum expectation values v_1, v_2
common/yukawa/y1(3),yu(3),yd(3) y1(3) yu(3) yd(3)	charged lepton Yukawa couplings Y_e, Y_μ, Y_τ (complex) Running $\overline{\text{MS}}$ up-quark Yukawa couplings at m_t scale: Y_u, Y_c, Y_t Running $\overline{\text{MS}}$ down-quark Yukawa couplings at m_t scale: Y_u, Y_c, Y_t
common/gmass/gm3,gm2,gm1 gm1,gm2 gm3	$U(1), SU(2)$ gaugino masses M_1, M_2 (complex) $SU(3)$ gaugino mass M_3
common/msoft/lms(3,3),rms(3,3),ums(3,3),dms(3,3),qms(3,3) lms(3,3),rms(3,3) ums(3,3),dms(3,3),qms(3,3)	hermitian slepton soft mass matrices m_L^2, m_E^2 (complex) hermitian squark soft mass matrices m_U^2, m_D^2, m_Q^2 (complex)
common/soft/ls(3,3),ks(3,3),ds(3,3),es(3,3),us(3,3),ws(3,3) ls(3,3),ds(3,3),us(3,3) ks(3,3),es(3,3),ws(3,3)	trilinear soft SUSY breaking terms A_l, A_d, A_u (complex) trilinear “non-holomorphic” soft SUSY breaking terms A'_l, A'_d, A'_u (complex)

Table 5: Common blocks storing the MSSM Lagrangian parameters.

Common block and variables	Masses and mixing matrices
common/fmass / em(3) , um(3) , dm(3)	
em(3)	Charged lepton pole masses m_e, m_μ, m_τ
um(3)	Running $\overline{\text{MS}}$ up-quark masses at the m_t scale: m_u, m_c, m_t
dm(3)	Running $\overline{\text{MS}}$ down-quark masses at the m_t scale: m_u, m_c, m_t
common/hmass / cm(2) , rm(2) , pm(2) , zr(2,2) , zh(2,2)	
rm(2)	neutral CP-even Higgs masses $\text{rm}(1) = M_H,$ $\text{rm}(2) = M_h$
pm(2)	neutral CP-odd Higgs mass $\text{pm}(1)$ and Goldstone mass $\text{pm}(2)$
cm(2)	charged Higgs mass $\text{cm}(1)$ and charged Goldstone mass $\text{cm}(2)$
zr(2,2)	CP-even Higgs mixing matrix Z_R
zh(2,2)	CP-odd and charged Higgs mixing matrix Z_H
common/charg / fcm(2) , zpos(2,2) , zneg(2,2)	
fcm(2)	chargino masses $M_{\chi_i^+}, i = 1, 2$
zpos(2,2) , zneg(2,2)	chargino mixing matrices Z_+, Z_- (complex)
common/neut / fnm(4) , zn(4,4)	
fnm(4)	neutralino masses $M_{\chi_i^0}, i = 1 \dots 4$
zn(4,4)	neutralino mixing matrix Z_N (complex)
common/slmass / vm(3) , slm(6) , zv(3,3) , z1(6,6)	
vm(3)	sneutrino masses $M_{\tilde{\nu}_I}, I = 1 \dots 3$
slm(6)	charged slepton masses $M_{L_i}, i = 1 \dots 6$
zv(3,3)	sneutrino mixing matrix $Z_{\tilde{\nu}}$ (complex)
z1(6,6)	charged slepton mixing matrix Z_L (complex)
common/sqmass / sum(6) , sdm(6) , zu(6,6) , zd(6,6)	
sum(6)	up-squark masses $M_{U_i}, i = 1 \dots 6$
sdm(6)	down-squark masses $M_{D_i}, i = 1 \dots 6$
zu(6,6)	up-squark mixing matrix Z_U (complex)
zd(6,6)	down-squark mixing matrix Z_D (complex)

Table 6: Common blocks storing particle masses and mixing matrices.

After call to `set_resummation_level` with `ilev ≠ 0`, `SUSY_FLAVOR` calculates the values of bare Yukawa couplings and CKM matrix elements (i.e. the values of the MSSM Lagrangian parameters) and starts to use in loop calculations appropriately corrected effective Higgs boson and supersymmetric particle couplings, automatically taking into account resummation of enhanced higher order terms.

One should keep in mind that if the chirally enhanced corrections are very large relation between bare and effective physical fermion masses and CKM matrix elements involve a significant degree of fine-tuning and one might also encounter numerical instabilities using the program. Therefore, the routines performing the resummation should be used with care. One can reasonably assume that resummation works properly in the decoupling limit $v_1, v_2 \ll M_{SUSY}$ as long as the difference between the bare and physical quantities is at least not significantly larger than the physical values themselves. Setting the actual “safety condition” is left to the `SUSY_FLAVOR` users. To facilitate that, the blocks `SFLAV_CHIRAL_YUKAWA` and `SFLAV_CHIRAL_CKM` in the `SUSY_FLAVOR` output file list the relative size of differences between the bare Yukawa couplings and CKM matrix elements of the superpotential and the (effective) physical quantities, calculated as

$$\delta X_{corr} = \left| \frac{X_{\text{bare}} - X_{\text{effective}}}{X_{\text{effective}}} \right| \quad (6)$$

One can use such output to define conditions rejecting points of the MSSM parameters space where the resummation effects are too large and calculations cannot be trusted. In our numerical experience, the stability of `SUSY_FLAVOR` results requires the relative size of the resummed loop corrections to be at most of order one for CKM elements and Yukawa couplings of 2nd and 3rd generation. Thus, if the chosen input respects ’t Hooft’s naturalness argument [30, 34], also the resummation of all chirally enhanced effects can be performed analytically in the decoupling limit and is stable numerically.

6 List of processes

In this section we list the set observables whose computation is fully implemented in `SUSY_FLAVOR` v2.0. QCD corrections and hadronic matrix elements are extracted mostly from various analyses done within the Standard Model. They are assumed to work reasonably well also in the MSSM since supersymmetric strong corrections from gluino and squarks are suppressed by large masses of these particles.

Results of calculations of the hadronic matrix elements can differ depending on the methods used and thus carry significant theoretical uncertainties. Therefore, in `SUSY_FLAVOR`, hadronic matrix element estimates and other QCD related quantities are treated as external parameters. They are initialized to the default values listed below for

each observable and can be directly modified by users by changing the relevant variables in the common blocks where they are stored. Currently most of the hadronic (and related) input parameters used in **SUSY_FLAVOR** are taken from the Table 3 of Ref. [69].

In most cases, QCD and hadronic corrections are known to a precision at the level of few percent to tens of percent, while variations of supersymmetric flavor and CP-violating parameters can change observables by orders-of-magnitude. Thus, as long as the MSSM parameters are not measured very precisely, the current implementation of strong corrections is sufficient for analyses performed in the framework of the general MSSM.

6.1 $g - 2$ magnetic moment anomaly for leptons

Anomalous magnetic moment of leptons are defined as the coefficient $a_{lI} \equiv (g_I - 2)/2$ in the effective Hamiltonian for the flavor-diagonal lepton-lepton-photon interaction:

$$\mathcal{H}_l = -\frac{e}{4m_l} a_I \bar{l}^I \sigma_{\mu\nu} l^I F^{\mu\nu}, \quad (7)$$

where $I = 1, 2, 3$ is the generation index of the lepton³. In **SUSY_FLAVOR** supersymmetric contribution to $(g - 2)$ anomaly (to be added to the SM one) is calculated by the routine:

Routine:	<code>double precision function g_minus_2_susy(I)</code>
Input:	$I = 1, 2, 3$ for e, μ, τ respectively
Output:	SUSY contribution to $a_I = (g_I - 2)/2$ for the charged lepton specified by I
QCD related factors:	none, QCD corrections are small and not included
Details of calculations:	Performed by authors, unpublished

6.2 Electric Dipole Moments of charged leptons

Lepton EDMs are defined as another coefficient d_{lI} in the effective Hamiltonian for the flavor-diagonal lepton-lepton-photon interaction:

$$\mathcal{H}_l = \frac{id_{lI}}{2} \bar{l}^I \sigma_{\mu\nu} \gamma_5 l^I F^{\mu\nu}, \quad (8)$$

where $I = 1, 2, 3$ is again the generation index of the lepton. In **SUSY_FLAVOR** lepton EDM is calculated by:

³The measurement of the anomalous magnetic moment of the muon is used to determine α . In order to consider the possible effect of new physics one needs an independent determination of α [33] - e.g. one can use the measurements of the Rubidium atom [70].

Routine:	<code>double precision function edm_l(I)</code>
Input:	$I = 1, 2, 3$ for e, μ, τ respectively
Output:	EDM for the charged lepton specified by I (in the units $e\text{ cm}$)
QCD related factors:	none, QCD corrections are small and not included
Details of calculations:	Ref. [7] (note that EDM are defined there with opposite relative sign to <code>SUSY_FLAVOR</code> convention)

6.3 Neutron Electric Dipole Moment

The neutron EDM can be approximated by the sum of the electric dipole moments of the constituent d and u quarks plus contributions of the chromoelectric dipole moments (CDM) of quarks and gluons. The EDMs of the individual quarks are defined analogously to eq. (8). The CDM c_q of quark q is defined as:

$$\mathcal{H}_c = -\frac{ic_q}{2}\bar{q}\sigma_{\mu\nu}\gamma_5 T^a q G^{\mu\nu a}. \quad (9)$$

The gluonic dipole moment c_g is defined as:

$$\mathcal{H}_g = -\frac{c_g}{6}f_{abc}G_{\mu\rho}^a G_{\nu}^{b\rho} G_{\lambda\sigma}^c \epsilon^{\mu\nu\lambda\sigma}. \quad (10)$$

The exact calculation of the neutron EDM requires knowledge of its hadronic wave function. `SUSY_FLAVOR` uses the formulae:

$$E_n = \eta_{edd}d_d + \eta_{eu}d_u + e(\eta_{cd}c_d + \eta_{cu}c_u) + \frac{e\eta_g\Lambda_X}{4\pi}c_g \quad (11)$$

where η_i and Λ_X are the QCD correction factors [72] and the chiral symmetry breaking scale [71], respectively. Various models give significantly different η_i factors. Thus the `SUSY_FLAVOR` result should be treated as an order of magnitude estimate only. The calculations are performed by calling

Routine	<code>double precision function edm_n()</code>
Input	none
Output	neutron EDM
QCD related factors:	
<code>common/edm_qcd/eta_ed, eta_eu, eta_cd, eta_cu, eta_g, alamx</code>	
η_{ed}	<code>eta_ed = 0.79</code>
η_{eu}	<code>eta_eu = -0.2</code>
η_{cd}	<code>eta_cd = 0.59</code>
η_{cu}	<code>eta_cu = 0.3</code>
η_g	<code>eta_g = 3.4</code>
Λ_X	<code>alamx = 1.18</code>
Details of calculations:	Ref. [7]

6.4 $\mu \rightarrow e\gamma$ and $\tau \rightarrow e\gamma, \mu\gamma$ decay rates

The branching ratios for the flavor violating decays of a heavy lepton into a lighter lepton and photon are given by :

$$Br(l^J \rightarrow l^I \gamma) = \frac{48\pi^2 e^2 Br(l^J \rightarrow e\bar{\nu}\nu)}{m_{l^J}^2 G_F^2} (|C_L^{JI}|^2 + |C_R^{JI}|^2) \quad (12)$$

where $C_{L,R}^{IJ}$ are the relevant Wilson coefficients calculated from the 1-loop lepton-photon triangle diagram with an on-shell photon. The branching ratios are calculated by

Routine:	double precision function br_llg(J,I)
Input:	$J, I = 1, 2, 3$ for e, μ, τ respectively
Output:	branching ratios for $\mu \rightarrow e\gamma$ decay ($J = 2, I = 1$) and $\tau \rightarrow e\gamma, \mu\gamma$ decays ($J = 3, I = 1, 2$)
QCD related factors:	none, QCD corrections are small and not included
Details of calculations:	Performed by authors, unpublished

6.5 $K_L^0 \rightarrow \pi^0 \bar{\nu}\nu$ and $K^+ \rightarrow \pi^+ \bar{\nu}\nu$ decay rates

The relevant part of the effective Hamiltonian generated by the top quark and SUSY particle exchanges can be written as

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \frac{\alpha}{2\pi \sin^2 \theta_w} \sum_{l=e,\mu,\tau} [X_L(\bar{s}d)_{V-A}(\bar{\nu}_l \nu_l)_{V-A} + X_R(\bar{s}d)_{V+A}(\bar{\nu}_l \nu_l)_{V-A}]. \quad (13)$$

The branching ratios for the $K \rightarrow \pi \nu \bar{\nu}$ decays are then given by

$$Br(K^+ \rightarrow \pi^+ \bar{\nu}\nu) = \kappa_+ \left[\left(\frac{\Im(X_L + X_R)}{\lambda^5} \right)^2 + \left(\frac{\Re(K_{cs}^* K_{cd})}{\lambda} P_c + \frac{\Re(X_L + X_R)}{\lambda^5} \right)^2 \right] \quad (14)$$

$$Br(K_L^0 \rightarrow \pi^0 \bar{\nu}\nu) = \kappa_L \left(\frac{\Im(X_L + X_R)}{\lambda^5} \right)^2 \quad (15)$$

where κ [73], λ (the Wolfenstein parameters [74]), and the NLO charm quark contribution P_c [75, 76, 77] can be modified by **SUSY_FLAVOR** users (note that κ and P_c depend on V_{us} , m_c and α_s) The calculations of the branching ratios are performed by calling

Routine	<code>subroutine k_pivv(br_k0,br_kp)</code>
Input	<code>none</code>
Output	$\text{br_k0} = \text{Br}(K_L^0 \rightarrow \pi^0 \bar{\nu}\nu)$ $\text{br_kp} = \text{Br}(K^+ \rightarrow \pi^+ \bar{\nu}\nu)$
QCD related factors	
<code>common/kpivv/ak0,del_ak0,akp,del_akp,pc,del_pc,alam</code>	
$\kappa_L \pm \Delta\kappa_L$	$\text{ak0} = 2.231 \cdot 10^{-10}$, $\text{del_ak0} = 0.013 \cdot 10^{-10}$
$\kappa_+ \pm \Delta\kappa_+$	$\text{akp} = 5.173 \cdot 10^{-11}$, $\text{del_akp} = 0.025 \cdot 10^{-11}$
$P_c \pm \Delta P_c$	$\text{pc} = 0.41$, $\text{del_pc} = 0.03$
λ	$\text{alam} = 0.225$
Details of calculations:	Ref. [14]

6.6 $B_d^0 \rightarrow l^{I+}l^{J-}$ and $B_s^0 \rightarrow l^{I+}l^{J-}$ decay rates

The general expression for these branching ratios are rather complicated and can be found in [15]⁴. For most users it is sufficient to know that, in addition to the MSSM parameters, the dilepton B decays depend on the B meson masses and the hadronic matrix elements of the down quark vector and scalar currents:

$$\langle 0 | \bar{b} \gamma_\mu P_{L(R)} s | B_{s(d)}(p) \rangle = - (+) \frac{i}{2} p_\mu f_{B_{s(d)}} , \quad (16)$$

$$\langle 0 | \bar{b} P_{L(R)} s | B_{s(d)}(p) \rangle = + (-) \frac{i}{2} \frac{M_{B_{s(d)}}^2 f_{B_s}}{m_b + m_{s(d)}} , \quad (17)$$

where p_μ is the momentum of the decaying $B_{s(d)}$ -meson of mass $M_{B_{s(d)}}$. The $B_d^0 \rightarrow l^{I+}l^{J-}$ and $B_s^0 \rightarrow l^{I+}l^{J-}$ decay branching ratios are calculated by:

Routine	<code>double precision function b_ll(K,L,I,J)</code>
Input	$I, J = 1, 2, 3$ - outgoing leptons generation indices K, L - generation indices of the valence quarks of the B^0 meson: setting $(K, L) = (3, 1), (1, 3), (3, 2)$ and $(2, 3)$ chooses respectively $B_d^0, \bar{B}_d^0, B_s^0$ and \bar{B}_s^0 decay
Output	Branching ratios of the decay defined by K, L, I, J
QCD related factors	
<code>common/meson_data/dmk,amk,epsk,fk,dmd,amd,fd,amb(2),dmb(2),gam_b(2),fb(2)</code>	
M_{B_d}	$\text{amb}(1) = 5.2794$
M_{B_s}	$\text{amb}(2) = 5.368$
f_{B_d}	$\text{fb}(1) = 0.193$
f_{B_s}	$\text{fb}(2) = 0.232$
Details of calculations:	Ref. [15]

⁴Note that only the 1-loop electroweak/SUSY contributions to $B_{d,s}^0 \rightarrow l^{I+}l^{J-}$ are implemented in **SUSY_FLAVOR v2**. Thus, in the limit of heavy SUSY masses **SUSY_FLAVOR** reproduces older SM 1-loop estimates for such decays, somewhat higher than the NLO result given recently for $B_s \rightarrow \mu^+ \mu^-$ in [78]

6.7 $B \rightarrow (D)\tau\nu$ decay rates

SUSY_FLAVOR v2 calculates $Br(B \rightarrow (D)\tau\nu)$ including the SM and the charged Higgs contribution. The chirally enhanced corrections to Yukawa couplings, which also affect the charged Higgs contribution, are included. The relevant part effective Hamiltonian reads as ($I = 1$ for B_u^+ meson):

$$H_{eff}^I = \frac{G_F V_{qb}}{\sqrt{2}} \left[(\bar{q}^I \gamma_\mu (1 - \gamma_5) b) (\bar{l} \gamma_\mu (1 - \gamma_5) \nu) + C_{NP}^l (\bar{q}^I \gamma_\mu (1 + \gamma_5) b) (\bar{l} \gamma_\mu (1 - \gamma_5) \nu) \right] \quad (18)$$

where the New Physics C_{NP}^l contribution comes from the modification of the effective Yukawa couplings and read as

$$C_{NP}^l \approx -\frac{\sqrt{2}}{4m_{H^+}^2 G_F V_{qb}} \Gamma_{qb}^{H^+ LR} \Gamma_{\nu\tau}^{H^+ LR} \quad (19)$$

with $\Gamma_{qb}^{H^+ LR}, \Gamma_{\nu\tau}^{H^+ LR}$ defined in eqs. (48),(50) of ref. [17].

The decay rates are given by [79]:

$$Br(B_u \rightarrow \tau\nu) = \frac{G_F^2 |V_{ub}|^2}{8\pi} m_\tau^2 m_B f_B^2 \left(1 - \frac{m_\tau^2}{m_B^2}\right)^2 \left|1 + \frac{m_B^2}{m_b m_\tau} C_{NP}^\tau\right|^2 \quad (20)$$

$$\frac{Br(B_u \rightarrow D\tau\nu)}{Br(B_u \rightarrow D\nu)} = (0.28 \pm 0.02) \left(1 + 1.38(3)\text{Re}(C_{NP}^\tau) + 0.88(2) |C_{NP}^\tau|^2\right) \quad (21)$$

Branching ratios are calculated by:

Routine	<code>subroutine b_taunu(br_taunu,br_dtaunu_rat)</code>
Input	none
Output	<code>br_taunu = Br(B_u → τν)</code> <code>br_dtaunu_rat = Br(B_u → Dτν) / Br(B_u → Dν)</code>
QCD related factors	
common/meson_data/dmk, amk, epsk, fk, dmd, amd, fd, amb(2), dmb(2), gam_b(2), fb(2)	
f_{B_d}	<code>fb(1) = 0.193</code>
f_{B_s}	<code>fb(2) = 0.232</code>
Details of calculations:	Ref. [79, 17]

6.8 $B^0 \rightarrow X_s \gamma$ decay rate

Both the SUSY contributions and the QCD corrections to the calculation of the $B^0 \rightarrow X_s \gamma$ decay rate are quite complex. Their implementation in **SUSY_FLAVOR** is based on the SUSY loop calculations performed by the authors (not published in a general form) and on the QCD evolution published in [80]. There are no user-accessible QCD factors apart from the arguments of the `bkg_nl` routine.

Routine	double precision function <code>bkg_nl(del, amiu_b)</code>
Input	<code>del</code> - relative photon energy infrared cutoff scale, $E_\gamma \geq (1 - \text{del})E_\gamma^{\max}$, $0 < \text{del} < 1$ <code>amiu_b</code> - renormalization scale
Output	$\text{Br}(B \rightarrow X_s \gamma)$.
Details of calculations:	General SUSY diagrams unpublished, QCD corrections based on [80]

6.9 $\bar{K}^0 K^0$ meson mixing parameters

SUSY_FLAVOR calculates two parameters measuring the amount of CP-violation in neutral K meson oscillations: ε_K and the $\bar{K}^0 - K^0$ mass difference ΔM_K .

$$\Delta M_K = 2\Re \langle \bar{K}^0 | H_{\text{eff}}^{\Delta S=2} | K^0 \rangle , \quad (22)$$

$$\varepsilon_K = \frac{\exp(i\pi/4)}{\sqrt{2}\Delta M_K} \Im \langle \bar{K}^0 | H_{\text{eff}}^{\Delta S=2} | K^0 \rangle . \quad (23)$$

QCD dependent corrections are known with reasonable accuracy for the ε_K parameter. The long distance contributions to ΔM_K are large and difficult to control. Thus the result given by SUSY_FLAVOR for ΔM_K should be treated as an order of magnitude estimate only.

Apart from the MSSM parameters, the calculation of the $\bar{K}^0 K^0$ meson mixing requires knowledge of the meson masses and of the hadronic matrix elements of the following set of four-quark operators:

$$\begin{aligned} Q_1^{\text{VLL}} &= (\bar{q}_\alpha^I \gamma_\mu P_L q_\alpha^J)(\bar{q}_\beta^I \gamma^\mu P_L q_\beta^J), \\ Q_1^{\text{LR}} &= (\bar{q}_\alpha^I \gamma_\mu P_L q_\alpha^J)(\bar{q}_\beta^I \gamma^\mu P_R q_\beta^J), \\ Q_2^{\text{LR}} &= (\bar{q}_\alpha^I P_L q_\alpha^J)(\bar{q}_\beta^I P_R q_\beta^J), \\ Q_1^{\text{SLL}} &= (\bar{q}_\alpha^I P_L q_\alpha^J)(\bar{q}_\beta^I P_L q_\beta^J), \\ Q_2^{\text{SLL}} &= (\bar{q}_\alpha^I \sigma_{\mu\nu} P_L q_\alpha^J)(\bar{q}_\beta^I \sigma^{\mu\nu} P_L q_\beta^J) \end{aligned} \quad (24)$$

where α, β are color indices, for the $\bar{K}^0 K^0$ mixing one should choose flavor indices $I = 2$ and $J = 1$. The matrix elements can be written as:

$$\begin{aligned} \langle \bar{K}^0 | Q_1^{\text{VLL}}(\mu) | K^0 \rangle &= \frac{1}{3} M_K F_K^2 B_1^{\text{VLL}}(\mu), \\ \langle \bar{K}^0 | Q_1^{\text{LR}}(\mu) | K^0 \rangle &= -\frac{1}{6} \left(\frac{M_K}{m_s(\mu) + m_d(\mu)} \right)^2 M_K F_K^2 B_1^{\text{LR}}(\mu), \\ \langle \bar{K}^0 | Q_2^{\text{LR}}(\mu) | K^0 \rangle &= \frac{1}{4} \left(\frac{M_K}{m_s(\mu) + m_d(\mu)} \right)^2 M_K F_K^2 B_2^{\text{LR}}(\mu), \\ \langle \bar{K}^0 | Q_1^{\text{SLL}}(\mu) | K^0 \rangle &= -\frac{5}{24} \left(\frac{M_K}{m_s(\mu) + m_d(\mu)} \right)^2 M_K F_K^2 B_1^{\text{SLL}}(\mu), \\ \langle \bar{K}^0 | Q_2^{\text{SLL}}(\mu) | K^0 \rangle &= -\frac{1}{2} \left(\frac{M_K}{m_s(\mu) + m_d(\mu)} \right)^2 M_K F_K^2 B_2^{\text{SLL}}(\mu), \end{aligned} \quad (25)$$

where F_K is the K -meson decay constant. By default, **SUSY_FLAVOR** uses the B_i^X values at the scale $\mu = 2$ GeV given in [81] using the NDR renormalization scheme (quark masses at the scale 2 GeV are stored in `common/fmass_high/`, see Sec. 4.2.2).

In addition to the hadronic matrix elements, QCD corrections depend also on the “ η ” factors describing the evolution of the relevant Wilson coefficients from the high to low energy scale. These factors are automatically calculated at NLO by **SUSY_FLAVOR**. For the SM contribution to the Wilson coefficient of the Q^{VLL} operator a separate careful calculation of the evolution factors has been performed [82, 83]. Therefore **SUSY_FLAVOR** treats this contribution separately, setting B_{SM}^{VLL} and the η_{SM} factor to default values given in [84] (see [81] for a very detailed discussion of the structure of the QCD corrections in $\bar{B}^0 B^0$ and $\bar{K}^0 K^0$ systems, including their renormalization scheme dependence and calculations of the evolution factors “ η ” implemented in **SUSY_FLAVOR**).

The kaon mass difference ΔM_K and the ε_K parameter measuring the amount of CP violation in $\bar{K}^0 K^0$ mixing are calculated by

Routine	<code>subroutine dd_kaon(eps_k,delta_mk)</code>
Input	<code>none</code>
Output	<code>eps_k = ε_K parameter</code> <code>delta_mk = ΔM_K mass difference</code>

QCD related factors:

<code>common/meson_data/dmk,amk,epsk,fk,dmd,amd,fd,amb(2),dmb(2),gam_b(2),fb(2)</code>	
M_K	<code>amk = 0.497614</code>
Measured ΔM_K^{exp}	<code>dmk = $3.483 \cdot 10^{-15}$</code>
Measured ε_K^{exp}	<code>epsk = $2.229 \cdot 10^{-3}$</code>
f_K	<code>fk = 0.1598</code>
<code>common/bx_4q/bk(5),bd(5),bb(2,5),amu_k,amu_d,amu_b</code>	
$B_1^{\text{VLL}}(\mu_K)$	<code>bk(1) = 0.61</code>
$B_1^{\text{SLL}}(\mu_K)$	<code>bk(2) = 0.76</code>
$B_2^{\text{SLL}}(\mu_K)$	<code>bk(3) = 0.51</code>
$B_1^{\text{LR}}(\mu_K)$	<code>bk(4) = 0.96</code>
$B_2^{\text{LR}}(\mu_K)$	<code>bk(5) = 1.30</code>
Renormalization scale μ_K	<code>amu_k = 2</code>
<code>common/sm_4q/eta_cc,eta_ct,eta_tt,eta_b,bk_sm,bd_sm,bb_sm(2)</code>	
B_{SM}^{VLL}	<code>bk_sm = 0.75</code>
η_{cc}	<code>eta_cc = 1.44</code>
η_{ct}	<code>eta_ct = 0.47</code>
η_{tt}	<code>eta_tt = 0.57</code>
Details of calculations:	<code>Ref. [81, 13]</code>

6.10 $\bar{D}^0 D^0$ meson mass difference

Calculations of the mass difference Δm_D of the neutral D mesons have large theoretical uncertainties due to unknown long-distance strong corrections. Thus, as in the case of Δm_K , the `SUSY_FLAVOR` result for Δm_D should be treated as an order of magnitude estimate only.

The structure of strong corrections is analogous to those in the K meson system. However, in this case hadronic matrix elements and QCD evolution calculations available in the literature are much less refined. `SUSY_FLAVOR` uses the NLO evolution for the “ η ” factors and sets, by default, all the relevant hadronic matrix elements $B_i = 1$, i.e. it uses the “vacuum saturation” approximation (this can be changed easily when new results become available).

Routine	<code>subroutine uu_bmeson(delta_md)</code>
Input	<code>none</code>
Output	<code>delta_md = ΔM_D mass difference</code>
QCD related factors:	
	<code>common/meson_data/dmk, amk, epsk, fk, dmd, amd, fd, amb(2), dmb(2), gam_b(2), fb(2)</code>
M_D	<code>amd = 1.8645</code>
Measured ΔM_D^{exp}	<code>dmd = $4.61 \cdot 10^{-14}$</code>
f_D	<code>fd = 0.2</code>
	<code>common/bx_4q/bk(5), bd(5), bb(2,5), amu_k, amu_d, amu_b</code>
$B_1^{VLL}(\mu_D)$	<code>bd(1) = 1</code>
$B_1^{SLL}(\mu_D)$	<code>bd(2) = 1</code>
$B_2^{SLL}(\mu_D)$	<code>bd(3) = 1</code>
$B_1^{LR}(\mu_D)$	<code>bd(4) = 1</code>
$B_2^{LR}(\mu_D)$	<code>bd(5) = 1</code>
Renormalization scale μ_D	<code>amu_d = 2</code>
	<code>common/sm_4q/eta_cc, eta_ct, eta_tt, eta_b, bk_sm, bd_sm, bb_sm(2)</code>
B_{SM}^{VLL}	<code>bd_sm = 1</code>
Details of calculations:	Performed by authors, unpublished

6.11 $\bar{B}_d^0 B_d^0$ and $\bar{B}_s^0 B_s^0$ meson mixing parameters

Mixing and CP violation phenomena are also observed in the neutral B meson systems. In particular, the mass differences in the $\bar{B}_d^0 B_d^0$ and $\bar{B}_s^0 B_s^0$ oscillations have been measured,

$$\Delta M_{B_{d(s)}} = 2 \left| \langle \bar{B}_{d(s)}^0 | H_{\text{eff}}^{\Delta B=2} | B_{d(s)}^0 \rangle \right|. \quad (26)$$

The time-dependent CP asymmetry in $B_d \rightarrow J/\psi K_s$ decays, $a_{J/\psi K_s} = \sin 2\beta_{eff} \sin \Delta M_{B_d} t$, is also measured. It can be related to the argument of the $\Delta F = 2$ hadronic matrix element:

$$2\beta_{eff} = \text{Arg} \left[\langle \bar{B}_d^0 | H_{\text{eff}}^{\Delta B=2} | B_{d(s)}^0 \rangle \right]. \quad (27)$$

As experimental definitions of CP asymmetries are often convention-dependent, **SUSY_FLAVOR** gives as a more universal output directly real and imaginary parts of the $\Delta F = 2$ matrix element, which can be further used in various asymmetry calculations.

In addition to the MSSM parameters, theoretical calculations of Δm_{B_d} and Δm_{B_s} depend, as for K and D oscillations, on the relevant hadronic matrix elements and QCD evolution factors. The formulae for $\bar{B}^0 B^0$ mixing can be obtained by making the obvious replacements in the formulae presented in Sec. 6.9. Currently **SUSY_FLAVOR** uses the same set of B_i factors for both the B_d and B_s sectors, but it leaves the possibility to distinguish between them in future, if necessary. For this one needs to independently initialize the arrays **bb(1,i)** (B_d meson hadronic matrix elements) and **bb(2,i)** (B_s meson hadronic matrix elements) stored in **common/bx_4q/**.

The values of the B meson masses and coupling constants are the same as those listed in Sec. 6.6. $\Delta M_{B_{d(s)}}$ is calculated by:

Routine	subroutine dd_bmeson(i,delta_mb,dmb_re,dmb_im)
Input	<i>i</i> = 1, 2 - generation index of the lighter valence quark in the B^0 meson, i.e. <i>i</i> = 2 chooses B_s^0 and <i>i</i> = 1 chooses B_d^0 .
Output	$\text{delta_mb} = \Delta m_{B_d}(\Delta m_{B_s})$ for <i>i</i> = 1(2) $\text{dmb_re} = \text{Re}[\langle \bar{B}_{d(s)}^0 H_{\text{eff}}^{\Delta B=2} B_{d(s)}^0 \rangle]$ for <i>i</i> = 1(2) $\text{dmb_im} = \text{Im}[\langle \bar{B}_{d(s)}^0 H_{\text{eff}}^{\Delta B=2} B_{d(s)}^0 \rangle]$ for <i>i</i> = 1(2)

QCD related factors:

common/meson_data/dmk, amk, epsk, fk, dmd, amd, fd, amb(2), dmb(2), gam_b(2), fb(2)

Measured $\Delta M_{B_d}^{exp}$	$dmb(1) = 3.337 \cdot 10^{-13}$
Measured $\Delta M_{B_s}^{exp}$	$dmb(2) = 1.17 \cdot 10^{-11}$
Measured width $\Gamma_{B_d}^{exp}$	$gam_b(1) = 1.53 \cdot 10^{-12}$
Measured width $\Gamma_{B_s}^{exp}$	$gam_b(1) = 1.466 \cdot 10^{-12}$

common/bx_4q/bk(5), bd(5), bb(2,5), amu_k, amu_d, amu_b

$B_1^{\text{VLL}}(\mu_B)$	$bb(1,1) = bb(2,1) = 0.87$
$B_1^{\text{SLL}}(\mu_B)$	$bb(1,2) = bb(2,2) = 0.8$
$B_2^{\text{SLL}}(\mu_B)$	$bb(1,3) = bb(2,3) = 0.71$
$B_1^{\text{LR}}(\mu_B)$	$bb(1,4) = bb(2,4) = 1.71$
$B_2^{\text{LR}}(\mu_B)$	$bb(1,5) = bb(2,5) = 1.16$

Renormalization scale μ_B $amu_b = 4.6$

common/sm_4q/eta_cc, eta_ct, eta_tt, eta_b, bk_sm, bd_sm, bb_sm(2)

$B_{SM B_d}^{\text{VLL}}$	$bb_sm(1) = 1.18$
$B_{SM B_s}^{\text{VLL}}$	$bb_sm(2) = 1.22$
η_b	$eta_b = 0.55$

Details of calculations: Ref. [13]

Block name	Block content
SFLAV_CONTROL	SUSY_FLAVOR control variables and error code status
SFLAV_MASS	full mass spectrum of the MSSM particles after mass matrix diagonalization
SFLAV_CHIRAL_YUKAWA	Relative size of resummed chiral corrections to Yukawa couplings
SFLAV_CHIRAL_CKM	Relative size of resummed chiral corrections to CKM matrix
SFLAV_DELTA_F0	Observables related to $\Delta F = 0$ processes (EDM, $g - 2$ anomaly)
SFLAV_DELTA_F1	Observables related to $\Delta F = 1$ processes ($l \rightarrow l'\gamma$, $K \rightarrow \pi\bar{\nu}\nu$, $B_u \rightarrow \tau\nu$, $B_u \rightarrow D\tau\nu$, $B \rightarrow X_s\gamma$, $B_{d,s} \rightarrow l_i^+l_j^-$)
SFLAV_DELTA_F2	Observables related to $\Delta F = 2$ processes (ϵ_K , Δm_K , Δm_D , Δm_{B_d} , Δm_{B_s})

Table 7: Block structure of `susy_flavor.out` file.

7 SUSY_FLAVOR output

SUSY_FLAVOR output is written to the file named `susy_flavor.out`. It has the “SLHA-like” structure, i.e. it is split into “data blocks”, however those blocks are SUSY_FLAVOR specific and does not follow commonly accepted SLHA2 standards. Output file of SUSY_FLAVOR v.210 contains the data Blocks listed in Table 7.

First four blocks in `susy_flavor.out` are included for control and test purposes. Block `SFLAV_CONTROL` lists state of control variables defining conventions used for input parameters, in particular dimension of sfermion flavour violating parameters. Block `SFLAV_MASS` contain a full list of MSSM particle masses - mass eigenstates of sleptons, squarks, neutralinos and charginos, physical Higgs boson masses⁵ and, for completeness, pole lepton masses and running quark masses at m_t scale. Blocks `SFLAV_CHIRAL_YUKAWA` and `SFLAV_CHIRAL_CKM` show a relative difference of bare vs. physical Yukawa couplings and CKM matrix elements after resummation of chiral corrections - if they are too large, $\geq \mathcal{O}(1)$, the remaining program output cannot be considered to be fully reliable.

Finally, entries of Blocks `SFLAV_DELTA_F0`, `SFLAV_DELTA_F1` and `SFLAV_DELTA_F2` contain values of the flavor and CP-violating observables given in Table 1.

⁵Only the rough 1-loop Effective Potential estimate of the neutral CP-even Higgs masses is given. For precise calculations of the neutral Higgs masses in the MSSM other public SUSY generators should be used.

8 Summary and Outlook

We have presented **SUSY_FLAVOR** v2.1, a tool for calculating important flavor observables in the general R -parity conserving MSSM. Version 2 of **SUSY_FLAVOR** is capable of calculating:

- Electric dipole moments of the leptons and the neutron.
- Supersymmetric contributions to anomalous magnetic moments $g - 2$ of leptons.
- Radiative lepton decays ($\mu \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma, e\gamma$).
- Rare Kaon decays ($K_L^0 \rightarrow \pi^0\bar{\nu}\nu$ and $K^+ \rightarrow \pi^+\bar{\nu}\nu$).
- Leptonic B decays ($B_{s,d} \rightarrow l^+l^-$, $B \rightarrow \tau\nu$ and $B \rightarrow D\tau\nu$).
- Radiative B decays ($B \rightarrow \bar{X}_s\gamma$).
- $\Delta F = 2$ processes ($\bar{K}^0 - K^0$, $\bar{D} - D$, $\bar{B}_d - B_d$ and $\bar{B}_s - B_s$ mixing).

All implemented physical observables can be calculated simultaneously for a given set of MSSM input parameters. The calculation of the SUSY tree-level particle spectrum and flavor mixing matrices are performed exactly, so the code can be used for a completely general pattern of soft SUSY breaking terms (including complex phases), without restrictions on the size of the off-diagonal elements in the sfermion mass matrices. Program is written in FORTRAN 77 and runs fairly quickly; it is capable of producing a reasonably wide-range scan over the MSSM parameters within hours or days on a typical personal computer.

As a new feature not present in version 1, **SUSY_FLAVOR** v2 implements the resummation of chirally enhanced corrections (stemming from large values of $\tan\beta$ and/or large trilinear A -terms) using the systematic method developed in [17]. Such corrections modify the effective couplings of supersymmetric particles and charged Higgs bosons and generate enhanced flavor-changing neutral Higgs couplings, the latter giving significant contributions to various amplitudes coming from Higgs-penguin type diagrams. Thus, **SUSY_FLAVOR** v2 is valid for the whole parameter space of the general R -parity conserving MSSM, a unique feature currently not shared by other publicly available programs calculating FCNC and CP violation in supersymmetric models.

Besides complete routines for calculating the physical observables, **SUSY_FLAVOR** v2 also provides an extensive library of parton-level Green's functions and Wilson coefficients of many effective quark and lepton operators (see Table 3). This set actually contains many more amplitudes than necessary to compute the quantities listed in Table 1. These intermediate building blocks can be used by **SUSY_FLAVOR** users to calculate observables related to additional processes, beyond those already fully implemented, by dressing appropriate combinations of available form factors in QCD corrections and hadronic matrix elements, without repeating tedious SUSY loop calculations. For instance, the form

factors implemented in **SUSY_FLAVOR** for the analysis of $B \rightarrow X_s\gamma$ and $B_{d(s)} \rightarrow l^+l^-$ decays [6, 15] are sufficient to also calculate the $B \rightarrow Kl^+l^-$ decay rate.

The **SUSY_FLAVOR** library is an open project. We want to gradually add more features in future versions. In particular, we plan to:

- add more observables in the B -meson system, like the CP asymmetries in $B \rightarrow X_s\gamma$ decay, observables associated with $B \rightarrow Kl^+l^-$ decay and others.
- include quantities related to FCNCs in the top sector, like $t \rightarrow cX$ with $X = \gamma, Z, g, H$, in order to probe the flavor violation in up-squark mass matrices that are weakly constrained at the moment.
- include the effects of massive neutrinos.

With the increasing accuracy of experimental data on flavor and CP violation in rare processes, it may eventually become possible to not only constrain the MSSM parameters, but also, if significant deviations from the SM predictions are found, to recover their actual values. For that multi-process analysis, such as the one performed by **SUSY_FLAVOR**, will be necessary. Therefore, we hope that **SUSY_FLAVOR** becomes an important tool that is useful not only to theorists working on MSSM but also to experimentalists fitting the MSSM onto current and forthcoming data from the Tevatron, LHC, and B -factories.

Acknowledgments

The authors thank A. Buras, T. Ewerth, L. Hofer, M. Misiak, C. Savoy, Ł. Sławianowska and S. Pokorski for collaboration in performing theoretical calculations used in **SUSY_FLAVOR** and for helping to check and debug some of its sections. We would also like to thank W. Altmannshofer, D. Straub, S. Frank, D. Guadagnoli, W. Porod, M. Wick, J. Berger and D. Ghosh for checking various parts of the **SUSY_FLAVOR** code and reporting bugs or inconsistencies.

This work has received funding from the EU Seventh Framework Programme under grant agreement PITN-GA-2009-237920 (2009-2013). A.C. is supported by the Swiss National Science Foundation. The Albert Einstein Center for Fundamental Physics is supported by the “Innovations- und Kooperationsprojekt C-13 of the Schweizerische Universitätskonferenz SUK/CRUS”. The work by J.R. is supported in part by National Science Centre under research grant DEC-2011/01/M/ST2/02466 (12.2011-12.2014). The A.D. research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) - Research Funding Program: THALIS. Investing in the society of knowledge through the European Social

Fund. S.J. was supported by the Science and Technology Facilities Council [grant number ST/H004661/1] and acknowledges support from the NExT institute and SEPnet. P.T. is supported by the Paul and Daisy Soros foundation and the U.S. National Science Foundation through grant PHY-0757868.

A Installation of the program

The installation and execution of **SUSY_FLAVOR** is very simple. On Unix or Linux systems, just follow these steps :

1. Download the latest version of the code from http://www.fuw.edu.pl/susy_flavor and unpack it.
2. Change directory into **susy_flavor**.
3. Edit **Makefile** and change **F77 = gfortran** and **FOPT = -O -fno-automatic -Wall** into your compiler name and options, respectively.
4. To use the **susy_flavor_file.f** driver, reading input data from **susy_flavor.in** file, type **make sfile** (or simply **make**). To use the **susy_flavor_prog.f** driver, where input data are initialized directly inside the FORTRAN code, type **make sprog**.
5. If everything goes through, the code output is written to the file **susy_flavor.out**.
6. To run the code from now on just type **./sfile** or **./sprog**.

The authors tested **SUSY_FLAVOR** on Linux machines. With few straightforward modifications the procedure describe above can be adapted to install program on other systems.

A sample set of input parameters and corresponding **SUSY_FLAVOR** output are listed in the following appendices.

B Example of the SUSY_FLAVOR initialization sequence

Below we present the contents of **susy_flavor_file.f** and **susy_flavor_prog.f**, the master driver files for the **SUSY_FLAVOR** library. They illustrate the correct initialization sequence for all relevant MSSM parameters (see Sec. 4) and also show how to perform separate calls to the routines calculating physical observables (Sec. 6).

Driver program **susy_flavor_file.f**, initializing MSSM parameters from the input file **susy_flavor.in** is compact and simple:

```

program susy_flavor_file
program susy_flavor_header
implicit double precision (a-h,o-z)
call sflav_input(ilev,ierr)           ! parameters read from susy_flavor.in
if (ierr.ne.0) write(*,*) 'Error in parameter initialization!'
call set_resummation_level(ilev,ierr)! resummation of chiral corrections
if (ierr.ne.0) write(*,*) ierr,'Error in chiral corrections resummation!'
call susy_flavor                      ! main routine calculating physical observables
call sflav_output(ilev,ierr)          ! output written to susy_flavor.out
end

```

Driver `susy_flavor_prog.f` is longer and more complicated as all parameters has to be specified inside the code. Using this diver, flavor violating entries of sfermion mass matrices has to be given as dimensionless mass insertions.

```

program susy_flavor_prog
implicit double precision (a-h,o-z)
dimension sll(3),slr(3),amsq(3),amsu(3),amsd(3)
double complex slmi_l(3),slmi_r(3),slmi_lr(3,3),slmi_lrp(3,3)
double complex sqmi_l(3),sdmi_r(3),sumi_r(3)
double complex sdmi_lr(3,3),sumi_lr(3,3)
double complex sdmi_lrp(3,3),sumi_lrp(3,3)
double complex amg,amgg,amue
common/sf_cont/eps,indx(3,3),iconv

c   Input convention choice:
iconv = 1                           ! SLHA2 input conventions
c   iconv = 2                          ! hep-ph/9511250 input conventions

c   fixes the treatment of enhanced chiral correction resummation
c   ilev = 0                           ! no resummation, SUSY corrections strictly 1-loop
c   ilev = 1                           ! resummation using the decoupling limit
c   ilev = 2                           ! exact iterative solution, may not always converge

c   SM basic input initialization
zm0 = 91.1876d0                      ! M_Z
wm0 = 80.398d0                        ! M_W
alpha_z = 1/127.934d0                  ! alpha_em(M_Z)
st2_new = 0.232d0                      ! s_W^2(MSBar)
call vpar_update(zm0,wm0,alpha_z,st2_new)

c   QCD parameters
alpha_s = 0.1172d0                     ! alpha_s(MZ)
call lam_fit(alpha_s)                  ! fits Lambda_QCD at 3 loop level

```

```

call lam_fit_nlo(alpha_s)           ! fits Lambda_QCD at NLO level

c CKM matrix initialization
alam = 0.2258d0                   ! lambda
apar = 0.808d0                     ! A
rhobar = 0.177d0                   ! rho bar
etabar = 0.360d0                   ! eta bar
call ckm_wolf(alam,apar,rhobar,etabar)

c Fermion mass initialization, input: MSbar running quark masses
top_scale = 163.1d0
top = 163.1d0                      ! m_t(top_scale)
bot_scale = 4.18d0
bot = 4.18d0                        ! m_b(bot_scale)
call init_fermion_sector(top,top_scale,bot,bot_scale)

c Higgs sector parameters
pm = 200                            ! M_A
tanbe = 4                            ! tan(beta)
amue = (200.d0,000.d0)               ! mu
call init_higgs_sector(pm,tanbe,amue,ierr)
if (ierr.ne.0) stop 'negative tree level Higgs mass^2?',

c Gaugino sector parameters. CAUTION: if M1 is set to 0 here then
c program sets M1 and M2 GUT-related, i.e. M1 = 5/3 s_W^2/c_W^2*M2
amgg = (200.d0,0.d0)                 ! M1 (bino mass)
amg = (300.d0,0.d0)                  ! M2 (wino mass)
amglu = 600                           ! M3 (gluino mass)
call init_ino_sector(amgg,amg,amglu,amue,tanbe,ierr)
if (ierr.ne.0) write(*,*) '-ino mass below M_Z/2?'

c Slepton diagonal soft breaking parameters
sll(1) = 300.d0                      ! left selectron mass scale
sll(2) = 300.d0                      ! left smuon mass scale
sll(3) = 300.d0                      ! left stau mass scale
slr(1) = 300.d0                      ! right selectron mass scale
slr(2) = 300.d0                      ! right smuon mass scale
slr(3) = 300.d0                      ! right stau mass scale
c Slepton LL and RR mass insertions (hermitian matrices)
c slmi_x(1),slmi_x(2), slmi_x(3) are 12,23,31 entry respectively
do i=1,3
  slmi_l(i) = dcplx(0.d0,0.d0) ! slepton LL mass insertion
  slmi_r(i) = dcplx(0.d0,0.d0) ! slepton RR mass insertion

```

```

    end do
    slmi_l(1) = (2.d-2,3.d-2)      ! example, non-vanishing LL 12 entry
c   Slepton LR mass insertions, non-hermitian in general
c   All entries dimensionless (normalized to diagonal masses)
    do i=1,3
        do j=1,3
c   holomorphic LR mixing terms
            slmi_lr(i,j) = (0.d0,0.d0)
c   non-holomorphic LR mixing terms
            slmi_lrp(i,j) = (0.d0,0.d0)
        end do
    end do
c   Example: diagonal entries normalized to Y_l as in SUGRA
    slmi_lr(1,1) = (1.d-4,0.d0)    ! A_e
    slmi_lr(2,2) = (1.0d-2,0.d0)    ! A_mu
    slmi_lr(3,3) = (1.0d-1,0.d0)    ! A_tau
    slmi_lr(2,3) = (2.d-2,1.d-2)    ! example, non-vanishing LR 23 entry
c   Calculate physical masses and mixing angles
    call init_slepton_sector(sll,slr,slmi_l,slmi_r,slmi_lr,slmi_lrp
$      ,ierr)
    if (ierr.ne.0) stop 'negative tree level slepton mass^2?',

c   Squark diagonal soft breaking parameters
    amsq(1) = 500.d0                ! left squark mass, 1st generation
    amsq(2) = 450.d0                ! left squark mass, 2nd generation
    amsq(3) = 400.d0                ! left squark mass, 3rd generation
    amsd(1) = 550.d0                ! right down squark mass
    amsd(2) = 550.d0                ! right strange squark mass
    amsd(3) = 300.d0                ! right sbottom mass
    amsu(1) = 450.d0                ! right up squark mass
    amsu(2) = 450.d0                ! right charm squark mass
    amsu(3) = 200.d0                ! right stop mass
c   Squark LL and RR mass insertions (hermitian matrices)
c   sqmi_l(1),sqmi_l(2), sqmi_l(3) are 12,23,31 entry respectively, etc.
    do i=1,3
        sqmi_l(i) = (0.d0,0.d0)      ! squark LL mass insertion
        sumi_r(i) = (0.d0,0.d0)      ! up-squark RR mass insertion
        sdmi_r(i) = (0.d0,0.d0)      ! down-squark RR mass insertion
    end do
    sqmi_l(2) = (2.d-2,1.d-2)      ! example, non-vanishing LL 23 entry
c   Squark holomorphic LR mass insertions, non-hermitian in general
c   All entries dimensionless (normalized to masses)
    do i=1,3

```

```

        do j=1,3
c      holomorphic LR mixing terms
        sumi_lr(i,j) = (0.d0,0.d0) ! up-squark
        sdmi_lr(i,j) = (0.d0,0.d0) ! down-squark
c      non-holomorphic LR mixing terms
        sumi_lrp(i,j) = (0.d0,0.d0) ! up-squark
        sdmi_lrp(i,j) = (0.d0,0.d0) ! down-squark
        end do
    end do
c      Example: diagonal entries normalized to Y_d,Y_u as in SUGRA
        sumi_lr(1,1) = dcmplx(1.d-5,0.d0)
        sumi_lr(2,2) = dcplx(4.d-3,0.d0)
        sumi_lr(3,3) = dcplx(1.d0,0.d0)
        sdmi_lr(1,1) = dcplx(-1.d-3,0.d0)
        sdmi_lr(2,2) = dcplx(-2.d-2,0.d0)
        sdmi_lr(3,3) = dcplx(-8.d-1,0.d0)
        sdmi_lr(2,3) = (1.d-2,-1.d-2) ! example, non-vanishing down LR 23 entry
c      Calculate physical masses and mixing angles
        call init_squark_sector(amsq,amsu,amsd,sqmi_l,sumi_r,sdmi_r,
$          sumi_lr,sdmi_lr,sumi_lrp,sdmi_lrp,ierr)
        if (ierr.ne.0) stop 'negative tree level squark mass^2?'
c      reset status of physical Higgs mass after parameter changes
        call reset_phys_data
c      Neutral CP-even Higgs masses in the 1-loop Effective Potential
c      Approximation. Only real mu, A_t, A_b allowed - replaced x-|abs(x)
        call fcorr_EPA(tanbe,pm,top,abs(amue),amsq(3),amsd(3),amsu(3)
$          ,abs(sdmi_lr(3,3)),abs(sumi_lr(3,3)),ierr)
        if (ierr.ne.0) stop 'negative 1-loop EPA CP-even Higgs mass^2?'
c      !!! End of input section !!!
        call set_resummation_level(ilev,ierr)
        if (ierr.ne.0) write(*,*)ierr,'Error in chiral corrections resummation!'
        call susy_flavor                      ! main routine calculating physical observables
        call sflav_output(ilev,ierr)           ! output written to susy_flavor.out
        end

```

C Example of SUSY_FLAVOR input file

By default, the driver program `susy_flavor_file.f` reads input parameters from the file `susy_flavor.in`. Starting from v2.10, SUSY_FLAVOR should be able to directly read most

of output files defining MSSM Lagrangian parameters produced by other public SUSY generators, simply after renaming them to `susy_flavor.in`. However, as there are already many of such programs and they do not always uniformly follow SLHA2 standards, some incompatibilities may eventually occur. In such case, please send a message to program maintainer, so the problem could be removed in next versions of `SUSY_FLAVOR` library.

Below we provide an example input file defining a set of parameters equivalent to those given in the driver file presented in Appendix B.

```
# Example input of SUSY_FLAVOR in Les Houches-like format
#
Block MODSEL                      # Select model
  1   0                           # General MSSM
  3   0                           # MSSM particle content
  4   0                           # R-parity conserving MSSM
  5   2                           # CP violated
  6   3                           # Lepton and quark flavor violated
Block SOFTINP                      # Choose convention for the soft terms
#
# Block SOFTINP is optional - standard SLHA2 used if it is missing,
# i.e. convention=1, input_type=2, ilev=2. Otherwise:
#
# convention = 1(2): input parameters in SLHA2(hep-ph/9511250) conventions
# input_type = 1:
# sfermion off-diagonal terms given as dimensionless mass insertions
# LR diagonal terms given as dimensionless parameters
# input_type = 2:
# sfermion soft terms given as absolute values (default)
# ilev = 0
# no resummation of chirally enhanced corrections
# ilev = 1
# analytical resummation of chirally enhanced corrections
# in the limit v1,v2 << M_SUSY
# ilev = 2 (default)
# numerical iterative resummation of chirally enhanced corrections
# See comment in Blocks MSXIN2, TXIN below
  1   1                           # iconv (conventions, SLHA2 or hep-ph/9511250)
  2   1                           # input_type (dimension of soft mass entries)
  3   2                           # ilev (level of chiral corrections resummation)
Block SMINPUTS                      # Standard Model inputs
  1   1.279340000e+02           # alpha(- 1) SM MSbar(MZ)
  3   1.172000000e-01           # alpha_s(MZ) SM MSbar
  4   9.118760000e+01           # MZ(pole)
  5   4.180000000e+00           # mb(mb) SM MSbar
```

```

6   1.735000000e+02      # mtop(pole)
7   1.7768400000e+00      # mtau(pole)
11  5.1099891000e-04      # me(pole)
13  1.056580000e-01      # mmu(pole)
21  4.700000000e-03      # md(2 GeV) MSbar
22  2.150000000e-03      # mu(2 GeV) MSbar
23  9.350000000e-02      # ms(2 GeV) MSbar
24  1.275000000e+00      # mc(mc) MSbar
30  8.039800000e+01      # MW (pole), not a standard SLHA2 entry !!!
31  2.320000000e-01      # sW2 (MSBar), not a standard SLHA2 entry !!!
Block VCKMIN
1   2.258000000e-01      # CKM matrix
2   8.080000000e-01      # lambda
3   1.770000000e-01      # A
4   3.600000000e-01      # rho bar
# Block EXTPAR
0   -1.000000000e+00      # non-minimal input parameters, real part
1   2.000000000e+02      # input at EW scale only, cannot be modified!!!
2   3.000000000e+02      # Re(m1), U(1) gaugino mass
3   6.000000000e+02      # Re(m2), SU(2) gaugino mass
23  2.000000000e+02      # m3, SU(3) gaugino mass
25  4.000000000e+00      # Re(mu)
26  2.000000000e+02      # tan(beta)
# Block IMEXTPAR
1   0.000000000e+00      # MA
2   0.000000000e+00      # non-minimal input parameters, imaginary part
23  1.000000000e+02      # Im(m1), U(1) gaugino mass
# if abs(m1) = 0 SUSY_FLAVOR uses m1=5/3 s_W2/c_W2 m2
#
# Soft sfermion mass matrices
#
# Off-diagonal entries may be given as absolute entries or as
# dimensionless mass insertions - then real off-diagonal entries of
# SLHA2 blocks are calculated by SUSY_FLAVOR as
# M2(I,J) = (mass insertion)(I,J) sqrt(M2(I,I) M2(J,J))
# (see comments at the top of subroutine sflav_input)
#
# Below we give an example of dimensionless off-diagonal entries
#
Block MSL2IN
1 1  9.000000000e+04      # left soft slepton mass matrix, real part
2 2  9.000000000e+04      # Left slepton diagonal mass2, 1st generation
3 3  9.000000000e+04      # Left slepton diagonal mass2, 2nd generation
# Left slepton diagonal mass2, 3rd generation

```

```

1 2 2.000000000e-02          # Dimensionless left slepton mass insertion 12
2 3 0.000000000e+00          # Dimensionless left slepton mass insertion 23
1 3 0.000000000e+00          # Dimensionless left slepton mass insertion 13
Block IMMSL2IN
1 2 3.000000000e-02          # left soft slepton mass matrix, imaginary part
2 3 0.000000000e+00          # Dimensionless left slepton mass insertion 12
1 3 0.000000000e+00          # Dimensionless left slepton mass insertion 23
# Dimensionless left slepton mass insertion 13
Block MSE2IN
1 1 9.000000000e+04          # right soft slepton mass matrix, real part
2 2 9.000000000e+04          # Right selectron diagonal mass2
3 3 9.000000000e+04          # Right smuon diagonal mass2
1 2 0.000000000e+00          # Right stau diagonal mass2
# Dimensionless right slepton mass insertion 12
2 3 0.000000000e+00          # Dimensionless right slepton mass insertion 23
1 3 0.000000000e+00          # Dimensionless right slepton mass insertion 13
Block IMMSE2IN
1 2 0.000000000e+00          # right soft slepton mass matrix, imaginary part
2 3 0.000000000e+00          # Dimensionless right slepton mass insertion 12
1 3 0.000000000e+00          # Dimensionless right slepton mass insertion 23
# Dimensionless right slepton mass insertion 13
Block MSQ2IN
1 1 2.500000000e+05          # left soft squark mass matrix, real part
2 2 2.025000000e+05          # Left squark diagonal mass2, 1st generation
3 3 1.600000000e+05          # Left squark diagonal mass2, 2nd generation
1 2 0.000000000e+00          # Left squark diagonal mass2, 3rd generation
# Dimensionless left squark mass insertion 12
2 3 2.000000000e-02          # Dimensionless left squark mass insertion 23
1 3 0.000000000e+00          # Dimensionless left squark mass insertion 13
# left soft squark mass matrix, imaginary part
Block IMMSQ2IN
1 2 0.000000000e+00          # Dimensionless left squark mass insertion 12
2 3 1.000000000e-02          # Dimensionless left squark mass insertion 23
1 3 0.000000000e+00          # Dimensionless left squark mass insertion 13
Block MSU2IN
1 1 2.025000000e+05          # right soft up-squark mass matrix, real part
2 2 2.025000000e+05          # Right u-squark diagonal mass2
3 3 4.000000000e+04          # Right c-squark diagonal mass2
1 2 0.000000000e+00          # Right stop diagonal mass2
# Dimensionless right up-squark mass insertion 12
2 3 0.000000000e+00          # Dimensionless right up-squark mass insertion 23
1 3 0.000000000e+00          # Dimensionless right up-squark mass insertion 13
# right soft up-squark mass matrix, imaginary part
Block IMMSU2IN
1 2 0.000000000e+00          # Dimensionless right up-squark mass insertion 12
2 3 0.000000000e+00          # Dimensionless right up-squark mass insertion 23
1 3 0.000000000e+00          # Dimensionless right up-squark mass insertion 13
# right soft down-squark mass matrix, real part
Block MSD2IN
1 1 3.025000000e+05          # Right d-squark diagonal mass2
2 2 3.025000000e+05          # Right s-squark diagonal mass2

```

```

 3 3 9.000000000e+04          # Right sbottom diagonal mass2
 1 2 0.000000000e+00          # Dimensionless right down-squark mass insertion 12
 2 3 0.000000000e+00          # Dimensionless right down-squark mass insertion 23
 1 3 0.000000000e+00          # Dimensionless right down-squark mass insertion 13
Block IMMSD2IN
 1 2 0.000000000e+00          # right soft down-squark mass matrix, imaginary part
 2 3 0.000000000e+00          # Dimensionless right down-squark mass insertion 12
 1 3 0.000000000e+00          # Dimensionless right down-squark mass insertion 23
                                         # Dimensionless right down-squark mass insertion 13
#
# Soft sfermion trilinear mixing matrices
#
# LR mixing parameters can be given as absolute entries or as
# dimensionless diagonal A-terms and dimensionless off-diagonal mass
# insertions - see comments at the top of subroutine sflav_input
#
# Below we give an example of dimensionless ‘‘A terms’’.
#
# Entries of SLHA2 LR blocks are calculated by SUSY_FLAVOR as
# TL(I,J) = AL(I,J) (ML2(I,I)*ME2(J,J))** (1/4)
# TU(I,J) = AU(I,J) (MQ2(I,I)*MU2(J,J))** (1/4)
# TD(I,J) = AD(I,J) (MQ2(I,I)*MD2(J,J))** (1/4)
#
Block TEIN
 1 1 1.000000000e-04          # slepton trilinear mixing, real part
 2 2 1.000000000e-02          # Diagonal AL term, 1st generation
 3 3 1.000000000e-01          # Diagonal AL term, 2nd generation
 1 2 0.000000000e+00          # Diagonal AL term, 3rd generation
 2 1 0.000000000e+00          # Slepton LR mass insertion 12
 2 3 2.000000000e-02          # Slepton LR mass insertion 21
 3 2 0.000000000e+00          # Slepton LR mass insertion 23
 1 3 0.000000000e+00          # Slepton LR mass insertion 32
 3 1 0.000000000e+00          # Slepton LR mass insertion 13
                                         # Slepton LR mass insertion 31
Block IMTEIN
 1 1 0.000000000e+00          # slepton trilinear mixing, imaginary part
 2 2 0.000000000e+00          # Diagonal AL term, 1st generation
 3 3 0.000000000e+00          # Diagonal AL term, 2nd generation
 1 2 0.000000000e+00          # Diagonal AL term, 3rd generation
 2 1 0.000000000e+00          # Slepton LR mass insertion 12
 2 3 1.000000000e-02          # Slepton LR mass insertion 21
 3 2 0.000000000e+00          # Slepton LR mass insertion 23
 1 3 0.000000000e+00          # Slepton LR mass insertion 32
 3 1 0.000000000e+00          # Slepton LR mass insertion 13
                                         # Slepton LR mass insertion 31
Block TUIN
                                         # up-squark trilinear mixing, real part

```

```

1 1 1.000000000e-05          # Diagonal AU term, 1st generation
2 2 4.000000000e-03          # Diagonal AU term, 2nd generation
3 3 1.000000000e+00          # Diagonal AU term, 3rd generation
1 2 0.000000000e+00          # Up-squark LR mass insertion 12
2 1 0.000000000e+00          # Up-squark LR mass insertion 21
2 3 0.000000000e+00          # Up-squark LR mass insertion 23
3 2 0.000000000e+00          # Up-squark LR mass insertion 32
1 3 0.000000000e+00          # Up-squark LR mass insertion 13
3 1 0.000000000e+00          # Up-squark LR mass insertion 31
Block IMTUIIN
1 1 0.000000000e+00          # up-squark trilinear mixing, imaginary part
2 2 0.000000000e+00          # Diagonal AU term, 1st generation
3 3 0.000000000e+00          # Diagonal AU term, 2nd generation
1 2 0.000000000e+00          # Diagonal AU term, 3rd generation
# Up-squark LR mass insertion 12
# Up-squark LR mass insertion 21
# Up-squark LR mass insertion 23
# Up-squark LR mass insertion 32
# Up-squark LR mass insertion 13
# Up-squark LR mass insertion 31
Block TDIN
1 1 -1.000000000e-03         # down-squark trilinear mixing, real part
2 2 -2.000000000e-02         # Diagonal AD term, 1st generation
3 3 -8.000000000e-01         # Diagonal AD term, 2nd generation
1 2 0.000000000e+00          # Diagonal AD term, 3rd generation
# Down-squark LR mass insertion 12
# Down-squark LR mass insertion 21
# Down-squark LR mass insertion 23
# Down-squark LR mass insertion 32
# Down-squark LR mass insertion 13
# Down-squark LR mass insertion 31
Block IMTDIN
1 1 0.000000000e+00          # down-squark trilinear mixing, imaginary part
2 2 0.000000000e+00          # Diagonal AD term, 1st generation
3 3 0.000000000e+00          # Diagonal AD term, 2nd generation
1 2 0.000000000e+00          # Diagonal AD term, 3rd generation
# Down-squark LR mass insertion 12
# Down-squark LR mass insertion 21
# Down-squark LR mass insertion 23
# Down-squark LR mass insertion 32
# Down-squark LR mass insertion 13
# Down-squark LR mass insertion 31
#
# ``Non-holomorphic'' soft sfermion trilinear mixing matrices (optional)
# Such couplings are not SLHA2-standard and set to 0 if not explicitly
# defined in the input file

```

```

#
# again LR mixing parameters can be given as absolute entries or as
# dimensionless diagonal A-terms and dimensionless off-diagonal mass
# insertions - see comments at the top of subroutine sflav_input
#
# Below we give an example of dimensionless "A terms".
#
# Entries of non-holomorphic LR blocks are calculated by SUSY_FLAVOR as
# TLNH(I,J) = ALNH(I,J) (ML2(I,I)*ME2(J,J))**(1/4)
# TUNH(I,J) = AUNH(I,J) (MQ2(I,I)*MU2(J,J))**(1/4)
# TDNH(I,J) = ADNH(I,J) (MQ2(I,I)*MD2(J,J))**(1/4)
#
Block TEINH
    1 1  0.000000000e-00          # slepton trilinear mixing, real part
    2 2  0.000000000e-00          # Diagonal ALNH term, 1st generation
    3 3  0.000000000e-00          # Diagonal ALNH term, 2nd generation
    1 2  0.000000000e+00          # Diagonal ALNH term, 3rd generation
    2 1  0.000000000e+00          # Slepton LRNH mass insertion 12
    2 3  0.000000000e-00          # Slepton LRNH mass insertion 21
    3 2  0.000000000e+00          # Slepton LRNH mass insertion 23
    1 3  0.000000000e+00          # Slepton LRNH mass insertion 32
    3 1  0.000000000e+00          # Slepton LRNH mass insertion 13
                                         # Slepton LRNH mass insertion 31
Block IMTEINH
    1 1  0.000000000e+00          # slepton trilinear mixing, imaginary part
    2 2  0.000000000e+00          # Diagonal ALNH term, 1st generation
    3 3  0.000000000e+00          # Diagonal ALNH term, 2nd generation
    1 2  0.000000000e+00          # Diagonal ALNH term, 3rd generation
    2 1  0.000000000e+00          # Slepton LRNH mass insertion 12
    2 3  0.000000000e-00          # Slepton LRNH mass insertion 21
    3 2  0.000000000e+00          # Slepton LRNH mass insertion 23
    1 3  0.000000000e+00          # Slepton LRNH mass insertion 32
    3 1  0.000000000e+00          # Slepton LRNH mass insertion 13
                                         # Slepton LRNH mass insertion 31
Block TUINH
    1 1  0.000000000e-00          # up-squark trilinear mixing, real part
    2 2  0.000000000e-00          # Diagonal AUNH term, 1st generation
    3 3  0.000000000e+00          # Diagonal AUNH term, 2nd generation
    1 2  0.000000000e+00          # Diagonal AUNH term, 3rd generation
    2 1  0.000000000e+00          # Up-squark LRNH mass insertion 12
    2 3  0.000000000e-00          # Up-squark LRNH mass insertion 21
    3 2  0.000000000e+00          # Up-squark LRNH mass insertion 23
    1 3  0.000000000e+00          # Up-squark LRNH mass insertion 32
    3 1  0.000000000e+00          # Up-squark LRNH mass insertion 13
                                         # Up-squark LRNH mass insertion 31
Block IMTUINH
                                         # up-squark trilinear mixing, imaginary part

```

```

1 1 0.000000000e+00          # Diagonal AUNH term, 1st generation
2 2 0.000000000e+00          # Diagonal AUNH term, 2nd generation
3 3 0.000000000e+00          # Diagonal AUNH term, 3rd generation
1 2 0.000000000e+00          # Up-squark LRNH mass insertion 12
2 1 0.000000000e+00          # Up-squark LRNH mass insertion 21
2 3 0.000000000e-00          # Up-squark LRNH mass insertion 23
3 2 0.000000000e+00          # Up-squark LRNH mass insertion 32
1 3 0.000000000e+00          # Up-squark LRNH mass insertion 13
3 1 0.000000000e+00          # Up-squark LRNH mass insertion 31
Block TDINH
1 1 0.000000000e-00          # down-squark trilinear mixing, real part
2 2 0.000000000e-00          # Diagonal ADNH term, 1st generation
3 3 0.000000000e-00          # Diagonal ADNH term, 2nd generation
1 2 0.000000000e+00          # Diagonal ADNH term, 3rd generation
# Down-squark LRNH mass insertion 12
2 1 0.000000000e+00          # Down-squark LRNH mass insertion 21
2 3 0.000000000e+00          # Down-squark LRNH mass insertion 23
3 2 0.000000000e+00          # Down-squark LRNH mass insertion 32
1 3 0.000000000e+00          # Down-squark LRNH mass insertion 13
3 1 0.000000000e+00          # Down-squark LRNH mass insertion 31
Block IMTDINH
1 1 0.000000000e+00          # down-squark trilinear mixing, imaginary part
2 2 0.000000000e+00          # Diagonal ADNH term, 1st generation
3 3 0.000000000e+00          # Diagonal ADNH term, 2nd generation
1 2 0.000000000e+00          # Diagonal ADNH term, 3rd generation
# Down-squark LRNH mass insertion 12
2 1 0.000000000e+00          # Down-squark LRNH mass insertion 21
2 3 0.000000000e+00          # Down-squark LRNH mass insertion 23
3 2 0.000000000e+00          # Down-squark LRNH mass insertion 32
1 3 0.000000000e+00          # Down-squark LRNH mass insertion 13
3 1 0.000000000e+00          # Down-squark LRNH mass insertion 31
Block SFLAV_HADRON
1    0.1598e0                  # hadronic and QCD-related input
2    0.2e0                      # f_K
3    0.193e0                    # f_D
4    0.232e0                    # f_B_d
5    0.75e0                      # f_B_s
6    1.44e0                      # B_K for SM contribution to KKbar
7    0.47e0                      # eta_cc in KK mixing (SM)
8    0.57e0                      # eta_cc in KK mixing (SM)
9    2.e0                        # eta_cc in KK mixing (SM)
10   0.61e0                      # scale for B_K (non-SM)
11   0.76e0                      # B_K for VLL (non-SM)
12   0.51e0                      # B_K for SLL1
13   0.96e0                      # B_K for SLL2
14   0.000000000e+00            # B_K for LR1

```

```

14  1.30e0          # B_K for LR2
15  1.e0            # B_D for SM contribution
16  2.e0            # scale for B_D (non-SM)
17  1.e0            # B_D for VLL
18  1.e0            # B_D for SLL1
19  1.e0            # B_D for SLL2
20  1.e0            # B_D for LR1
21  1.e0            # B_D for LR2
22  1.18e0          # B_Bd for SM contribution
23  4.6e0            # scale for B_B (non-SM, both Bd and Bs)
24  0.87e0          # B_Bd for VLL (non-SM)
25  0.8e0            # B_Bd for SLL1
26  0.71e0          # B_Bd for SLL2
27  1.71e0          # B_Bd for LR1
28  1.16e0          # B_Bd for LR2
29  1.22e0          # B_Bs for SM contribution
30  0.55e0          # eta_b for BsBs (SM)
31  0.87e0          # B_Bs for VLL (non-SM)
32  0.8e0            # B_Bs for SLL1
33  0.71e0          # B_Bs for SLL2
34  1.71e0          # B_Bs for LR1
35  1.16e0          # B_Bs for LR2
36  1.519e-12        # Bd lifetime (experimental value)
37  1.466e-12        # Bs lifetime (experimental value)
38  5.27958e0         # Bd mass (experimental value)
39  5.36677e0         # Bs mass (experimental value)
40  3.337e-13         # Delta Bd (experimental value)
41  1.17e-11          # Delta Bs (experimental value)
42  0.497614e0        # K0 mass (experimental value)
43  3.483e-15          # Delta mK (experimental value)
44  2.229e-3           # eps_K (experimental value)
45  1.8645e0            # D0 mass (experimental value)
46  1.56e-14            # Delta mD (experimental value)
47  2.231e-10          # parameter kappa in K^0->pi^0vv calculations
48  5.173e-11          # parameter kappa in K^+->pi^+vv calculations
49  0.41e0              # parameter P_c in K->pivv calculations
50  0.013e-10          # error of kappa0
51  0.024e-11          # error of kappa+
52  0.03e0              # error of P_c
53  0.79e0              # neutron EDM_d QCD coefficient
54  -0.2e0              # neutron EDM_u QCD coefficient
55  0.59e0              # neutron CDM_d QCD coefficient
56  0.3e0              # neutron CDM_u QCD coefficient

```

```

57  3.4e0          # neutron CDM_g QCD coefficient
58  1.18e0          # neutron EDM chiral symmetry breaking scale
59  1.5e0           # pole c quark mass (in B->X_s gamma and t->cH)
60  0.1872e0        # Br(tau->e+e-)

```

D Example of SUSY_FLAVOR output

The parameters defined inside the driver program in Appendix B and in the input file listed in Appendix C should produce almost identical output, up to minor differences on distant decimal digits coming from finite accuracy of numerical computations.

We enclose content of the `susy_flavor.out` output file here, so that SUSY_FLAVOR users can check that the program gives the same result on their own computers and compilers.

```

#
# ****
# * SUSY_FLAVOR 2.10 output *
# ****
#
BLOCK SFLAV_CONTROL
 1      2          # resummation level of chiral corrections
 2      0          # error code (0 if all calculations were correct)
BLOCK SFLAV_MASS          # Mass Spectrum
#   code   mass       # particle
 24     8.039800000E+01 # W+
 25     1.053410641E+02 # h (simple 1-loop EPA approximation only)
 35     2.055718539E+02 # H (simple 1-loop EPA approximation only)
 36     2.000000000E+02 # A
 37     2.155547225E+02 # H+
 41     5.109989100E-04 # e (pole)
 42     1.056580000E-01 # mu (pole)
 43     1.776840000E+00 # tau (pole)
 44     2.608286100E-03 # md(mt) (running)
 45     5.188824475E-02 # ms(mt) (running)
 46     2.744876788E+00 # mb(mt) (running)
 47     1.193152152E-03 # mu(mt) (running)
 48     6.054455750E-01 # mc(mt) (running)
 49     1.630910000E+02 # mt(mt) (running)
1000021 6.000000000E+02 # ~g
1000022 1.608765047E+02 # ~chi_10
1000023 2.232983298E+02 # ~chi_20
1000025 2.283423917E+02 # ~chi_30
1000035 3.445964525E+02 # ~chi_40

```

```

1000024 1.879361006E+02      # ~chi_1+
1000037 3.427129063E+02      # ~chi_2+
                                # sfermion mass eigenstates
101    3.006787307E+02      # ~d(1)
102    4.036057898E+02      # ~d(2)
103    4.538544623E+02      # ~d(3)
104    5.030939992E+02      # ~d(4)
105    5.505104468E+02      # ~d(5)
106    5.505128191E+02      # ~d(6)
111    2.320752401E+02      # ~u(1)
112    4.417316304E+02      # ~u(2)
113    4.484943603E+02      # ~u(3)
114    4.487351086E+02      # ~u(4)
115    4.490302575E+02      # ~u(5)
116    4.974666710E+02      # ~u(6)
121    2.978623937E+02      # ~l(1)
122    3.017204620E+02      # ~l(2)
123    3.028213138E+02      # ~l(3)
124    3.028221696E+02      # ~l(4)
125    3.043671201E+02      # ~l(5)
126    3.085663690E+02      # ~l(6)
131    2.882496211E+02      # ~nu(1)
132    2.938268860E+02      # ~nu(2)
133    2.992957297E+02      # ~nu(3)

BLOCK SFLAV_CHIRAL_YUKAWA          # Chiral corrections size to Yukawa couplings
1     9.231173648E-03      # correction to Y_e
2     7.848366975E-03      # correction to Y_mu
3     7.331175093E-03      # correction to Y_tau
4     2.829121376E-02      # correction to Y_d
5     2.937392024E-02      # correction to Y_s
6     4.070622108E-02      # correction to Y_b
7     1.444624239E-02      # correction to Y_u
8     1.124456708E-02      # correction to Y_c
9     8.455172762E-03      # correction to Y_t

BLOCK SFLAV_CHIRAL_CKM            # Chiral corrections size to CKM matrix
1 1   3.662042898E-05      # correction to V_11
1 2   6.816725084E-04      # correction to V_12
1 3   2.096818898E-02      # correction to V_13
2 1   6.876920294E-04      # correction to V_21
2 2   2.105264221E-05      # correction to V_22
2 3   1.338999316E-02      # correction to V_23
3 1   5.993015614E-03      # correction to V_31
3 2   1.372449926E-02      # correction to V_32

```

```

 3 3      2.199454681E-05      # correction to V_33
BLOCK SFLAV_DELTA_F0
 1      -1.491542210E-25      # Delta F = 0 processes
 2      -3.072858763E-23      # EDM_e
 3      -5.158561036E-22      # EDM_mu
 4      -8.398033795E-25      # EDM_tau
 5      1.036131150E-14      # neutron EDM
 6      5.257455792E-10      # (g-2)_e/2, SUSY contribution
 7      1.575384269E-07      # (g-2)_mu/2, SUSY contribution
 8      1.575384269E-07      # (g-2)_tau/2, SUSY contribution
BLOCK SFLAV_DELTA_F1
 1      2.269341908E-08      # Br(mu-> e gamma)
 2      8.600335139E-20      # Br(tau-> e gamma)
 3      1.863540597E-08      # Br(tau-> mu gamma)
 4      2.789588524E-11      # Br(K0 -> pi0 nu nu)
 5      7.688776020E-11      # Br(K+ -> pi+ nu nu)
 6      8.769154322E-05      # BR(B_u -> tau nu)
 7      2.793594255E-01      # BR(B_u -> D tau nu)/BR(B_u -> D l nu)
 8      6.718636692E-04      # BR(B -> X_s gamma)
 9      0.000000000E+00      # BR(t -> u h) (not yet implemented)
10      0.000000000E+00      # BR(t -> c h) (not yet implemented)
11      2.647762674E-15      # BR(B_d -> e e)
12      1.131091071E-10      # BR(B_d -> mu mu)
13      2.367815464E-08      # BR(B_d -> tau tau)
14      8.146126104E-22      # BR(B_d -> mu e)
15      5.943783276E-34      # BR(B_d -> tau e)
16      6.529434197E-24      # BR(B_d -> tau mu)
17      8.806627066E-14      # BR(B_s -> e e)
18      3.762180926E-09      # BR(B_s -> mu mu)
19      7.979798794E-07      # BR(B_s -> tau tau)
20      3.677288507E-19      # BR(B_s -> mu e)
21      6.457124051E-28      # BR(B_s -> tau e)
22      3.978789529E-21      # BR(B_s -> tau mu)
BLOCK SFLAV_DELTA_F2
 1      2.453913407E-03      # epsilon_K
 2      2.508122147E-15      # Delta m_K (GeV)
 3      1.781110131E-15      # Delta m_D (GeV)
 4      3.414876336E-13      # Delta m_Bd (GeV)
 5      1.159694065E-13      # Re(H_eff_Bd)
 6      -1.253177870E-13     # Im(H_eff_Bd)
 7      1.219476743E-11      # Delta m_Bs (GeV)
 8      6.092519965E-12      # Re(H_eff_Bs)
 9      2.434925491E-13      # Im(H_eff_Bs)

```

References

- [1] T. Teubner, K. Hagiwara, R. Liao, A. D. Martin and D. Nomura, arXiv:1001.5401 [hep-ph].
- [2] G. W. Bennett *et al.* [Muon G-2 Collaboration], Phys. Rev. D **73** (2006) 072003 [hep-ex/0602035].
- [3] For reviews, see for example, H. P. Nilles, Phys. Rept. **110**, 1 (1984). H. E. Haber and G. L. Kane, Phys. Rept. **117**, 75 (1985). S. P. Martin, arXiv:hep-ph/9709356. S. Weinberg, “The quantum theory of fields. Vol. 3: Supersymmetry,” *Cambridge, UK: Univ. Pr. (2000)* 419 p
- [4] G. D’Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, Nucl. Phys. B **645** (2002) 155 [arXiv:hep-ph/0207036].
- [5] L. J. Hall, V. A. Kostelecky and S. Raby, Nucl. Phys. B **267** (1986) 415.
- [6] M. Misiak, S. Pokorski and J. Rosiek, “Supersymmetry and the FCNC effects” Adv. Ser. Direct. High Energy Phys. **15** (1998) 795 [arXiv:hep-ph/9703442].
- [7] S. Pokorski, J. Rosiek and C. A. Savoy, Nucl. Phys. B **570** (2000) 81 [arXiv:hep-ph/9906206].
- [8] J. Rosiek, Acta Phys. Polon. B **30** (1999) 3379.
- [9] A. Buras, P. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **619** (2001) 434 [arXiv:hep-ph/0107048].
- [10] J. Rosiek, arXiv:hep-ph/0108226.
- [11] A. Buras, P. Chankowski, J. Rosiek and L. Slawianowska, Phys. Lett. B **546** (2002) 96 [arXiv:hep-ph/0207241].
- [12] P. Chankowski and J. Rosiek, Acta Phys. Polon. B **33** (2002) 2329 [arXiv:hep-ph/0207242].
- [13] A. Buras, P. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **659** (2003) 3 [arXiv:hep-ph/0210145].
- [14] A. Buras, T. Ewerth, S. Jager and J. Rosiek, Nucl. Phys. B **714** (2005) 103 [arXiv:hep-ph/0408142].
- [15] A. Dedes, J. Rosiek and P. Tanedo, Phys. Rev. D **79** (2009) 055006 [arXiv:0812.4320 [hep-ph]].
- [16] J. Rosiek, arXiv:0911.3339 [hep-ph].

- [17] A. Crivellin, L. Hofer and J. Rosiek, JHEP **1107** (2011) 017 [arXiv:1103.4272 [hep-ph]].
- [18] J. Rosiek, Phys. Rev. D **41** (1990) 3464; *erratum* arXiv:hep-ph/9511250.
- [19] P. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. B **274** (1992) 191.
 P. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. B **281** (1992) 100.
 P. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. B **286** (1992) 307.
 P. Chankowski, S. Pokorski and J. Rosiek, Nucl. Phys. B **423** (1994) 437.
 P. Chankowski, S. Pokorski and J. Rosiek, Nucl. Phys. B **423** (1994) 497.
 P. Chankowski *et al.*, Nucl. Phys. B **417** (1994) 101. J. Rosiek and A. Sopczak, Phys. Lett. B **341** (1995) 419. P. Chankowski *et al.*, Nucl. Phys. Proc. Suppl. **37B** (1994) 232. V. Driesen, W. Hollik and J. Rosiek, Z. Phys. C **71** (1996) 259. A. Djouadi, V. Driesen, W. Hollik and J. Rosiek, Nucl. Phys. B **491** (1997) 68. J. Rosiek, Acta Phys. Polon. B **27** (1996) 3855. S. Heinemeyer, W. Hollik, J. Rosiek and G. Weiglein, Eur. Phys. J. C **19** (2001) 535.
- [20] J. Rosiek, P. Chankowski, A. Dedes, S. Jager and P. Tanedo, Comput. Phys. Commun. **181** (2010) 2180 [arXiv:1003.4260 [hep-ph]].
- [21] L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50** (1994) 7048 [hep-ph/9306309].
- [22] M. S. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, Nucl. Phys. B **426** (1994) 269 [hep-ph/9402253].
- [23] C. Hamzaoui, M. Pospelov and M. Toharia, Phys. Rev. D **59** (1999) 095005 [hep-ph/9807350].
- [24] M. S. Carena, D. Garcia, U. Nierste and C. E. M. Wagner, Nucl. Phys. B **577** (2000) 88 [hep-ph/9912516].
- [25] K. S. Babu and C. F. Kolda, Phys. Rev. Lett. **84** (2000) 228 [hep-ph/9909476].
- [26] G. Isidori and A. Retico, JHEP **0111** (2001) 001 [hep-ph/0110121].
- [27] G. Isidori and A. Retico, JHEP **0209** (2002) 063 [hep-ph/0208159].
- [28] A. Dedes and A. Pilaftsis, Phys. Rev. D **67** (2003) 015012 [arXiv:hep-ph/0209306].
- [29] J. Foster, K. I. Okumura and L. Roszkowski, JHEP **0508** (2005) 094.
- [30] A. Crivellin and U. Nierste, Phys. Rev. D **79** (2009) 035018 [arXiv:0810.1613 [hep-ph]].
- [31] L. Hofer, U. Nierste and D. Scherer, JHEP **0910** (2009) 081 [arXiv:0907.5408 [hep-ph]].

- [32] A. Crivellin and U. Nierste, Phys. Rev. D **81** (2010) 095007 [arXiv:0908.4404 [hep-ph]].
- [33] J. Girrbach, S. Mertens, U. Nierste and S. Wiesenfeldt, JHEP **1005** (2010) 026 [arXiv:0910.2663 [hep-ph]].
- [34] A. Crivellin and J. Girrbach, Phys. Rev. D **81** (2010) 076001 [arXiv:1002.0227 [hep-ph]].
- [35] A. Crivellin, Phys. Rev. D **83** (2011) 056001 [arXiv:1012.4840 [hep-ph]].
- [36] J. S. Lee, M. Carena, J. Ellis, A. Pilaftsis and C. E. M. Wagner, Comput. Phys. Commun. **180** (2009) 312 [arXiv:0712.2360 [hep-ph]].
- [37] G. Degrassi, P. Gambino and P. Slavich, Comput. Phys. Commun. **179**, 759 (2008) [arXiv:0712.3265 [hep-ph]].
- [38] W. Porod, Comput. Phys. Commun. **153**, 275 (2003) [arXiv:hep-ph/0301101].
- [39] F. Mahmoudi, Comput. Phys. Commun. **180**, 1579 (2009) [arXiv:0808.3144 [hep-ph]].
- [40] D. Chowdhury, R. Garani, S. K. Vempati, arXiv:1109.3551 [hep-ph].
- [41] S. Heinemeyer, W. Hollik and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000) [arXiv:hep-ph/9812320].
- [42] B. C. Allanach, Comput. Phys. Commun. **143**, 305 (2002) [arXiv:hep-ph/0104145].
- [43] A. Djouadi, J. L. Kneur and G. Moultaka, Comput. Phys. Commun. **176**, 426 (2007) [arXiv:hep-ph/0211331].
- [44] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. **177**, 894 (2007).
- [45] P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E. A. Baltz, JCAP **0407** (2004) 008 [arXiv:astro-ph/0406204].
- [46] U. Ellwanger and C. Hugonie, Comput. Phys. Commun. **175** (2006) 290 [arXiv:hep-ph/0508022].
- [47] L. J. Hall and L. Randall, Phys. Rev. Lett. **65** (1990) 2939.
- [48] R. S. Van Dyck, P. B. Schwinberg and H. G. Dehmelt, Phys. Rev. Lett. **59** (1987) 26.
- [49] A. Heister *et al.* [ALEPH Collaboration], Eur. Phys. J. C **30** (2003) 291 [hep-ex/0209066].

- [50] B. C. Regan, E. D. Commins, C. J. Schmidt and D. DeMille, Phys. Rev. Lett. **88**, 071805 (2002).
- [51] R. McNabb [Muon g-2 Collaboration], arXiv:hep-ex/0407008.
- [52] K. Nakamura *et al.* [Particle Data Group Collaboration], online update off “Review of particle physics,” J. Phys. G G **37** (2010) 075021.
- [53] C. A. Baker *et al.*, Phys. Rev. Lett. **97**, 131801 (2006) [arXiv:hep-ex/0602020].
- [54] B. Golden [MEG Collaboration], Nucl. Phys. Proc. Suppl. **218** (2011) 62.
- [55] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **104** (2010) 021802 [arXiv:0908.2381 [hep-ex]].
- [56] J. K. Ahn *et al.* [E391a Collaboration], Phys. Rev. Lett. **100**, 201802 (2008) [arXiv:0712.4164 [hep-ex]].
- [57] A. V. Artamonov *et al.* [E949 Collaboration], Phys. Rev. Lett. **101**, 191802 (2008) [arXiv:0808.2459 [hep-ex]].
- [58] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **77** (2008) 032007 [arXiv:0712.1516 [hep-ex]].
- [59] R. Aaij *et al.* [the LHCb Collaboration], Phys. Lett. B **699** (2011) 330 [arXiv:1103.2465 [hep-ex]].
“Search for the rare decay $B_s^0 \rightarrow \mu^+ \mu^-$ at the LHC with the CMS and LHCb experimentsCombination of LHC results of the search for $B_s \rightarrow \mu^+ \mu^-$ decays,” CMS-PAS-BPH-11-019.
- [60] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **96** (2006) 241802 [arXiv:hep-ex/0511015].
- [61] M. Acciarri *et al.* [L3 Collaboration], Phys. Lett. B **391** (1997) 474.
- [62] B. Aubert *et al.* [BABAR Collaboration], arXiv:0707.2758 [hep-ex].
- [63] E. Barberio *et al.* [Heavy Flavor Averaging Group], arXiv:0808.1297 [hep-ex].
- [64] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **97** (2006) 242003 [arXiv:hep-ex/0609040].
- [65] B. Allanach *et al.*, Comput. Phys. Commun. **180** (2009) 8 [arXiv:0801.0045 [hep-ph]].
- [66] A. Dedes, H. Haber and J. Rosiek, JHEP **0711** (2007) 059 [arXiv:0707.3718 [hep-ph]].
- [67] P. Z. Skands *et al.*, JHEP **0407** (2004) 036 [arXiv:hep-ph/0311123].

- [68] A. B. Lahanas and D. Wyler, Phys. Lett. B **122** (1983) 258. F. Borzumati, G. R. Farrar, N. Polonsky and S. D. Thomas, Nucl. Phys. B **555** (1999) 53. A. Crivellin, J. Girrbach and U. Nierste, Phys. Rev. D **83** (2011) 055009. A. Crivellin, L. Hofer, U. Nierste and D. Scherer, Phys. Rev. D **84** (2011) 035030.
- [69] W. Altmannshofer, A. Buras, S. Gori, P. Paradisi and D. M. Straub, Nucl. Phys. B **830** (2010) 17 [arXiv:0909.1333 [hep-ph]].
- [70] M. Cadoret, E. de Mirandes, P. Clade, S. Guellati-Khelifa, C. Schwob, F. Nez, L. Julien and F. Biraben, Phys. Rev. Lett. **101** (2008) 230801 [arXiv:0810.3152 [physics.atom-ph]].
- [71] A. Manohar, H. Georgi, *Nucl. Phys.* **B234** (1984) 189.
- [72] K. Fuyuto, J. Hisano, N. Nagata and K. Tsumura, arXiv:1308.6493 [hep-ph].
- [73] F. Mescia and C. Smith, Phys. Rev. D **76** (2007) 034017 [arXiv:0705.2025 [hep-ph]].
- [74] L. Wolfenstein, *Phys. Rev. Lett.* **51** (1983) 1945.
- [75] G. Buchalla and A. Buras, Nucl. Phys. **B548** (1999) 309.
- [76] G. Buchalla and A. Buras, Nucl. Phys. **B412** (1994) 106.
- [77] A. Buras, M. Gorbahn, U. Haisch and U. Nierste, Phys. Rev. Lett. **95** (2005) 261805 [arXiv:hep-ph/0508165]; J. Brod and M. Gorbahn, Phys. Rev. D **78** (2008) 034006 [arXiv:0805.4119 [hep-ph]]; G. Isidori, F. Mescia and C. Smith, Nucl. Phys. B **718** (2005) 319 [arXiv:hep-ph/0503107].
- [78] A. Buras, J. Girrbach, D. Guadagnoli and G. Isidori, Eur. Phys. J. C **72** (2012) 2172 [arXiv:1208.0934 [hep-ph]].
- [79] W. -S. Hou, Phys. Rev. D **48** (1993) 2342. B. Grzadkowski and W. S. Hou, Phys. Lett. B **283** (1992) 427.
- [80] Konstantin Chetyrkin, Mikolaj Misiak, Manfred Munz, Phys.Lett. **B400** (1997) 206-219; Erratum-ibid. **B425** (1998) 414 [arXiv:hep-ph/9612313].
- [81] A. Buras, S. Jager and J. Urban, Nucl. Phys. B **605** (2001) 600.
- [82] A. Buras, M. Jamin and P. Weisz, Nucl. Phys. **B347** (1990) 491; J. Urban, F. Krauss, U. Jentschura and G. Soff, Nucl. Phys. **B523** (1998) 40.
- [83] S. Herrlich and U. Nierste, Nucl. Phys. **B419** (1994) 292; S. Herrlich and U. Nierste, Phys. Rev. **D52** (1995) 6505; Phys. Rev. **D476** (1996) 27.
- [84] A. J. Buras, Les Houches Lectures “*Probing the Standard Model of Particle Interactions*”, F. David and R. Gupta, eds, Elsevier Science B.V., [arXiv:hep-ph/9806471].

PROGRAM SUMMARY

Manuscript Title: SUSY_FLAVOR v2.1: a computational tool for FCNC and CP-violating processes in the MSSM

Authors: A. Crivellin, J. Rosiek, P. Chankowski, A. Dedes, S. Jäger, P. Tanedo

Program Title: SUSY_FLAVOR v2

Journal Reference:

Catalogue identifier:

Licensing provisions: None

Programming language: Fortran 77

Operating system: Any, tested on Linux

Keywords: Supersymmetry, K physics, B physics, rare decays, CP-violation

PACS: 12.60.Jv, 13.20.He

Classification: 11.6 Phenomenological and Empirical Models and Theories

Nature of problem:

Predicting CP-violating observables, meson mixing parameters and branching ratios for set of rare processes in the general R-parity conserving MSSM.

Solution method:

We use standard quantum theoretical methods to calculate Wilson coefficients in MSSM and at one loop including QCD corrections at higher orders when this is necessary and possible. The input parameters can be read from an external file in SLHA format.

Restrictions:

The results apply only to the case of MSSM with R-parity conservation.

Unusual features:

Running time:

For single parameter set approximately 1s in double precision on a PowerBook Mac G4

References:

- [1] M. Misiak, S. Pokorski and J. Rosiek, “Supersymmetry and the FCNC effects” Adv. Ser. Direct. High Energy Phys. **15** (1998) 795 [arXiv:hep-ph/9703442].
- [2] S. Pokorski, J. Rosiek and C. A. Savoy, Nucl. Phys. B **570** (2000) 81 [arXiv:hep-ph/9906206].
- [3] A. Buras, P. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **619** (2001) 434 [arXiv:hep-ph/0107048].
- [4] A. Buras, P. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **659** (2003) 3 [arXiv:hep-ph/0210145].
- [5] A. Buras, T. Ewerth, S. Jager and J. Rosiek, Nucl. Phys. B **714** (2005) 103 [arXiv:hep-ph/0408142].
- [6] A. Dedes, J. Rosiek and P. Tanedo, Phys. Rev. D **79** (2009) 055006 [arXiv:0812.4320 [hep-ph]].
- [7] A. Crivellin, L. Hofer and J. Rosiek, JHEP **1107** (2011) 017 [arXiv:1103.4272 [hep-ph]].