

Higgs-boson reheating and frozen-in DM

Bohdan Grzadkowski

University of Warsaw



based on:

- Aqeel Ahmed, BG, Anna Socha, Phys.Lett.B 831 (2022) 137201, e-Print: 2111.06065
- Aqeel Ahmed, BG, Anna Socha, e-Print: 2207.11218

Workshop on Multi-Higgs Models 2022, Lisbon, August 31, 2022

Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM

Results

Summary

Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM

Results

Summary

- Inflation solves puzzles of the standard Big-Bang cosmology.

Motivations

- Inflation solves puzzles of the standard Big-Bang cosmology.
- Dynamics of the reheating period, which follows the inflation is often underestimated or oversimplified.

Motivations

- Inflation solves puzzles of the standard Big-Bang cosmology.
- Dynamics of the reheating period, which follows the inflation is often underestimated or oversimplified.
- It is usually assumed that the inflaton decay rate, Γ_ϕ , is constant.

Motivations

- Inflation solves puzzles of the standard Big-Bang cosmology.
- Dynamics of the reheating period, which follows the inflation is often underestimated or oversimplified.
- It is usually assumed that the inflaton decay rate, Γ_ϕ , is constant.
- Hereafter we are going to discuss relations between inflation and reheating dynamics focusing on possible interactions between the Higgs boson and inflaton.

Motivations

- Inflation solves puzzles of the standard Big-Bang cosmology.
- Dynamics of the reheating period, which follows the inflation is often underestimated or oversimplified.
- It is usually assumed that the inflaton decay rate, Γ_ϕ , is constant.
- Hereafter we are going to discuss relations between inflation and reheating dynamics focusing on possible interactions between the Higgs boson and inflaton.
- Dynamics of reheating influences the dark matter sector, especially in the context of the freeze-in DM production.

Non-instantaneous reheating




Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM

Results

Summary

The α -attractor T-model

$$\mathcal{L}_\phi = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi)$$

$$V(\phi) = \Lambda^4 \tanh^{2n} \left(\frac{|\phi|}{\sqrt{6\alpha} M_{\text{Pl}}} \right)$$

$$\simeq \begin{cases} \Lambda^4 & |\phi| \gg M_{\text{Pl}} \\ \Lambda^4 \left| \frac{\phi}{M_{\text{Pl}}} \right|^{2n} & |\phi| \ll M_{\text{Pl}} \end{cases},$$

where $n > 0$, $\sqrt{6\alpha} \lesssim 10$, $\Lambda \lesssim 1.6 \times 10^{16}$ GeV.

$$\ddot{\phi} + 3H\dot{\phi} + V_{,\phi}(\phi) = 0,$$

$H \equiv \dot{a}/a$ is the Hubble rate.





Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM


Results

Summary

Interactions

$$\mathcal{L}_{\text{DM}} = -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} + \frac{1}{2}m_X^2 X_\mu X^\mu,$$

$$\mathcal{L}_{\text{int}} = -\left\{ \boxed{g_{h\phi} M_{\text{Pl}} \phi |h|^2} + \frac{h^{\mu\nu}}{M_{\text{Pl}}} \left[T_{\mu\nu}^\phi + T_{\mu\nu}^{\text{DM}} + T_{\mu\nu}^{\text{SM}} \right] + \frac{C_X^\phi m_X^2}{2M_{\text{Pl}}} \phi X_\mu X^\mu + \frac{C_X^h m_X^2}{2M_{\text{Pl}}^2} |h|^2 X_\mu X^\mu \right\},$$



The Higgs portal

$$\mathcal{L}_{int} = g_{h\phi} M_{\text{Pl}} \phi |h|^2$$

homogeneous, classical
background field

ϕ ✗

coherently oscillating

$$\phi = \varphi(t) \cdot \mathcal{P}(t)$$

rapidly-oscillating
slowly-varying envelope, $\rho_\phi \equiv V(\varphi)$

⇒ **Reheating** i.e., energy transfer between the inflaton and the SM sector

$$\frac{1}{V} \frac{dE_g}{dt} \equiv \rho_\phi \Gamma_\phi = g_{h\phi}^2 M_{\text{Pl}}^2 \frac{\varphi^2(t)}{8\pi} \sum_{i=0}^3 \sum_{k=1}^{\infty} k\omega |\mathcal{P}_k|^2 \sqrt{1 - \left(\frac{2m_{h_i}}{k\omega}\right)^2}, \quad \mathcal{P}(t) = \sum_{k=-\infty}^{\infty} \mathcal{P}_k e^{-ik\omega t}$$

⇒ **Higgs mass** induced by the oscillating inflaton background


$$m_{h_0}^2 = g_{h\phi} M_{\text{Pl}} \varphi \begin{cases} |\mathcal{P}|, & \mathcal{P}(t) > 0 \\ 2|\mathcal{P}|, & \mathcal{P}(t) < 0 \end{cases} \quad v_h = \begin{cases} 0, & \mathcal{P}(t) > 0 \\ \sqrt{|m_{h_0}^2|/(2\lambda_h)}, & \mathcal{P}(t) < 0 \end{cases}$$

The Higgs portal


$$m_{h_0}^2 = g_{h\phi} M_{\text{Pl}} \varphi \begin{cases} |\mathcal{P}|, & \mathcal{P}(t) > 0 \\ 2|\mathcal{P}|, & \mathcal{P}(t) < 0 \end{cases}$$

$$v_h = \begin{cases} 0, & \mathcal{P}(t) > 0 \\ \sqrt{|m_{h_0}^2| / (2\lambda_h)}, & \mathcal{P}(t) < 0 \end{cases}$$

$n=2/3, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$



$n=3/2, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$



Kinematic suppression

The inflaton decay rate can be written as

$$\langle \Gamma_\phi \rangle = \frac{g_{h\phi}^2}{32\pi} \frac{M_{\text{Pl}}^2}{m_\phi(a)} \gamma_h(a),$$

effective mass

$$m_\phi(a) = V_{,\phi\phi}|_{\phi=\varphi}$$


$$\gamma_h(a) \simeq \sum_{i=0}^3 \sum_{k=1}^{\infty} k |\mathcal{P}_k|^2 \left\langle \sqrt{1 - \left(\frac{2m_{h_i}(a)}{k\omega(a)} \right)^2} \right\rangle \quad \omega \propto m_\phi$$

It turns out that

$$\gamma_h \propto a^{\frac{3(3-2n)}{1+n}}$$

which implies

$$\langle \Gamma_\phi \rangle \propto a^{\frac{6-3n}{1+n}} \equiv a^{-\beta}$$




The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = - \langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H\rho_{\text{SM}} = \langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle \rho_\phi - 2\langle E_X \rangle \mathcal{S}_{\text{SM}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3Hn_X = \mathcal{D}_\phi + \mathcal{S}_\phi + \mathcal{S}_{\text{SM}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$




The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = -\langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H \rho_{\text{SM}} = \boxed{\langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle} \rho_\phi - 2\langle E_X \rangle \mathcal{S}_{\text{SM}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3H n_X = \mathcal{D}_\phi + \mathcal{S}_\phi + \mathcal{S}_{\text{SM}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$




The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = -\langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H\rho_{\text{SM}} = \langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle \rho_\phi - 2\langle E_X \rangle \boxed{\mathcal{S}_{\text{SM}}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3Hn_X = \mathcal{D}_\phi + \mathcal{S}_\phi + \boxed{\mathcal{S}_{\text{SM}}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$




The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = -\langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H\rho_{\text{SM}} = \langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle \rho_\phi - 2\langle E_X \rangle \mathcal{S}_{\text{SM}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3Hn_X = \mathcal{D}_\phi + \mathcal{S}_\phi + \mathcal{S}_{\text{SM}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$




The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = -\langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H\rho_{\text{SM}} = \langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle \rho_\phi - 2\langle E_X \rangle \mathcal{S}_{\text{SM}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3Hn_X = \boxed{\mathcal{D}_\phi} + \mathcal{S}_\phi + \mathcal{S}_{\text{SM}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$



The time-averaged Boltzmann equations

$$\dot{\rho}_\phi + \frac{6n}{n+1} H \rho_\phi = -\langle \Gamma_\phi \rangle \rho_\phi$$

$$\dot{\rho}_{\text{SM}} + 4H\rho_{\text{SM}} = \langle \Gamma_{\phi \rightarrow \text{SM SM}} \rangle \rho_\phi - 2\langle E_X \rangle \mathcal{S}_{\text{SM}} - \langle E_{h_0} \rangle \mathcal{D}_{h_0}$$

$$\dot{n}_X + 3Hn_X = \mathcal{D}_\phi + \boxed{\mathcal{S}_\phi} + \mathcal{S}_{\text{SM}} + \mathcal{D}_{h_0}$$

with the Hubble rate $H^2 = \frac{1}{3M_{\text{Pl}}^2} (\rho_\phi + \rho_{\text{SM}} + \rho_X)$




Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM


Results

Summary


Non-instantaneous reheating

$$\rho_\phi(a) \stackrel{H \gg \Gamma_\phi}{\simeq} 3M_{\text{Pl}}^2 H_e^2 \left(\frac{a_e}{a}\right)^{3(1+\bar{w})}$$


$$\rho_{\mathcal{R}}(a) = \frac{6M_{\text{Pl}}^2 H_e \Gamma_\phi^e}{5-3\bar{w}-2\beta} \left[\left(\frac{a_e}{a}\right)^{\beta+3(1+\bar{w})/2} - \left(\frac{a_e}{a}\right)^4 \right]$$




$$g_{h\phi} = 10^{-5}, n=1, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$$



$$g_{h\phi} = 10^{-5}, n=2, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$$



$$g_{h\phi} = 10^{-5}, n=1, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$$




$$m_{h_i} = 0$$

$$\beta = 0, \rho_{\mathcal{R}} \propto a^{-3/2}$$

$$m_{h_i} \neq 0$$

$$\beta = -\frac{3}{2}, \rho_{\mathcal{R}} \propto a^0$$

$$g_{h\phi} = 10^{-5}, n=2, \alpha=1/6, \Lambda=3 \cdot 10^{-3} M_{\text{Pl}}$$




$$\beta = -1, \rho_{\mathcal{R}} \propto a^{-1}$$

$$\beta = 0, \rho_{\mathcal{R}} \propto a^{-2}$$

Gravitational DM production

$$\mathcal{L}_{\text{DM}} = -\frac{1}{4} X_{\mu\nu} X^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu X^\mu + \mathcal{L}_{\text{int}}$$

$$\mathcal{L}_{\text{int}} = \frac{h_{\mu\nu}}{M_{\text{Pl}}} \left(T_\phi^{\mu\nu} + T_X^{\mu\nu} + T_{\text{SM}}^{\mu\nu} \right)$$




$$\frac{dN_X}{da} = \frac{a^2}{H} (\mathcal{S}_\phi + \mathcal{S}_{\text{SM}})$$

where $N_X \equiv n_X a^3$

$$\Omega_X^{\text{grav}} h^2 \simeq \frac{m_X}{\rho_c} \frac{N^{\text{grav}}(a_{\text{rh}})}{a_{\text{rh}}^3} \frac{s_0}{s(a_{\text{rh}})} h^2.$$

$$\Omega_X^{\text{grav}} h^2 = \Omega_X^{(\text{obs})} h^2 = 0.1198 \pm 0.0012$$




Heavy DM particles are produced by the freeze-in from the SM sector

$$\Omega_X^{\text{grav}} h^2 \simeq \frac{m_X}{\rho_c} \frac{N^{\text{grav}}(a_{\text{rh}})}{a_{\text{rh}}^3} \frac{s_0}{s(a_{\text{rh}})} h^2.$$

$$\Omega_X^{\text{grav}} h^2 = \Omega_X^{(\text{obs})} h^2 = 0.1198 \pm 0.0012$$

For the $n=2$ case,
 $\Omega_X^{\text{grav}} h^2$ does not depend on $g_{h\phi}$

Light DM particles are dominantly produced from the inflaton



For the $n=2/3$ case,
 $\Omega_X^{\text{grav}} h^2$ does not depend on m_X for heavy DM

Table of Contents

Introduction

Inflation dynamics

The model of reheating and DM

Results

Summary

Summary

- The α -attractor T-model potential for the inflaton field has been adopted:

$$V(\phi) = \Lambda^4 \tanh^{2n} \left(\frac{|\phi|}{\sqrt{6\alpha} M_{\text{Pl}}} \right) \simeq \begin{cases} \Lambda^4 & |\phi| \gg M_{\text{Pl}} \\ \Lambda^4 \left| \frac{\phi}{M_{\text{Pl}}} \right|^{2n} & |\phi| \ll M_{\text{Pl}} \end{cases},$$

- The reheating has been triggered by

$$\mathcal{L}_{int} = g_{h\phi} M_{\text{Pl}} \phi |h|^2$$

- It has been shown that both duration of reheating and evolution of radiation energy density, $\rho_{\mathcal{R}}$, are sensitive to the shape of the inflaton potential (n).
- The role of kinematical suppression emerging from \mathcal{L}_{int} has been investigated. It has been shown that [the non-zero mass of the Higgs boson](#) leads to the elongation of the reheating period, changes the $\rho_{\mathcal{R}}(a)$ and $T(a)$ evolution, and favors reduced T_{max} .

- It has been shown that purely gravitational production of DM is possible.
- Purely gravitation reheating needs to be investigated.

Back-up slides

The α -attractor T-model


Time averaging:

$$\langle f(t) \rangle = \frac{1}{\mathcal{T}} \int_t^{t+\mathcal{T}} d\tau f(\tau),$$

Equation of state:

$$\bar{w} \equiv \frac{\langle p_\phi \rangle}{\langle \rho_\phi \rangle} = \frac{n-1}{n+1}$$

| n | $\bar{w} \equiv \frac{\langle p_\phi \rangle}{\langle \rho_\phi \rangle}$ |
|---------------|---|
| $\frac{2}{3}$ | $-\frac{1}{5}$ |
| 1 | 0 |
| $\frac{3}{2}$ | $\frac{1}{5}$ |
| 2 | $\frac{1}{3}$ |



Particle production in a classical inflaton background

For the interactions proportional to the $\phi = \varphi \cdot \mathcal{P}$ term, the lowest-order non-vanishing S-matrix element takes the form

$$S_{if}^{(1)} = \sum_k \mathcal{P}_k \langle f | \int d^4x \varphi(t) e^{-ik\omega t} \mathcal{L}_{\text{int}}(x) | i \rangle$$

where

$$|i\rangle \equiv |0\rangle, \quad |f\rangle \equiv \hat{a}_f^\dagger \hat{a}_f^\dagger |0\rangle.$$

If the **envelope** $\varphi(t)$ varies on the time-scale much longer than the time-scale relevant for processes of particle creation, the S-matrix element can be written as

$$S_{if}^{(1)} = i\varphi(t) \sum_k \mathcal{P}_k \mathcal{M}_{0 \rightarrow f}(k) \times (2\pi)^4 \delta(k\omega - 2E_f) \delta^3(p_{f_1} + p_{f_2}).$$