

Bosonic quadratic Hamiltonians

Jan Dereziński

Department of Mathematical Methods in Physics,
Faculty of Physics, University of Warsaw,
Pasteura 5,
02-093 Warszawa, Poland,
email: jan.derezinski@fuw.edu.pl

January 1, 2017

Abstract

We discuss self-adjoint operators given formally by expressions quadratic in bosonic creation and annihilation operators. We give conditions when they can be defined as self-adjoint operators, possibly after an infinite renormalization. We also discuss explicit formulas for their infimum.

Our main motivation comes from Local Quantum Field Theory, which furnishes interesting examples of bosonic quadratic Hamiltonians that require an infinite renormalization [De].

1 Introduction

Quantum bosonic quadratic Hamiltonians, or *bosonic Bogoliubov Hamiltonians* are formally given by expressions of the form

$$\hat{H} = \sum h_{ij} \hat{a}_i^* \hat{a}_j + \frac{1}{2} \sum g_{ij} \hat{a}_i^* \hat{a}_j^* + \frac{1}{2} \sum \bar{g}_{ij} \hat{a}_i \hat{a}_j + c, \quad (1.1)$$

where $h = [h_{ij}]$ is a Hermitian matrix, $g = [g_{ij}]$ is a symmetric matrix, c is an arbitrary real number (possibly, infinite!) and \hat{a}_i^*, \hat{a}_j are the usual bosonic creation/annihilation operators. They are often used in quantum field theory to describe free theories interacting with a given external classical field [IZ, De]. They are responsible for the *Casimir effect* [IZ]. Bogoliubov applied them to the theory of interacting Bose gas [Bo], which justifies the name *Bogoliubov Hamiltonians*.

Bogoliubov Hamiltonians that are bounded from below are especially useful. Their infimum $E := \inf \hat{H}$ is often interesting physically.

Bogoliubov Hamiltonians have a surprisingly rich mathematical theory. In infinite dimension this theory sometimes involves interesting pathologies. For instance, \hat{H} is often ill defined, but one can define its “infimum” E . In some situations, one needs to perform an infinite renormalization in order to define \hat{H} , or at least to compute E . This is typical for Bogoliubov Hamiltonians that are motivated by relativistic quantum field theory [De]. Another example of interesting mathematics related to Bogoliubov Hamiltonians can be found in a recent paper [NNS], which contains a beautiful proof of diagonalizability of normally ordered Bogoliubov Hamiltonians under essentially optimal conditions.

Our paper is devoted to a systematic theory of bosonic Bogoliubov Hamiltonians in an abstract setting. We do not restrict ourselves to the normally ordered case (with $c = 0$ in (1.1)). We start from a more general definition saying that a Bogoliubov Hamiltonian is the self-adjoint generator of a one-parameter unitary group on a bosonic Fock space that implements a symplectic group. There are interesting and

physically important examples where the normally ordered Bogoliubov Hamiltonian is ill defined, whereas renormalized ones exist [De].

The family of Bogoliubov Hamiltonians given by fixing h, g and varying $c \in \mathbb{R}$ (1.1) can be understood as various quantizations of a single classical quadratic Hamiltonian,

$$H = \sum h_{ij} a_i^* a_j + \frac{1}{2} \sum g_{ij} a_i^* a_j^* + \frac{1}{2} \sum \bar{g}_{ij} a_i a_j, \quad (1.2)$$

where a_i, a_j^* are classical (commuting) variables. c , which appears in (1.1), can be understood as the ambiguity of quantization due to noncommutativity of \hat{a}_i, \hat{a}_j . The most popular choice is probably $c = 0$, corresponding to the *normally (Wick) ordered Hamiltonian*. It will be denoted \hat{H}^n . The choice $c = \frac{1}{2} \sum_i h_{ii}$, which we call the *Weyl Bogoliubov Hamiltonian* and denote \hat{H}^w , has its advantages as well. In some situations, however, one needs to consider other quantizations, where the constant c may turn out to be infinite, and can be viewed as a renormalization counterterm. We describe one particular possibility, which we call \hat{H}^{ren} . In the language of Feynman diagrams \hat{H}^{ren} corresponds to discarding loops of order 2 or less, which is often implicit in quantum field theory.

We will use the following notation for the infimum of the three main Bogoliubov Hamiltonians that we discuss:

$$E^w := \inf \hat{H}^w, \quad E^n := \inf \hat{H}^n, \quad E^{\text{ren}} := \inf \hat{H}^{\text{ren}}. \quad (1.3)$$

In physics the infimum of the Hamiltonian appears under various names, eg. vacuum energy, Casimir energy, vacuum polarization, effective potential. Physicists often compute the vacuum energy without worrying whether the corresponding quantum Hamiltonian is well defined as a self-adjoint operator. Following this philosophy, we may consider E^n or E^{ren} under conditions that are more general than the conditions for the existence of the corresponding Hamiltonians.

1.1 Comparison with literature

It is not always very easy to read the literature on Bogoliubov Hamiltonians and to compare statements in various papers. Their authors often use different conventions, terminology and notations.

Most of these issues disappear when one fixes a basis in the 1-particle space, identifying it with \mathbb{C}^m . Then a Bogoliubov Hamiltonian is determined by two matrices, $h = [h_{ij}]$ and $g = [g_{ij}]$, and possibly a number c , see (1.1).

When we want to use a basis independent language, replacing \mathbb{C}^m by an abstract Hilbert space \mathcal{W} , it is clear how to interpret h —it is a self-adjoint operator on \mathcal{W} . It is less obvious how to interpret g . One possibility is to view g as a symmetric tensor, that is, an element of $\otimes_s^2 \mathcal{W}$. Often, however, it is preferable to view g as an operator from \mathbb{C}^m to \mathbb{C}^m . These two \mathbb{C}^m should be however viewed as two distinct spaces—one is the complex conjugate of the other, see eg. [De]. The notion of a complex conjugate space is somewhat subtle and has a few equivalent but superficially distinct interpretations, see Subsection A.1. Various authors prefer distinct interpretations, see eg. the footnote ⁶ in Appendix A of [HS]. (Strictly speaking, this footnote refers to the fermionic case, however the fermionic and bosonic cases are quite analogous).

When we consider an infinite dimensional space, there are additional problems: various operators are often unbounded, are not trace class, or simply do not exist.

Because of these two kinds of problems, our paper is divided into two parts. In the first part we assume that the 1-particle space is finite dimensional and has a fixed orthonormal basis. All operators are represented by matrices. We do not worry about conceptual subtleties related to antilinear maps and the complex conjugate space. Infinite renormalization is not needed and all formulas are valid with no technical restrictions.

In the second part of our paper, the 1-particle space is an abstract space \mathcal{W} of any dimension. We follow mostly the conceptual framework of [DG]. We distinguish between \mathcal{W} and its complex conjugate

$\overline{\mathcal{W}}$. We need to give technical conditions guaranteeing that various concepts and formulas survive into infinite dimension.

Throughout the paper it is assumed that the reader is familiar with mathematical formalism of 2nd quantization. Properties of the metaplectic representation in the Fock space play an important role, such as the Shale Theorem and formulas for the Bogoliubov implementers (2.17) and (2.18). These formulas were known to Friedrichs [F1], analysed later by Ruisenaars [Ru1, Ru2] and Berezin [Be]. We treat [DG] as the basic reference on this subject, where in particular various questions related to the unboundedness of bosonic creation and annihilation operators are discussed in detail.

Large parts of Sect. 2 is well known. Thm 2.3 about diagonalizability of a quadratic Hamiltonian by a positive symplectic transformation is implicitly contained in [DG] (see Thm 11.20 (3) together with Thm 18.5 (3)). We come back to this issue in the next section, where an arbitrary dimension introduces additional technical issues. Note that a similar fact proven in [NNS] does not provide a construction of a distinguished diagonalizing operator.

The basic formula for the infimum of a quadratic Hamiltonian comes from [BD]. However, some of the formulas for the infimum of the normally ordered Hamiltonians, such as (2.68), (2.69) and (2.70) seem to be new. In finite dimension they are not so interesting, however they become quite useful in infinite dimension.

It seems that the construction of the renormalized Hamiltonians described in Subsects 2.12 and 2.13 has never been presented in the literature in the abstract setting. Their importance is evident in concrete situations of Quantum Field Theory described in [De]. We give a brief discussion of the examples from QFT at the end of Introduction.

Quadratic Hamiltonians in infinite dimensions is a rather technical topic of operator theory. Therefore, we prefer to give a self-contained treatment of this subject. Many results and definitions that we present are new, however at some places we recall proofs contained in the literature.

Note that it would be awkward and restrictive to define Bogoliubov Hamiltonians in the infinite dimensional context by an expression of the form (1.1). Instead, we define them as self-adjoint generators of one parameter unitary groups implementing Bogoliubov transformations. (In the bosonic context the term “Bogoliubov transformations” is usually meant to denote “symplectic transformations”). The abstract approach makes it sometimes difficult to define some objects, since we cannot refer to a formula of the form (1.1). Fortunately, it is obvious how to define the Weyl Bogoliubov Hamiltonian—as the generator of a group inside the metaplectic group. It is less obvious how to define normally ordered Hamiltonians. The definition that we propose in Subsec. 3.7 seems to be new—in particular, it is more general from the definition of [BD].

Subsect. 3.7 and 3.8 give criteria for the existence of various quantizations. In these subsections there is no assumption on the positivity of h . On the other hand, most results require the boundedness of g . Some results in this part of the paper come from [Be] and [BD]. However, Thm 3.18 (1), which gives a convenient criterion for the implementability of classical dynamics, seems to be new. It is useful in the context of examples from QFT discussed below.

In the following subsections we adopt a different set of assumptions. In particular, we assume that h is positive and g is form bounded wrt h with bound less than 1. This condition guarantees the positivity and diagonalizability of classical Hamiltonians.

Diagonalization of Bogoliubov Hamiltonians on the quantum level was considered already by Berezin [Be], then by Bach and Bru [BB]. In a recent paper [NNS], Napiórkowski, Phan Thanh Nam and Solovej gave a new beautiful proof of diagonalizability. In our paper we repeat some of the arguments of [NNS], describing their result in Thm 3.21, giving essentially optimal conditions for diagonalization. In distinction to [NNS], we show that there exists a distinguished positive symplectic operator diagonalizing a given Bogoliubov Hamiltonian.

In Thm 3.23 we also describe a construction of normally ordered Bogoliubov Hamiltonian based on the form techniques (involving the so-called KLMN Theorem) presented in [NNS]. This is an important

improvement (even if it sounds technical) as compared to the results of [BD], which were restricted to operator type perturbations.

These theorems are complemented with new results. In Thm 3.24, we show that the dynamics generated by the normally ordered Hamiltonian implements the corresponding classical dynamics. On a formal level this theorem seems obvious, nevertheless due to the unboundedness of various operators it needs a careful proof. Another new result, easy in finite dimension and rather technical in the general case is the formula for the ground state energy described in Thm 3.29. We also discuss a criterion for the existence of the Weyl Bogoliubov Hamiltonians in Thm 3.31 and for the existence of the renormalized ground state energy in Thm 3.32.

Let us mention some topics that are left out of our paper. We do not discuss time-dependent Bogoliubov Hamiltonians, the implementability and the phase of the corresponding scattering operator. This is interesting, especially in the context of charged relativistic fields in an external electromagnetic potential. An infinite renormalization is needed in order to define the vacuum energy. This topic on a partly heuristic level is discussed in [De]. Its fermionic counterpart (a Dirac particle in an external electromagnetic potential) is better known in the literature, see eg. [DDMS].

1.2 Applications to QFT

Let us first discuss the question of naturalness of the definition of various kinds of Bogoliubov Hamiltonians.

The Weyl Hamiltonian \hat{H}^w is the most natural. In fact, it is invariant wrt symplectic transformations, see (2.31). Unfortunately, it is often ill defined.

The normally ordered Hamiltonian \hat{H}^n is naturally defined given a Fock representation. In particular, this is the case when we have a distinguished positive classical quadratic Hamiltonian which is treated as the “free” one. Then there exists a unique Fock representation where the free Hamiltonian can be quantized without any double creation/annihilation operators. It is usually quantized in the normally ordered form. We will denote it by \hat{H}_0^n .

Suppose that we are interested in the “full” Hamiltonian, which is quadratic, but more complicated than the free one and involves an interaction with external fields. We can then ask whether the corresponding classical Hamiltonian can be quantized. The most straightforward procedure seems to consider the normally ordered full quantum Hamiltonian \hat{H}^n . Then the corresponding ground state energy formally equals then the difference of the “free Weyl ground state energy” and the “full Weyl ground state energy” (in typical situations both infinite).

It sometimes happens that \hat{H}^n is ill defined as well. Then we can try to subtract from \hat{H}^n another counterterm. As examples from QFT show, the most natural possibility is to subtract the 2nd order contribution in the perturbative expansion, obtaining \hat{H}^{ren} as in Subsect. 2.12 and 2.13. Below we briefly describe two examples where one needs to perform such a renormalization. These examples are discussed in more detail in [De].

Consider the neutral massive scalar quantum field $\hat{\phi}(\vec{x})$. Its conjugate field is denoted $\hat{\pi}(\vec{x})$ with the usual equal time commutation relations

$$\begin{aligned} [\hat{\phi}(\vec{x}), \hat{\phi}(\vec{y})] &= [\hat{\pi}(\vec{x}), \hat{\pi}(\vec{y})] = 0, \\ [\hat{\phi}(\vec{x}), \hat{\pi}(\vec{y})] &= i\delta(\vec{x} - \vec{y}). \end{aligned} \tag{1.4}$$

The free Hamiltonian is defined in the standard way:

$$\hat{H}_0^n := \int : \left(\frac{1}{2} \hat{\pi}^2(\vec{x}) + \frac{1}{2} (\vec{\partial} \hat{\phi}(\vec{x}))^2 + \frac{1}{2} m^2 \hat{\phi}^2(x) \right) : d\vec{x}, \tag{1.5}$$

where the double dots denote the normal ordering.

Suppose that the mass is perturbed by a Schwartz function $\kappa(\vec{x})$. One can check that the normally ordered full Hamiltonian does not exist. However, the renormalized Hamiltonian is well-defined (see Chap. III Subsect. C14 of [De]). Formally, it can be written a

$$\hat{H}^{\text{ren}} := \int : \left(\frac{1}{2} \hat{\pi}^2(\vec{x}) + \frac{1}{2} (\vec{\partial} \hat{\phi}(\vec{x}))^2 + \frac{1}{2} (m^2 + \kappa(\vec{x})) \hat{\phi}^2(\vec{x}) \right) : d\vec{x} - E_2, \quad (1.6)$$

where the infinite counterterm E_2 is the contribution of loop diagrams with 2 vertices, see Subsect. 2.13.

The next example is more singular. Consider the charged massive scalar quantum field $\hat{\psi}(\vec{x})$, with $\hat{\psi}^*(\vec{x})$ denoting its Hermitian adjoint. The conjugate field will be denoted $\hat{\eta}(\vec{x})$, so that we have the commutation relations

$$[\hat{\psi}(\vec{x}), \hat{\psi}(\vec{y})] = [\hat{\psi}(\vec{x}), \hat{\eta}(\vec{y})] = [\hat{\eta}(\vec{x}), \hat{\eta}(\vec{y})] = 0, \quad (1.7)$$

$$[\hat{\psi}(\vec{x}), \hat{\eta}^*(\vec{y})] = [\hat{\psi}^*(\vec{x}), \hat{\eta}(\vec{y})] = i\delta(\vec{x} - \vec{y}). \quad (1.8)$$

The free Hamiltonian is of course

$$\hat{H}_0^{\text{fr}} = \int : \left(\hat{\eta}^*(\vec{x}) \hat{\eta}(\vec{x}) + \vec{\partial} \hat{\psi}^*(\vec{x}) \vec{\partial} \hat{\psi}(\vec{x}) + m^2 \hat{\psi}^*(\vec{x}) \hat{\psi}(\vec{x}) \right) : d\vec{x}.$$

Suppose now that we consider an external stationary electromagnetic potential, described by, say, Schwartz functions (A^0, \vec{A}) . Then the natural candidate for the full Hamiltonian is (see Chap. VI, Subsect. B17 of [De]):

$$\begin{aligned} \hat{H}^{\text{ren}} = & \int d\vec{x} \left(\hat{\eta}^*(\vec{x}) \hat{\eta}(\vec{x}) + ieA_0(\vec{x}) (\hat{\psi}^*(\vec{x}) \hat{\eta}(\vec{x}) - \hat{\eta}^*(\vec{x}) \hat{\psi}(\vec{x})) \right. \\ & + (\partial_i - ieA_i(\vec{x})) \hat{\psi}^*(\vec{x}) (\partial_i + ieA_i(\vec{x})) \hat{\psi}(\vec{x}) \\ & \left. + m^2 \hat{\psi}^*(\vec{x}) \hat{\psi}(\vec{x}) \right) - E_0 - E_1 - E_2, \end{aligned} \quad (1.9)$$

where E_0, E_1, E_2 are infinite counterterms, which come from the expansion described in (2.99).

Unfortunately, the classical dynamics is implementable only if the vector potential \vec{A} vanishes everywhere. Therefore, \hat{H}^{ren} is well defined only in this case. However, the infimum of (1.9), that is E^{ren} , is a well defined gauge-invariant number also for nonzero \vec{A} .

Note that both Hamiltonians (1.6) and (1.9) can be derived from local Lagrangians. Therefore, even if the models based on these Hamiltonians do not satisfy Haag-Kastler axioms in the strict sense (because of the absence of translation invariance), they belong to Local Quantum Field Theory: they lead to nets satisfying the Einstein causality, and they have bounded from below Hamiltonians. At the same time, all of them require an infinite renormalization, typical for computations in perturbative Quantum Field Theory.

The examples 1.5 and 1.9 are especially interesting in the context of more complicated interacting quantum field theory when, typically, κ , A^0 and \vec{A} are promoted to the role of a quantum fields. Then E^{ren} can be interpreted as the value of certain renormalized loop diagrams. In particular, E^{ren} of the second example is usually called the *vacuum poarization (in scalar QED)*.

Acknowledgements. I would like to thank Marcin Napiórkowski, Phan Thành Tham and Jan Philip Solovej for useful discussions. I gratefully acknowledge financial support of the National Science Center, Poland, under the grant UMO-2014/15/B/ST1/00126.

2 Finite dimensions, basis dependent formalism

Let us first describe the basic theory of bosonic quadratic Hamiltonians in finite dimensions, assuming that the one-particle space is \mathbb{C}^m . Seemingly, our formulas will depend on the choice of the canonical basis

in \mathbb{C}^m . In reality, after an appropriate interpretation, they are basis independent. This interpretation will be given in the next section, when we discuss an arbitrary dimension.

Operators on \mathbb{C}^m will be identified with matrices. If $h = [h_{ij}]$ is a matrix, then \bar{h} , h^* and $h^\#$ will denote its complex conjugate, hermitian conjugate and transpose.

2.1 Creation/annihilation operators

We consider the bosonic Fock space $\Gamma_s(\mathbb{C}^m)$. \hat{a}_i, \hat{a}_j^* are the standard annihilation and creation operators on $\Gamma_s(\mathbb{C}^m)$. \hat{a}_i^* is the Hermitian conjugate of \hat{a}_i ,

$$\begin{aligned} [\hat{a}_i, \hat{a}_j] &= [\hat{a}_i, \hat{a}_j] = 0, \\ [\hat{a}_i, \hat{a}_j^*] &= \delta_{ij}. \end{aligned}$$

(We decorate creation/annihilation operators with hats, because we want to distinguish them from their classical analogs).

We use the more or less standard notation for operators on Fock spaces. In particular, we use the standard notation $\Gamma(\cdot)$ and $d\Gamma(\cdot)$, which will be recalled in Subsection 3.2. If $w = [w_i] \in \mathbb{C}^m$, then the corresponding creation/annihilation operators are

$$\hat{a}^*(w) := \sum_i w_i \hat{a}_i^*, \quad \hat{a}(w) := \sum_i \bar{w}_i \hat{a}_i. \quad (2.1)$$

If $g = [g_{ij}]$ is a symmetric $m \times m$ matrix, then the corresponding double creation/annihilation operators are

$$\hat{a}^*(g) := \sum_{ij} g_{ij} \hat{a}_i^* \hat{a}_j^*, \quad \hat{a}(g) := \sum_{ij} \bar{g}_{ij} \hat{a}_j \hat{a}_i.$$

2.2 Classical phase space

To specify a linear combination of operators \hat{a}_i, \hat{a}_j^* we need to choose a vector $(w, w') \in \mathbb{C}^m \oplus \mathbb{C}^m$:

$$\hat{\phi}(w, w') := \sum_i \hat{a}_i^* w_i + \sum_i \hat{a}_i w'_i \quad (2.2)$$

(2.2) is self-adjoint iff $\bar{w} = w'$. Therefore, it is natural to introduce the doubled space $\mathbb{C}^m \oplus \mathbb{C}^m$ equipped with the *complex conjugation*

$$J \begin{bmatrix} w \\ w' \end{bmatrix} = \begin{bmatrix} \bar{w}' \\ \bar{w} \end{bmatrix}. \quad (2.3)$$

Vectors left invariant by J have the form

$$\begin{bmatrix} w \\ \bar{w} \end{bmatrix}, \quad w \in \mathbb{C}^m. \quad (2.4)$$

They form a $2m$ -dimensional real subspace of $\mathbb{C}^m \oplus \mathbb{C}^m$, which can be identified with \mathbb{R}^{2m} . (In what follows, when we speak of \mathbb{R}^{2m} we usually mean the space of vectors of the form (2.4)).

Operators on $\mathbb{C}^m \oplus \mathbb{C}^m$ that commute with J , or equivalently preserve \mathbb{R}^{2m} , have the form

$$R = \begin{bmatrix} p & q \\ \bar{q} & \bar{p} \end{bmatrix}, \quad (2.5)$$

and will be called *J-real*. Note that if we know the restriction of R to (2.4) then we can uniquely extend it to a (complex linear) operator on $\mathbb{C}^m \oplus \mathbb{C}^m$.

The operator

$$S = \begin{bmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{bmatrix}. \quad (2.6)$$

determines the commutation relations:

$$\left[\hat{\phi}(w_1, w'_1)^*, \hat{\phi}(w_2, w'_2) \right] = (w_1|w_2) - (w'_1|w'_2) = ((w_1, w'_1)|S(w_2, w'_2)). \quad (2.7)$$

Instead of quantum operators \hat{a}_i^* and \hat{a}_j , one can also consider classical (commuting) variables a_i, a_j^* , $i = 1, \dots, m$, such that a_i^* is the complex conjugate of a_i and the following Poisson bracket relations hold:

$$\begin{aligned} \{a_i, a_j\} &= \{a_i, a_j\} = 0, \\ \{a_i, a_j^*\} &= -i\delta_{ij}. \end{aligned} \quad (2.8)$$

Setting

$$\phi(w, w') := \sum_i a_i^* w_i + \sum_i a_i w'_i, \quad (2.9)$$

we can rewrite (2.8) as

$$\{\phi(w_1, w'_1)^*, \phi(w_2, w'_2)\} = -i(w_1|w_2) + i(w'_1|w'_2) = -i((w_1, w'_1)|S(w_2, w'_2)). \quad (2.10)$$

In particular, $\phi(w, \bar{w})$ are real, and (2.10) can be rewritten as

$$\{\phi(w_1, \bar{w}_1)^*, \phi(w_2, \bar{w}_2)\} = 2\text{Im}(w_1|w_2) = \text{Im}((w_1, \bar{w}_1)|S(w_2, \bar{w}_2)). \quad (2.11)$$

Thus S determines a symplectic structure on \mathbb{R}^{2m} (and sometimes S itself is called, incorrectly, a symplectic form).

2.3 Symplectic transformations

In this subsection we recall some basic facts concerning the symplectic and metaplectic group. We follow mostly [DG].

We say that an operator R on $\mathbb{C}^m \oplus \mathbb{C}^m$ is *symplectic* if it is J -real and preserves S :

$$R^* S R = S. \quad (2.12)$$

We denote by $Sp(\mathbb{R}^{2m})$ the group of all symplectic transformations.

Note that if R is symplectic, then so is R^* . In fact, iS is symplectic, and

$$R^* = iS R^{-1} (iS)^{-1}. \quad (2.13)$$

The operator

$$R = \begin{bmatrix} p & q \\ \bar{q} & \bar{p} \end{bmatrix}. \quad (2.14)$$

satisfies (2.12) iff

$$\begin{aligned} p^* p - q^\# \bar{q} &= \mathbb{1}, & p^* q - q^\# \bar{p} &= 0, \\ pp^* - qq^* &= \mathbb{1}, & pq^\# - qp^\# &= 0. \end{aligned}$$

Note that

$$pp^* \geq \mathbb{1}, \quad p^* p \geq \mathbb{1}.$$

Hence p^{-1} is well defined, and we can set

$$d_1 := q^\# (p^\#)^{-1}, \quad (2.15)$$

$$d_2 := q\bar{p}^{-1}. \quad (2.16)$$

We have $d_1^\# = d_1, d_2 = d_2^\#$.

2.4 Metaplectic transformations

Let U be a unitary operator on $\Gamma_s(\mathbb{C}^m)$. Let R be a symplectic transformation written as (2.14). We say that U implements R if

$$\begin{aligned} U\hat{a}_i^*U^* &= \hat{a}_j^*p_{ji} + \hat{a}_j\bar{q}_{ji}, \\ U\hat{a}_iU^* &= \hat{a}_j^*q_{ji} + \hat{a}_j\bar{p}_{ji}. \end{aligned}$$

U will be called a (Bogoliubov) implementer of R . Every symplectic transformation has an implementer, unique up to a phase factor. One can distinguish some canonical choices: the natural implementer U_R^{nat} , and a pair of metaplectic implementers $\pm U_R^{\text{met}}$:

$$U_R^{\text{nat}} := |\det pp^*|^{-\frac{1}{4}} e^{-\frac{1}{2}\hat{a}^*(d_2)} \Gamma((p^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_1)}, \quad (2.17)$$

$$\pm U_R^{\text{met}} := \pm (\det p^*)^{-\frac{1}{2}} e^{-\frac{1}{2}\hat{a}^*(d_2)} \Gamma((p^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_1)}, \quad (2.18)$$

see eg. Thm 11.33 and Def. 11.36 of [DG].

It is easy to see that the set of Bogoliubov implementers is a group. It is sometimes called the c -metaplectic group $Mp^c(\mathbb{R}^{2m})$.

It is a little less obvious, but also true, that the set of metaplectic Bogoliubov implementers is a subgroup of $Mp^c(\mathbb{R}^{2m})$. It is called the metaplectic group $Mp(\mathbb{R}^{2m})$.

We have a homomorphism $Mp^c(\mathbb{R}^{2m}) \ni U \mapsto R \in Sp(\mathbb{R}^{2m})$, where U implements R .

Various homomorphism related to the metaplectic group can be described by the following diagram:

$$\begin{array}{ccccccc} & & 1 & & 1 & & 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & \mathbb{Z}_2 & \rightarrow & U(1) & \rightarrow & U(1) \rightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & Mp(\mathbb{R}^{2m}) & \rightarrow & Mp^c(\mathbb{R}^{2m}) & \rightarrow & U(1) \rightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & Sp(\mathbb{R}^{2m}) & \rightarrow & Sp(\mathbb{R}^{2m}) & \rightarrow & 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array} \quad (2.19)$$

2.5 Positive symplectic transformations

Positive symplectic transformations are especially important. They satisfy

$$p = p^*, \quad p > 0, \quad q = q^\#. \quad (2.20)$$

For positive transformations, d_1 equals d_2 , and it will be simply denoted by d . We have

$$d := q(p^\#)^{-1}.$$

The natural implementer coincides in this case with one of the metaplectic implementers:

$$U_R^{\text{nat}} := (\det p)^{-\frac{1}{2}} e^{-\frac{1}{2}\hat{a}^*(d)} \Gamma(p^{-1}) e^{\frac{1}{2}\hat{a}(d)}.$$

Positive symplectic transformations have special properties. In particular, one can diagonalize them in an explicit way. We will need this later on.

Proposition 2.1 *Assume that R is positive symplectic and $\text{Ker}(p - \mathbb{1}) = \{0\}$. Then q is invertible, so that we can define $u := q|q|^{-1}$ with $|q| := \sqrt{q^*\bar{q}}$. Besides,*

$$M := \frac{1}{\sqrt{2}} \begin{bmatrix} \mathbb{1} & -u \\ u^* & \mathbb{1} \end{bmatrix} \quad (2.21)$$

is unitary and diagonalizes R :

$$R = M \begin{bmatrix} p + \sqrt{p^2 - \mathbb{1}} & 0 \\ 0 & \bar{p} - \sqrt{\bar{p}^2 - \mathbb{1}} \end{bmatrix} M^*. \quad (2.22)$$

Proof. We have the polar decomposition $q = u|q|$. u is a unitary operator and we have $|\bar{q}| = u|q|u^*$. Now (2.22) follows using $u\bar{p}u^* = p$, $|q| = \sqrt{p^2 - \mathbb{1}}$, $\sqrt{\mathbb{1} + |q|^2} = \bar{p}$. \square

2.6 Classical quadratic Hamiltonians

It is easy to analyze generators of 1-parameter symplectic groups. In fact, $e^{itB} \in Sp(\mathbb{R}^{2m})$ for any $t \in \mathbb{R}$ iff BS is J -real and self-adjoint. All such operators can be written as

$$B = \begin{bmatrix} h & -g \\ \bar{g} & -\bar{h} \end{bmatrix}. \quad (2.23)$$

where h, g are $m \times m$ matrices satisfying $h = h^*$, $g = g^\#$. Note that iB is J -real, and

$$SB = B^*S. \quad (2.24)$$

With every such an operator B we associate another operator A_B by

$$A_B := BS = \begin{bmatrix} h & g \\ \bar{g} & \bar{h} \end{bmatrix}, \quad (2.25)$$

As we noted above, A_B is self-adjoint and J -real. The corresponding *classical quadratic Hamiltonian* is the expression

$$H_B = \sum h_{ij} a_i^* a_j + \frac{1}{2} \sum g_{ij} a_i^* a_j^* + \frac{1}{2} \sum \bar{g}_{ij} a_i a_j, \quad (2.26)$$

which can be viewed as a quadratic function on the classical phase space. Moreover,

$$\{H_B, \phi(w, w')\} = -i\phi(w_1, w'_1), \quad \begin{bmatrix} w_1 \\ w'_1 \end{bmatrix} = B \begin{bmatrix} w \\ w' \end{bmatrix}. \quad (2.27)$$

Clearly, for any symplectic R ,

$$A_{RBR^{-1}} = RA_B R^*. \quad (2.28)$$

In what follows we will often abuse the terminology: A_B will also be called a *classical Hamiltonian* just as H_B . B will be called a *symplectic generator*. Besides, we will often drop the subscript B from H_B and A_B .

2.7 Quantum quadratic Hamiltonians

Let B be a symplectic generator of the form (2.23).

By a *quantization of H_B* (2.26) we will mean an operator on $\Gamma_s(\mathbb{C}^m)$ of the form

$$\hat{H}_B^c := \sum h_{ij} \hat{a}_i^* \hat{a}_j + \frac{1}{2} \sum g_{ij} \hat{a}_i^* \hat{a}_j^* + \frac{1}{2} \sum \bar{g}_{ij} \hat{a}_i \hat{a}_j + c, \quad (2.29)$$

where c is an arbitrary real constant. By an abuse of terminology, we will usually say that (2.29) is a *quantization of B* (2.25). We will often drop the subscript B from \hat{H}_B^c , and c will be replaced by other superscripts corresponding to some special choices.

Two quantizations of B are especially useful: the *Weyl* (or *symmetric*) *quantization* \hat{H}_B^w and the *normally ordered* (or *Wick*) *quantization* \hat{H}_B^n :

$$\begin{aligned}\hat{H}_B^w &:= \frac{1}{2} \sum h_{ij} \hat{a}_i^* \hat{a}_j + \frac{1}{2} \sum h_{ij} \hat{a}_j \hat{a}_i^* + \frac{1}{2} \sum g_{ij} \hat{a}_i^* \hat{a}_j^* + \frac{1}{2} \sum \bar{g}_{ij} \hat{a}_i \hat{a}_j, \\ \hat{H}_B^n &:= \sum h_{ij} \hat{a}_i^* \hat{a}_j + \frac{1}{2} \sum g_{ij} \hat{a}_i^* \hat{a}_j^* + \frac{1}{2} \sum \bar{g}_{ij} \hat{a}_i \hat{a}_j.\end{aligned}$$

Here is the relation between these two quantizations:

$$\hat{H}_B^w = \hat{H}_B^n + \frac{1}{2} \text{Tr} h. \quad (2.30)$$

Note a special relationship of the Weyl quantization to the metaplectic group (defined in Subsect. 2.3): for any B , $e^{it\hat{H}_B^w}$ belongs to $Mp(\mathbb{R}^{2m})$, see eg. Thm 11.34 of [DG]. Besides, if R is symplectic and U_R is its implementer, then

$$U_R \hat{H}_B^w U_R^* = \hat{H}_{RBR^{-1}}^w. \quad (2.31)$$

2.8 Diagonalization of quadratic Hamiltonians

In this subsection we show that if $A_B > 0$, then A_B can be *diagonalized*. By this we mean that we can find a symplectic transformation R that kills off-diagonal terms of A_B :

$$A_B = R \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & \bar{h}_{\text{dg}} \end{bmatrix} R^* \quad (2.32)$$

for some h_{dg} . Of course, h_{dg} has to be positive.

Clearly, this is equivalent to *diagonalizing* B , that is to killing its off-diagonal terms: of B , that is,

$$B = R \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & -\bar{h}_{\text{dg}} \end{bmatrix} R^{-1}. \quad (2.33)$$

On the quantum level, this is equivalent to finding a unitary operator U that removes double annihilators and double creators. Then the free constant equals the infimum of the quantum Hamiltonian:

$$\begin{aligned}U^* \hat{H}^w U &= d\Gamma(h_{\text{dg}}) + E^w, \\ U^* \hat{H}^n U &= d\Gamma(h_{\text{dg}}) + E^n.\end{aligned}$$

As a preparation for a construction of a diagonalizing operator, let us prove the following proposition.

In this proposition we will use the function $\text{sgnt} := \begin{cases} 1 & t > 0, \\ 0 & t = 0, \\ -1 & t < 0. \end{cases}$

Proposition 2.2 *Suppose that $A_B > 0$.*

- (1) *The operator B has only real nonzero eigenvalues. Therefore, sgn can be interpreted as a holomorphic function on a neighborhood of $\text{sp}B$, and we can define $\text{sgn}(B)$ by the standard holomorphic functional calculus.*
- (2) *A symplectic transformation R diagonalizes B iff*

$$\text{sgn}(B) = RSR^{-1}. \quad (2.34)$$

Proof. It is useful to endow the space $\mathbb{C}^m \oplus \mathbb{C}^m$ with the scalar product given by the positive operator SAS . More precisely, if $v = (v_1, v_2), w = (w_1, w_2) \in \mathbb{C}^m \oplus \mathbb{C}^m$, we set

$$(v|w)_{\text{en}} = (v|SASw) = (v_1|hw_1) - (v_1|gw_2) - (v_2|\bar{g}w_1) + (v_2|\bar{h}w_2). \quad (2.35)$$

(2.35) is sometimes called the *energy scalar product*.

Note that we also have the original scalar product

$$(v|w) = (v_1|w_1) + (v_2|w_2),$$

which is used for basic notation such as the Hermitian adjoints.

First note that B is self-adjoint in the energy scalar product and has a zero nullspace. Indeed

$$\begin{aligned} (v|Bw)_{\text{en}} &= (v|SASASw) \\ &= (ASv|SASw) = (Bv|w)_{\text{en}}, \\ (Bv|Bv)_{\text{en}} &= (v|SASASASv) > 0, \quad v \neq 0. \end{aligned}$$

This shows (1).

Now let R be symplectic. Set

$$B_{\text{dg}} := R^{-1}BR, \quad A_{\text{dg}} := B_{\text{dg}}S = R^{-1}AR^{*-1}.$$

Then, by functional calculus,

$$\text{sgn}(B) = R\text{sgn}(B_{\text{dg}})R^{-1}. \quad (2.36)$$

R diagonalizes A iff

$$A_{\text{dg}} = \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & \bar{h}_{\text{dg}} \end{bmatrix}, \quad B_{\text{dg}} = \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & -\bar{h}_{\text{dg}} \end{bmatrix}. \quad (2.37)$$

A is strictly positive, hence so are A_{dg} and h_{dg} . Therefore,

$$\text{sgn}(B_{\text{dg}}) = S. \quad (2.38)$$

Together with (2.36), this implies (2.34).

Conversely, suppose that (2.34) holds. Together with (2.36), this implies (2.38). Hence B_{dg} is diagonal. \square

It is possible to find a distinguished positive symplectic transformation R diagonalizing B .

Theorem 2.3 *Suppose that $A_B > 0$.*

- (1) $i\text{sgn}(B)$ is symplectic.
- (2) $R_0 := \text{sgn}(B)S$ is symplectic and has positive eigenvalues.
- (3) Using holomorphic calculus and the principal square root (which for positive arguments has positive values), define

$$R := R_0^{\frac{1}{2}}. \quad (2.39)$$

Then R is positive, symplectic and diagonalizes B .

- (4) Here is an alternative formula for R_0 , where the square root can be interpreted in terms of functional calculus for self-adjoint operators:

$$R_0 = SA_B^{-\frac{1}{2}} \left(A_B^{\frac{1}{2}} SA_B SA_B^{\frac{1}{2}} \right)^{\frac{1}{2}} A_B^{-\frac{1}{2}} S. \quad (2.40)$$

Proof. B satisfies (2.24). Hence for any function f holomorphic on the spectrum of B

$$Sf(B)S^{-1} = f(B^*). \quad (2.41)$$

In particular

$$S\text{sgn}(B)S^{-1} = \text{sgn}(B^*). \quad (2.42)$$

But sgn is real, hence $\text{sgn}(B^*) = \text{sgn}(B)^*$. Besides, away from 0 we have $\text{sgn}(t) = \text{sgn}(t)^{-1}$. Hence, $\text{sgn}(B) = \text{sgn}(B)^{-1}$. Therefore, (2.42) can be rewritten as

$$S\text{sgn}(B)S^{-1} = \text{sgn}(B)^{* -1}. \quad (2.43)$$

Hence,

$$(i\text{sgn}(B))^* S i\text{sgn}(B) = S. \quad (2.44)$$

This means that $i\text{sgn}(B)$ preserves S . Besides,

$$i\text{sgn}B = (- (iB)^2)^{\frac{1}{2}} (iB)^{-1} \quad (2.45)$$

is also J -real. Thus we have shown that $i\text{sgn}(B)$ is symplectic.

$-iS$ is also symplectic. Therefore, so is $R_0 = (i\text{sgn}(B))(-iS)$.

Now,

$$R_0 = (B^2)^{\frac{1}{2}} B^{-1} S \quad (2.46)$$

$$= (ASAS)^{\frac{1}{2}} SA^{-1} S \quad (2.47)$$

$$= SA^{-\frac{1}{2}} (A^{\frac{1}{2}} SAS A^{\frac{1}{2}})^{\frac{1}{2}} A^{-\frac{1}{2}} S. \quad (2.48)$$

Therefore, (2.40) is true and R_0 is a positive self-adjoint operator for the original scalar product. Hence it has positive eigenvalues.

$R_0 = R_0^*$ and R_0 is symplectic. Hence,

$$SR_0S^{-1} = R_0^{-1}.$$

Hence for any Borel function f ,

$$Sf(R_0)S^{-1} = f(R_0^{-1}).$$

Choosing f to be the (positive) square root we obtain

$$SR_0^{\frac{1}{2}}S^{-1} = R_0^{-\frac{1}{2}}.$$

Thus $R := R_0^{\frac{1}{2}}$ is symplectic, positive and self-adjoint for the original scalar product.

Now

$$\text{sgn}(B) = R^2 S = RSR^{-1}.$$

Hence (2.34) is true. \square

2.9 Positive Weyl Bogoliubov Hamiltonians

Theorem 2.4 (1) *If $A_B \geq 0$, then the Weyl quantization of B is positive. Hence all quantizations of B are bounded from below.*

(2) *If B possesses a quantization that is bounded from below, then $A_B \geq 0$.*

Proof. (1) Let $A \geq 0$. Then there exists a symplectic transformation R and a decomposition $\mathbb{C}^m = \mathbb{C}^{m_1} \oplus \mathbb{C}^{m-m_1}$ such that RAR^* decomposes into the direct sum of the following two terms:

$$\begin{aligned} & \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & \bar{h}_{\text{dg}} \end{bmatrix} && \text{on } \mathbb{C}^{m_1} \oplus \mathbb{C}^{m_1}, \\ & \frac{1}{2} \begin{bmatrix} \mathbb{1} & \mathbb{1} \\ \mathbb{1} & \mathbb{1} \end{bmatrix} && \text{on } \mathbb{C}^{m-m_1} \oplus \mathbb{C}^{m-m_1}, \end{aligned}$$

where $h_{\text{dg}} \geq 0$ and can be assumed to be diagonal. This is a well-known fact, proven eg. in [Ho1, DG]. It is a very special case of a more general and more complicated classification of quadratic forms on a symplectic space called Williamson's Theorem, proven eg. in [Wi, Ho]. If we strengthen the assumption and demand that $h > 0$, it follows also from the diagonalizability of A (Theorem 2.3). Thus, after an application of the transformation R , and a diagonalization of h_{dg} , the classical Hamiltonian becomes

$$H_{RBR^{-1}} = \sum_{i=1}^{m_1} h_{\text{dg},ii} a_i^* a_i + \sum_{m_1+1}^m \frac{1}{2} (a_i^* + a_i)^2. \quad (2.49)$$

After application of an implementer of R , the quantum Weyl Hamiltonian becomes

$$U_R \hat{H}_B^w U_R^* = \hat{H}_{RBR^{-1}}^w = \sum_{i=1}^{m_1} \frac{1}{2} h_{\text{dg},ii} (\hat{a}_i^* \hat{a}_i + \hat{a}_i \hat{a}_i^*) + \sum_{m_1+1}^m \frac{1}{2} (\hat{a}_i^* + \hat{a}_i)^2. \quad (2.50)$$

Thus \hat{H}_B^w is positive.

(2) Consider the family of coherent vectors

$$\Omega_w := e^{\hat{a}^*(w) - \hat{a}(w)} \Omega, \quad w \in \mathbb{C}^m. \quad (2.51)$$

Note that

$$e^{-\hat{a}^*(w) + \hat{a}(w)} \hat{a}_i^* e^{\hat{a}^*(w) - \hat{a}(w)} = \hat{a}_i^* + \bar{w}_i, \quad e^{-\hat{a}^*(w) + \hat{a}(w)} \hat{a}_i e^{\hat{a}^*(w) - \hat{a}(w)} = \hat{a}_i + w_i. \quad (2.52)$$

Obviously, if one of quantizations of B is bounded from below, then so are all of them. Let \hat{H}_B^n be bounded from below by $-c$. Then, using (2.52), we obtain

$$-c \leq (\Omega_w | \hat{H}_B^n \Omega_w) \quad (2.53)$$

$$= \left(\Omega | e^{-\hat{a}^*(w) + \hat{a}(w)} \hat{H}_B^n e^{\hat{a}^*(w) - \hat{a}(w)} \Omega \right) \quad (2.54)$$

$$= \sum h_{ij} \bar{w}_i w_j + \frac{1}{2} \sum g_{ij} \bar{w}_i \bar{w}_j + \frac{1}{2} \sum \bar{g}_{ij} w_i w_j. \quad (2.55)$$

Thus the classical Hamiltonian is quadratic polynomial and is bounded from below. But if a quadratic polynomial is bounded from below, then it is nonnegative. \square

Note that by the above theorem, every B satisfying $A_B \geq 0$, beside \hat{H}_B^w and \hat{H}_B^n , possesses another natural quantization: the *zero infimum quantization* \hat{H}_B^z fixed by the condition

$$\inf \hat{H}_B^z = 0. \quad (2.56)$$

The infimum of the Weyl Bogoliubov Hamiltonians can be computed from several formulas described in the following theorem borrowed from [BD, DG]:

Theorem 2.5 *Assume that $A_B \geq 0$. Then*

$$E_B^w := \inf \hat{H}_B^w = \frac{1}{4} \text{Tr} \sqrt{B^2} \quad (2.57)$$

$$= \frac{1}{4} \text{Tr} \left[\begin{array}{cc} h^2 - g\bar{g} & -hg + g\bar{h} \\ \bar{g}h - \bar{h}\bar{g} & \bar{h}^2 - \bar{g}g \end{array} \right]^{\frac{1}{2}} \quad (2.58)$$

$$= \frac{1}{4} \text{Tr} \sqrt{A^{\frac{1}{2}} S A S A^{\frac{1}{2}}} \quad (2.59)$$

$$= \frac{1}{4} \text{Tr} \int \frac{B^2}{B^2 + \tau^2} \frac{d\tau}{2\pi}. \quad (2.60)$$

Proof. Let R be as in the proof of Theorem 2.4. Clearly,

$$\inf(\hat{a}_i^* \hat{a}_i + \hat{a}_i \hat{a}_i^*) = 1, \quad (2.61)$$

$$\inf(\hat{a}_i^* + \hat{a}_i)^2 = 0. \quad (2.62)$$

Hence, by (2.50),

$$\inf \hat{H}_B^w = \inf U_R \hat{H}_B^w U_R^* = \inf \hat{H}_{RBR^{-1}}^w = \frac{1}{2} \sum h_{\text{dg}, ii} = \frac{1}{2} \text{Tr} h_{\text{dg}} \quad (2.63)$$

$$= \frac{1}{4} \text{Tr} \sqrt{B_{\text{dg}}^2} = \frac{1}{4} \text{Tr} R \sqrt{B^2} R^{-1} = \frac{1}{4} \text{Tr} \sqrt{B^2}. \quad (2.64)$$

This gives (2.57), which implies (2.58) and (2.59).

(2.60) follows by an application of the identity (A.5). \square

2.10 Infimum of normally ordered Hamiltonians

Here are a few formulas for the infimum of the normally ordered Hamiltonian:

Theorem 2.6 *Assume that $A_B \geq 0$. Then*

$$E_B^n := \inf \hat{H}_B^n = E_B^w - \frac{1}{2} \text{Tr} h \quad (2.65)$$

$$= \frac{1}{4} \text{Tr} (\sqrt{B^2} - \sqrt{B_0^2}) \quad (2.66)$$

$$= \frac{1}{4} \text{Tr} \left(\left[\begin{array}{cc} h^2 - g\bar{g} & -hg + g\bar{h} \\ \bar{g}h - \bar{h}\bar{g} & \bar{h}^2 - \bar{g}g \end{array} \right]^{\frac{1}{2}} - \left[\begin{array}{cc} h & 0 \\ 0 & \bar{h} \end{array} \right] \right) \quad (2.67)$$

$$= \frac{1}{8} \int_0^1 d\sigma \text{Tr} \frac{B_\sigma}{\sqrt{B_\sigma^2}} G. \quad (2.68)$$

$$= \frac{1}{8} \int_0^1 d\sigma \text{Tr} A_\sigma^{\frac{1}{2}} (A_\sigma^{\frac{1}{2}} S A_\sigma S A_\sigma^{\frac{1}{2}})^{-\frac{1}{2}} A_\sigma^{\frac{1}{2}} G S \quad (2.69)$$

$$= \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1 - \sigma) \text{Tr} \frac{1}{(A_\sigma + i\tau S)} S G \frac{1}{(A_\sigma + i\tau S)} S G. \quad (2.70)$$

where

$$G := B - B_0 = \begin{bmatrix} 0 & -g \\ \bar{g} & 0 \end{bmatrix}, \quad (2.71)$$

$$A_\sigma = A_0 + \sigma GS = \begin{bmatrix} h & \sigma g \\ \sigma \bar{g} & \bar{h} \end{bmatrix}, \quad (2.72)$$

$$B_\sigma := B_0 + \sigma G = \begin{bmatrix} h & -\sigma g \\ \sigma \bar{g} & -\bar{h} \end{bmatrix}, \quad \sigma \in \mathbb{R}. \quad (2.73)$$

Proof. (2.65), (2.66) and (2.67) follow immediately from Theorem 2.5 and (2.30). Starting from (2.66), let us prove (2.68), which immediately implies (2.69):

$$\frac{1}{4} \text{Tr}(\sqrt{B^2} - \sqrt{B_0^2}) \quad (2.74)$$

$$= \frac{1}{4} \int \text{Tr} \left(\frac{B^2}{B^2 + \tau^2} - \frac{B_0^2}{B_0^2 + \tau^2} \right) \frac{d\tau}{2\pi} \quad (2.75)$$

$$= -\frac{1}{4} \int \text{Tr} \left(\frac{1}{B^2 + \tau^2} - \frac{1}{B_0^2 + \tau^2} \right) \frac{\tau^2 d\tau}{2\pi} \quad (2.76)$$

$$= -\frac{1}{4} \int_0^1 d\sigma \int \frac{d}{d\sigma} \text{Tr} \frac{1}{B_\sigma^2 + \tau^2} \frac{\tau^2 d\tau}{2\pi} \quad (2.77)$$

$$= \frac{1}{4} \int_0^1 d\sigma \int \text{Tr} \frac{1}{B_\sigma^2 + \tau^2} (B_\sigma G + G B_\sigma) \frac{1}{B_\sigma^2 + \tau^2} \frac{\tau^2 d\tau}{2\pi} \quad (2.78)$$

$$= \frac{1}{2} \int_0^1 d\sigma \int \text{Tr} \frac{B_\sigma}{(B_\sigma^2 + \tau^2)^2} G \frac{\tau^2 d\tau}{2\pi} \quad (2.79)$$

$$= \frac{1}{8} \int_0^1 d\sigma \text{Tr} \frac{B_\sigma}{\sqrt{B_\sigma^2}} G, \quad (2.80)$$

where at the end we used the identity (A.6).

Now, starting from (2.68), we prove (2.70):

$$\frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} \text{Tr} \frac{B_\sigma}{B_\sigma^2 + \tau^2} G \quad (2.81)$$

$$= \frac{1}{16} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} \text{Tr} \left(\frac{1}{(B_\sigma + i\tau)} + \frac{1}{(B_\sigma - i\tau)} \right) G \quad (2.82)$$

$$= \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} \text{Tr} \frac{1}{(B_\sigma + i\tau)} G \quad (2.83)$$

$$= \frac{1}{8} \int_0^1 d\sigma \int_0^\sigma d\sigma_1 \int \frac{d\tau}{2\pi} \frac{d}{d\sigma_1} \text{Tr} \frac{1}{(B_{\sigma_1} + i\tau)} G \quad (2.84)$$

$$= -\frac{1}{8} \int_0^1 (1 - \sigma) d\sigma \int \frac{d\tau}{2\pi} \text{Tr} \frac{1}{(B_\sigma + i\tau)} G \frac{1}{(B_\sigma + i\tau)} G \quad (2.85)$$

$$= -\frac{1}{8} \int_0^1 (1 - \sigma) d\sigma \int \frac{d\tau}{2\pi} \text{Tr} \frac{1}{(A_\sigma + i\tau S)} S G \frac{1}{(A_\sigma + i\tau S)} S G. \quad (2.86)$$

In (2.84) \Rightarrow (2.85) we used

$$\frac{d}{d\sigma_1} \frac{1}{(B_{\sigma_1} + i\tau)} = -\frac{1}{(B_{\sigma_1} + i\tau)} G \frac{1}{(B_{\sigma_1} + i\tau)}. \quad (2.87)$$

□

2.11 Loop expansion

Suppose now that

$$B_0 = \begin{bmatrix} h_0 & 0 \\ 0 & -\bar{h}_0 \end{bmatrix}, \quad (2.88)$$

is a “free” symplectic generator. We assume that $h_0 > 0$. Note that we allow h_0 to be different from h .

We set

$$A_0 := B_0 S = \begin{bmatrix} h_0 & 0 \\ 0 & \bar{h}_0 \end{bmatrix}, \quad (2.89)$$

$$V := B^2 - B_0^2 = \begin{bmatrix} h^2 - h_0^2 - g\bar{g} & -hg + g\bar{h} \\ \bar{g}h - \bar{h}\bar{g} & \bar{h}^2 - \bar{h}_0^2 - \bar{g}g \end{bmatrix}. \quad (2.90)$$

(2.60) can be rewritten as

$$\begin{aligned} E^w &= \frac{1}{4} \text{Tr} \int \frac{B_0^2}{B_0^2 + \tau^2} \frac{d\tau}{2\pi} + \frac{1}{4} \text{Tr} \int \frac{1}{B^2 + \tau^2} V \frac{1}{B_0^2 + \tau^2} \tau^2 \frac{d\tau}{2\pi} \\ &= \sum_{j=0}^k L_j + \frac{1}{4} \text{Tr} \int \frac{(-1)^k}{B_0^2 + \tau^2} V \frac{1}{B^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^k \tau^2 \frac{d\tau}{2\pi} \\ &= \sum_{j=0}^{\infty} L_j, \end{aligned}$$

where

$$L_0 = \frac{1}{4} \text{Tr} \int \frac{B_0^2}{B_0^2 + \tau^2} \frac{d\tau}{2\pi} = \frac{1}{4} \text{Tr} |B_0| = \frac{1}{2} \text{Tr} h_0, \quad (2.91)$$

$$L_j = \frac{1}{4} \text{Tr} \int \frac{(-1)^{j+1}}{B_0^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^j \tau^2 \frac{d\tau}{2\pi} \quad (2.92)$$

$$= \frac{1}{4} \text{Tr} \int \frac{(-1)^{j+1}}{2j} \left(V \frac{1}{B_0^2 + \tau^2} \right)^j \frac{d\tau}{2\pi}, \quad j = 1, 2, \dots \quad (2.93)$$

The last identity for L_j follows by a cyclic relocation of operators under the trace and by an application of integration by parts.

We can further simplify the formula for L_1 :

$$L_1 = \text{Tr} \int \frac{1}{8} V \frac{1}{B_0^2 + \tau^2} \frac{d\tau}{2\pi} = \frac{1}{8} \text{Tr} V \frac{1}{|B_0|} = \frac{1}{4} \text{Tr} (h^2 - h_0^2 - g\bar{g}) h_0^{-1}. \quad (2.94)$$

The constant L_j arises in the diagrammatic expansion as the evaluation of the loop with j vertices. To see this, introduce the “time variable” t and the “Feynman propagator”

$$G(t) := \frac{e^{-|B_0|t}}{2|B_0|}.$$

Clearly, τ can be interpreted as the “energy variable” and

$$\frac{1}{B_0^2 + \tau^2} = \int G(t) e^{it\tau} dt.$$

Therefore,

$$L_j = \frac{1}{4} \int dt_{j-1} \cdots \int dt_1 \text{Tr} V G(t_j - t_1) V G(t_1 - t_2) \cdots V G(t_{j-1} - t_j) \quad (2.95)$$

$$= \lim_{T \rightarrow \infty} \frac{1}{2T} \frac{1}{4} \int_{-T}^T dt_j \int_{-T}^T dt_{j-1} \cdots \int_{-T}^T dt_1 \text{Tr} V G(t_j - t_1) V G(t_1 - t_2) \cdots V G(t_{j-1} - t_j). \quad (2.96)$$

2.12 Renormalization I

Note that in general V (2.90) contains terms of the 1st and 2nd order. Explicitly, let λ be a ‘‘coupling constant’’. Let $h = h_0 + \lambda h_1$ and replace g with λg (to keep track of the order of perturbation). Then $V = \lambda V_1 + \lambda^2 V_2$, where

$$V_1 := \begin{bmatrix} h_0 h_1 + h_1 h_0 & -h_0 g + g \bar{h}_0 \\ \bar{g} h_0 - \bar{h}_0 \bar{g} & \bar{h}_0 \bar{h}_1 + \bar{h}_0 \bar{h}_1 \end{bmatrix}, \quad (2.97)$$

$$V_2 := \begin{bmatrix} h_1^2 - g g^* & -h_1 g + g \bar{h}_1 \\ \bar{g} h_1 - \bar{h}_1 \bar{g} & \bar{h}_1^2 - g^* g \end{bmatrix}. \quad (2.98)$$

We can expand E^w wrt the coupling constant λ :

$$E^w = \sum_{n=0}^{\infty} \lambda^n E_n. \quad (2.99)$$

We have,

$$L_0 = E_0 = \frac{1}{2} \text{Tr} h_0. \quad (2.100)$$

However, in general, L_n of higher orders differ from $\lambda^n E_n$.

There are situations when it is natural to introduce the *renormalized vacuum energy*

$$E^{\text{ren}} := E^w - E_0 - \lambda E_1 - \lambda^2 E_2 = \sum_{n=3}^{\infty} \lambda^n E_n. \quad (2.101)$$

and the *renormalized Hamiltonian*

$$\hat{H}^{\text{ren}} := \hat{H}^w - E_0 - \lambda E_1 - \lambda^2 E_2, \quad (2.102)$$

so that $E^{\text{ren}} = \inf \hat{H}^{\text{ren}}$. The numbers E_0, E_1 and E_2 can be called *counterterms*.

The above constuctions are natural e.g. in the theory of charged scalar fields in external electromagnetic potentials. In this case, E_0, E_1, E_2 are infinite. \hat{H}^{ren} is usually also ill defined. However E^{ren} is typically finite. Thus we have a somewhat paradoxical situation: the Hamiltonian does not exists, however the ‘‘infimum of the Hamiltonian’’ is well defined.

2.13 Renormalization II

Suppose now that

$$h_1^2 = g \bar{g}, \quad h_1 g = g \bar{h}_1. \quad (2.103)$$

(2.103) implies $V_2 = 0$. Therefore, the loop expansion coincides with the expansion into powers of λ . Putting $\lambda = 1$, we thus have

$$E_n = L_n, \quad n = 0, 1, \dots \quad (2.104)$$

We can compute the loop with one vertex:

$$L_1 = \frac{1}{4} \text{Tr}(h_0 h_1 + h_1 h_0) h_0^{-1} = \frac{1}{2} \text{Tr} h_1. \quad (2.105)$$

Thus

$$L_0 + L_1 = \frac{1}{2} \text{Tr}(h_0 + h_1) = \frac{1}{2} \text{Tr} h. \quad (2.106)$$

Therefore, the loop expansion for the infimum of the normally ordered Hamiltonian amounts to omitting L_0 and L_1 :

$$\inf E^n = E^w - \frac{1}{2} \text{Tr} h = \sum_{n=2}^{\infty} L_n. \quad (2.107)$$

Note that L_1 , and especially L_0 , are often infinite. Sometimes, L_2 is infinite as well. Then we can renormalize the vacuum energy even further, introducing

$$E^{\text{ren}} := E^w - L_0 - L_1 - L_2 = \sum_{n=3}^{\infty} L_n \quad (2.108)$$

$$= -\frac{1}{4} \int \text{Tr} \frac{1}{B_0^2 + \tau^2} V \frac{1}{B^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^2 \tau^2 \frac{d\tau}{2\pi}. \quad (2.109)$$

We can also introduce the renormalized Hamiltonian

$$\hat{H}^{\text{ren}} := \hat{H}^w - L_0 - L_1 - L_2, \quad (2.110)$$

so that

$$E^{\text{ren}} = \inf \hat{H}^{\text{ren}}. \quad (2.111)$$

The situation described in this subsection is typical for a charged particle in an external electrostatic potential (without a vector potential), as well as for a neutral scalar particle with a masslike perturbation [De]. One can then often introduce the renormalized Hamiltonian \hat{H}^{ren} , which is a well-defined self-adjoint operator so that (2.111) holds.

3 Arbitrary dimensions, basis independent formalism

In this section we consider Bogoliubov Hamiltonians in any dimension. Unlike in the previous section, we will use a basis independent notation.

We will use the standard notation for the Hilbert-Schmidt and trace class norm:

$$\|g\|_2 := \sqrt{\text{Tr} g^* g}, \quad \|g\|_1 := \text{Tr} \sqrt{g^* g}. \quad (3.1)$$

3.1 Doubled space in abstract setting

Let \mathcal{W} be a Hilbert space. \mathcal{W} will serve as the *1-particle space*.

Let $\overline{\mathcal{W}}$ be another Hilbert space with a fixed antiunitary map $\chi : \mathcal{W} \rightarrow \overline{\mathcal{W}}$. $\overline{\mathcal{W}}$ will be called the *complex conjugate of \mathcal{W}* ,

We will often use the doubled space $\mathcal{W} \oplus \overline{\mathcal{W}}$ equipped with the conjugation

$$J = \begin{bmatrix} 0 & \chi^{-1} \\ \chi & 0 \end{bmatrix}. \quad (3.2)$$

A J -real operator is an operator on $\mathcal{W} \oplus \overline{\mathcal{W}}$ commuting with J . Bounded J -real operators have the form

$$R = \begin{bmatrix} p & q \\ \chi q \chi & \chi p \chi^{-1} \end{bmatrix}, \quad (3.3)$$

for some $p \in B(\mathcal{W})$, $q \in B(\overline{\mathcal{W}}, \mathcal{W})$.

J -real operators leave invariant the real subspace of vectors

$$\begin{bmatrix} w \\ \chi w \end{bmatrix}, \quad w \in \mathcal{W},$$

which we will denote \mathcal{Y} . Note also that every J -real operator in $B(\mathcal{W} \oplus \overline{\mathcal{W}})$ restricts to an operator $B(\mathcal{Y})$, and conversely, each operator in $B(\mathcal{Y})$ extends uniquely to an operator in $B(\mathcal{W} \oplus \overline{\mathcal{W}})$.

In what follows, we will usually write \bar{w} for χw . We will write \bar{p} , \bar{q} for $\chi p \chi^{-1}$ and $\chi q \chi$. We will write $p^\#$, $q^\#$ for $\chi p^* \chi^{-1}$ and $\chi^{-1} q^* \chi^{-1}$. In Appendix A.1 we explain why it is natural to use this simplified notation.

To reduce the formalism of this section to that of Section 2 it suffices to set $\mathcal{W} = \mathbb{C}^m$ and replace χ with the complex conjugation.

3.2 Fock spaces

If \mathcal{D} is a vector space of any dimension (with or without a Hilbert space structure), then we can introduce its *algebraic n th symmetric power*, denoted by $\overset{\text{al}}{\otimes}_s^n \mathcal{D}$ and the *algebraic bosonic Fock space*

$$\overset{\text{al}}{\Gamma}_s(\mathcal{W}) := \overset{\text{al}}{\bigoplus}_{n=0}^{\infty} \overset{\text{al}}{\otimes}_s^n \mathcal{D},$$

which is the space of finite symmetric tensor products of vectors of \mathcal{D} [DG]. If \mathcal{W} is a Hilbert space, then we prefer to use the Hilbert space versions of the above constructions. Thus $\otimes_s^n \mathcal{W}$ will denote the *n th symmetric tensor power of \mathcal{W} in the sense of Hilbert spaces* and, as usual, the *bosonic Fock space over the one-particle space \mathcal{W}* , is defined as

$$\Gamma_s(\mathcal{W}) := \bigoplus_{n=0}^{\infty} \otimes_s^n \mathcal{W}.$$

$\Omega := (1, 0, \dots)$ denotes the vacuum vector and

$$\begin{aligned} \Gamma_s^{\text{fin}}(\mathcal{W}) &:= \overset{\text{al}}{\bigoplus}_{n=0}^{\infty} \otimes_s^n \mathcal{W} \\ &= \{(\Psi^0, \dots, \Psi^n, \dots) \in \Gamma_s(\mathcal{W}) \mid \Psi^n = 0 \text{ for all but a finite number of } n\}, \end{aligned}$$

is the *finite particle bosonic Fock space*.

Note that if \mathcal{D} is dense in \mathcal{W} , then $\overset{\text{al}}{\Gamma}_s(\mathcal{D})$ is dense in $\Gamma_s(\mathcal{W})$.

If h is an operator on \mathcal{W} , $d\Gamma(h)$ will denote

$$d\Gamma(h) \upharpoonright_{\otimes_s^n \mathcal{W}} := \sum_{j=1}^n \underbrace{1 \otimes \dots \otimes 1}_{j-1} \otimes h \otimes \underbrace{1 \otimes \dots \otimes 1}_{n-j} \upharpoonright_{\otimes_s^n \mathcal{W}}.$$

If q is an operator on \mathcal{W} of norm less than 1, we define $\Gamma(q) : \Gamma_s(\mathcal{W}) \rightarrow \Gamma_s(\mathcal{W})$ by

$$\Gamma(q) \upharpoonright_{\otimes_s^n \mathcal{H}} := q \otimes \dots \otimes q \upharpoonright_{\otimes_s^n \mathcal{H}}.$$

3.3 Quadratic forms on Fock spaces

For any operator h on \mathcal{W} such that $h \geq c$, its *form domain* is defined as

$$\text{Dom}(|h|^{\frac{1}{2}}) = (\mathbb{1} + |h|)^{-\frac{1}{2}}\mathcal{W}. \quad (3.4)$$

For $w_1, w_2 \in \text{Dom}(|h|^{\frac{1}{2}})$, we can define $(w_i | h w_j)$. $\text{Dom}(|h|^{\frac{1}{2}})$ is a Hilbert space for the scalar product $(w_1 | (h + c + \mathbb{1}) w_2)$. We say that \mathcal{D} is a *form core for h* if it is a dense subspace of the form domain of h .

Lemma 3.1 *Suppose that $h \geq 0$ and \mathcal{D} is a form core of h . Then $\overset{\text{al}}{\Gamma}_s(\mathcal{D})$ is a form core of $d\Gamma(h)$.*

Proof. It is easy to see that

$$\mathbb{1} + d\Gamma(h) \leq \Gamma(\mathbb{1} + h). \quad (3.5)$$

Hence,

$$\Gamma(\mathbb{1} + h)^{-\frac{1}{2}}\Gamma_s(\mathcal{W}) \subset (\mathbb{1} + d\Gamma(h))^{-\frac{1}{2}}\Gamma_s(\mathcal{W}). \quad (3.6)$$

Let $\Psi \in (\mathbb{1} + d\Gamma(h))^{-\frac{1}{2}}\Gamma_s(\mathcal{W})$. Set

$$\Psi_n := \mathbb{1}_{[0,n]}(\Gamma(\mathbb{1} + h))\Psi. \quad (3.7)$$

By the spectral theorem and the fact that $\mathbb{1} + d\Gamma(h)$ and $\Gamma(\mathbb{1} + h)$ commute with one another, $\Psi_n \in \text{Dom}\Gamma(\mathbb{1} + h)^{\frac{1}{2}}$ and $\Psi_n \rightarrow \Psi$ in $(\mathbb{1} + d\Gamma(h))^{-\frac{1}{2}}\Gamma_s(\mathcal{W})$. Hence

$$\Gamma(\mathbb{1} + h)^{-\frac{1}{2}}\Gamma_s(\mathcal{W}) \text{ is dense in } (\mathbb{1} + d\Gamma(h))^{-\frac{1}{2}}\Gamma_s(\mathcal{W}). \quad (3.8)$$

Now \mathcal{D} is dense in $(\mathbb{1} + h)^{-\frac{1}{2}}\mathcal{W}$. Hence

$$\overset{\text{al}}{\Gamma}_s(\mathcal{D}) \text{ is dense in } \Gamma_s((\mathbb{1} + h)^{-\frac{1}{2}}\mathcal{W}) = \Gamma(\mathbb{1} + h)^{-\frac{1}{2}}\Gamma_s(\mathcal{W}). \quad (3.9)$$

Putting together (3.8) and (3.9) we obtain

$$\overset{\text{al}}{\Gamma}_s(\mathcal{D}) \text{ is dense in } (\mathbb{1} + d\Gamma(h))^{-\frac{1}{2}}\Gamma_s(\mathcal{W}). \quad (3.10)$$

But the RHS of (3.10) is the form domain of $d\Gamma(h)$. \square

3.4 Creation/annihilation operators

For any $w \in \mathcal{W}$, $\hat{a}(w)$ and $\hat{a}^*(w)$ denote the usual annihilation/creation operators

$$\hat{a}^*(w)\Psi := \sqrt{n+1}w \otimes_s \Psi, \quad \Psi \in \otimes_s^n \mathcal{W}, \quad (3.11)$$

$$\hat{a}(w)\Psi := \sqrt{n+1}(w | \otimes_s^1 \Psi, \quad \Psi \in \otimes_s^{n+1} \mathcal{W}, \quad (3.12)$$

These operators, originally well defined on $\Gamma_s^{\text{fin}}(\mathcal{W})$, extend to closed operators on $\Gamma_s(\mathcal{W})$. We set

$$\hat{\phi}(w, \bar{w}') := \hat{a}^*(w) + \hat{a}(w'). \quad (3.13)$$

Note that $\hat{\phi}(w, \bar{w})$ are self-adjoint. One can also introduce the so-called *Weyl operators* $e^{i\hat{\phi}(w, \bar{w})}$.

Remark 3.2 *Sometimes we may want to define creation/annihilation operators for w that do not belong to \mathcal{W} , but are functionals, possibly unbounded, with domain $\mathcal{D} \subset \mathcal{W}$. Then we can still define the annihilation operator $\hat{a}(w)$ by the formula (3.12), at least for $\Psi \in \overset{\text{al}}{\Gamma}_s(\mathcal{D})$. If w is unbounded, then $\hat{a}(w)$ is not closable and (3.11), the definition of $\hat{a}^*(w)$ as an operator, is incorrect. However, we can interpret both $\hat{a}(w)$ and $\hat{a}^*(w)$ as quadratic forms on $\overset{\text{al}}{\Gamma}_s(\mathcal{D})$.*

The following inequality is sometimes called the N_τ -estimate:

Proposition 3.3 *Let $h > 0$ and $w \in \mathcal{W}$. Then*

$$\|\hat{a}(w)\Phi\|^2 \leq (w|h^{-1}w)(\Phi|d\Gamma(h)\Phi). \quad (3.14)$$

Therefore,

$$\|d\Gamma(h)^{-\frac{1}{2}}\hat{a}^*(w)\| \leq \|h^{-\frac{1}{2}}w\|. \quad (3.15)$$

Proof. Clearly,

$$|w)(w| \leq (w|h^{-1}w)h. \quad (3.16)$$

Applying $d\Gamma$, we obtain

$$\hat{a}^*(w)\hat{a}(w) = d\Gamma(|w)(w|) \leq (w|h^{-1}w)d\Gamma(h). \quad (3.17)$$

□

Let $g \in \otimes_s^2 \mathcal{W}$. We define the annihilation and creation operators associated to g as follows:

$$\hat{a}^*(g)\Psi := \sqrt{n+2}\sqrt{n+1}g \otimes_s \Psi, \quad \Psi \in \otimes_s^n \mathcal{W}, \quad (3.18)$$

$$\hat{a}(g)\Psi := \sqrt{n+2}\sqrt{n+1}(g| \otimes \mathbb{1}^{\otimes n} \Psi, \quad \Psi \in \otimes_s^{n+2} \mathcal{W}. \quad (3.19)$$

Again, these operators, originally defined on $\Gamma_s^{\text{fin}}(\mathcal{W})$, extend to closed operators on $\Gamma_s(\mathcal{W})$:

Remark 3.4 *Again, if g does not belong to $\oplus_s^2 \mathcal{W}$, but is a functional with the domain $\overset{\text{al}}{\otimes}_s^2 \mathcal{D} \subset \otimes_s^2 \mathcal{W}$, then we can define $\hat{a}(g)$ and $\hat{a}^*(g)$ as quadratic forms on $\Gamma_s^{\text{al}}(\mathcal{D})$.*

It is important to note that each $g \in \otimes^2 \mathcal{W}$ defines a linear Hilbert-Schmidt operator from $\overline{\mathcal{W}}$ to \mathcal{W} , denoted by the same symbol g , by the identity

$$(w_1 \otimes w_2|g) = (w_2|g\chi w_1). \quad (3.20)$$

This provides an isometric isomorphism of $\otimes^2 \mathcal{W}$ with $B^2(\overline{\mathcal{W}}, \mathcal{W})$ —the space of Hilbert-Schmidt operators from $\overline{\mathcal{W}}$ to \mathcal{W} . Symmetric tensors (elements of $\otimes_s^2 \mathcal{W}$) are mapped onto symmetric operators (where the *symmetry* of g means $g = g^\#$).

Let us state the following fact about this identification:

Proposition 3.5 *Let $p_1, p_2 \in B(\mathcal{W})$. Then the tensor $p_1 \otimes p_2 g$ corresponds to the operator $p_1 g p_2^\#$.*

Proposition 3.6 *Let $w \in \mathcal{W}$, $h \in B(\mathcal{W})$, $g \in \mathcal{W} \otimes_s \mathcal{W}$. Then the following identities are true:*

$$[d\Gamma(h), \hat{a}^*(w)] = \hat{a}^*(hw), \quad [d\Gamma(h), \hat{a}(w)] = -\hat{a}(hw), \quad (3.21)$$

$$[\hat{a}(g), \hat{a}^*(w)] = 2\hat{a}^*(g\overline{w}), \quad [\hat{a}^*(g), \hat{a}(w)] = -2\hat{a}(g\overline{w}). \quad (3.22)$$

3.5 Symplectic and metaplectic transformations in infinite dimensions

As in (2.6) we introduce the operator

$$S = \begin{bmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{bmatrix}. \quad (3.23)$$

Let $R \in B(\mathcal{W} \oplus \overline{\mathcal{W}})$. As in Subsection 2.3, R is called symplectic if $R^*SR = S$. Bounded symplectic transformations form a group, which we denote $Sp(\mathcal{Y})$.

Various properties of symplectic operators described in Subsection 2.3 are valid in the present setting.

Theorem 3.7 *Let*

$$R = \begin{bmatrix} p & q \\ \bar{q} & \bar{p} \end{bmatrix} \in Sp(\mathcal{Y}). \quad (3.24)$$

Then the following conditions are equivalent:

(1) *There exists a unitary U such that*

$$\begin{aligned} U\hat{a}^*(w)U^* &= \hat{a}^*(pw) + \hat{a}(q\bar{w}), \\ U\hat{a}(w)U^* &= \hat{a}^*(q\bar{w}) + \hat{a}(pw), \quad w \in \mathcal{W}. \end{aligned} \quad (3.25)$$

(2) *There exists a unitary U such that*

$$Ue^{i\hat{\phi}(w,\bar{w})}U^* = e^{i\hat{\phi}(w',\bar{w}')}, \quad R \begin{bmatrix} w \\ \bar{w} \end{bmatrix} = \begin{bmatrix} w' \\ \bar{w}' \end{bmatrix}, \quad w \in \mathcal{W}. \quad (3.26)$$

(3) *There exists a *-automorphism α_R of $B(\Gamma_s(\mathcal{W}))$ such that*

$$\alpha_R(e^{i\hat{\phi}(w,\bar{w})}) = e^{i\hat{\phi}(w',\bar{w}')}, \quad R \begin{bmatrix} w \\ \bar{w} \end{bmatrix} = \begin{bmatrix} w' \\ \bar{w}' \end{bmatrix}, \quad w \in \mathcal{W}. \quad (3.27)$$

Let (1), (2) and (3) be true. Then U (common for (1) and (2)) is uniquely determined up to a phase factor. Besides, α_R is uniquely defined.

If R satisfies the conditions of the above theorem, then we say that R is *implementable*. The unitary U is called a (*Bogoliubov*) *implementer* of R . α_R is called the *Bogoliubov automorphism associated to R* .

We leave the proof of this theorem to the reader. Let us only mention that to show (3) \Rightarrow (2) we need to use Proposition A.1. To obtain the uniqueness of α_R we use the weak density of linear combinations of Weyl operators in $B(\Gamma_s(\mathcal{W}))$.

$Sp_{\text{res}}(\mathcal{Y})$ will denote the *restricted symplectic group*, which consists of $R \in Sp(\mathcal{Y})$ such that q is Hilbert-Schmidt. The importance of $Sp_{\text{res}}(\mathcal{Y})$ is due to the Shale Theorem [Sh], which we quote below in the form given in [DG].

Theorem 3.8 *$R \in Sp(\mathcal{Y})$. Then R is implementable iff $R \in Sp_{\text{res}}(\mathcal{Y})$. For such R , we can define the natural implementer of R*

$$U_R^{\text{nat}} := |\det pp^*|^{-\frac{1}{4}} e^{-\frac{1}{2}\hat{a}^*(d_2)} \Gamma((p^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_1)}, \quad (3.28)$$

where d_2, d_1 are defined as in (2.15) and (2.16). All implementers of $R \in Sp_{\text{res}}(\mathcal{Y})$ coincide with U_R^{nat} , up to a phase factor.

Bogoliubov implementers form a group, which is denoted $Mp^c(\mathcal{Y})$. We have a short exact sequence

$$\mathbb{1} \rightarrow U(1) \rightarrow Mp^c(\mathcal{Y}) \rightarrow Sp_{\text{res}}(\mathcal{Y}) \rightarrow \mathbb{1}.$$

Let us mention the following criterion, which was used in [NNS]:

Proposition 3.9 *If $R^*R - \mathbb{1}$ is Hilbert-Schmidt, then $R \in Sp_{\text{res}}(\mathcal{Y})$.*

Proof.

$$R^*R = \begin{bmatrix} p^*p + q^{\#}\bar{q} & p^*q + q^{\#}\bar{p} \\ p^{\#}\bar{q} + q^*p & p^{\#}\bar{p} + q^*q \end{bmatrix} = \begin{bmatrix} \mathbb{1} + 2q^{\#}\bar{q} & 2p^*q \\ 2p^{\#}\bar{q} & \mathbb{1} + 2q^*q \end{bmatrix}.$$

Now

$$\|\mathbb{1} - R^*R\|_2^2 = 8\mathrm{Tr}q^*q + 8\mathrm{Tr}q^*qq^*q + 8\mathrm{Tr}q^*pp^*q \quad (3.29)$$

$$= 16\mathrm{Tr}q^*qq^*q + 16\mathrm{Tr}q^*q \geq 16\mathrm{Tr}q^*q. \quad (3.30)$$

□

$Sp_{\mathrm{af}}(\mathcal{Y})$ will denote the *anomaly-free symplectic group*, which consists of $R \in Sp(\mathcal{Y})$ such that $\mathbb{1} - p$ is trace class [DG].

Proposition 3.10 $Sp_{\mathrm{af}}(\mathcal{Y})$ is a subgroup of $Sp_{\mathrm{res}}(\mathcal{Y})$.

Proof. We have

$$q^*q = p^*p - \mathbb{1} = (p^* - \mathbb{1})p + p - \mathbb{1}. \quad (3.31)$$

Therefore, $\|p - \mathbb{1}\|_1 < \infty$ implies $\|q\|_2 < \infty$. □

For $R \in Sp_{\mathrm{af}}(\mathcal{Y})$ we can define a pair of *metaplectic Bogoliubov implementers*

$$\pm U_R^{\mathrm{met}} := \pm(\det p^*)^{-\frac{1}{2}} e^{-\frac{1}{2}\hat{a}^*(d_2)} \Gamma((p^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_1)}. \quad (3.32)$$

They form a group, which we denote $Mp_{\mathrm{af}}(\mathcal{Y})$ [DG]. We have a short exact sequence

$$\mathbb{1} \rightarrow \mathbb{Z}_2 \rightarrow Mp_{\mathrm{af}}(\mathcal{Y}) \rightarrow Sp_{\mathrm{af}}(\mathcal{Y}) \rightarrow \mathbb{1}.$$

3.6 Classical quadratic Hamiltonians

In this subsection we consider strongly continuous 1-parameter groups of symplectic transformations. The following proposition describes their generators:

Proposition 3.11 Let iB be a generator of a 1-parameter group on $\mathcal{W} \oplus \overline{\mathcal{W}}$. The following statements are equivalent:

- (1) e^{iBt} , $t \in \mathbb{R}$, is a strongly continuous 1-parameter group of symplectic transformations.
- (2) iB is J -real, $SB^* \supset BS$.
- (3) $A_B := BS$ is J -real and $A_B^* \supset A_B$ (in other words, A_B is Hermitian).

Proof. We have for $w_1, w_2 \in \mathrm{Dom}(B)$

$$\left. \frac{d}{dt} (e^{itB} w_1 | S e^{itB} w_2) \right|_{t=0} = -i(Bw_1 | Sw_2) + i(w_1 | SBw_2). \quad (3.33)$$

Hence preservation of S by e^{itB} is equivalent to $(SAS)^* = B^*S \supset SB = SAS$, which means that SAS is Hermitian. This is equivalent to A being Hermitian. □

For brevity, we will say that B is a *symplectic generator* if iB generates a one-parameter group of symplectic transformations. Similarly as in the previous section, $A_B := BS$ will be sometimes called the *classical Hamiltonian of B* , and we will often write A instead of A_B .

Note that in finite dimensions the converse of Proposition 3.11 (3) is true: If A is Hermitian and J -real, then $B := AS$ is a symplectic generator. This is probably not the case in infinite dimensions.

3.7 Bogoliubov Hamiltonians

Let B be, as usual, a symplectic generator, and $A = BS$. We will write

$$e^{itB} = \begin{bmatrix} p_t & q_t \\ \bar{q}_t & \bar{p}_t \end{bmatrix}. \quad (3.34)$$

Theorem 3.12 *The following conditions are equivalent:*

- (1) *There exists a self-adjoint operator \hat{H} on $\Gamma_s(\mathcal{W})$ such that $e^{it\hat{H}}$ implements e^{itB} for any $t \in \mathbb{R}$.*
- (2) *There exists α_t , a 1-parameter C_0^* -group of $*$ -automorphisms of $B(\Gamma_s(\mathcal{W}))$, such that*

$$\alpha_t(e^{i\hat{\phi}(w, \bar{w})}) = e^{i\hat{\phi}(w_t, \bar{w}_t)}, \quad \begin{bmatrix} w_t \\ \bar{w}_t \end{bmatrix} = e^{itB} \begin{bmatrix} w \\ \bar{w} \end{bmatrix}, \quad w \in \mathcal{W}. \quad (3.35)$$

- (3) $\lim_{t \rightarrow 0} \|q_t\|_2 = 0$.

Let (1), (2), (3) be true. Then α_t is determined uniquely. \hat{H} is uniquely defined up to an additive constant.

\hat{H} will be called a *quantization* of B . We will also say that \hat{H} is a *quantum quadratic Hamiltonian*, or shorter, a *Bogoliubov Hamiltonian*. If the equivalent conditions of the above theorem are satisfied, then we will say that B possesses *quantizations*.

Proof. (1) \Leftrightarrow (2) is a consequence of Proposition A.2. We need to show that (1),(2) \Leftrightarrow (3)

If e^{itB} , $t \in \mathbb{R}$, is implementable, then $\|q_t\|_2 < \infty$, for all $t \in \mathbb{R}$.

If $\lim_{t \rightarrow 0} \|q_t\|_2 = 0$, then $\|q_t\|_2 < \infty$, for small enough t . But since $Sp_{\text{res}}(\mathcal{Y})$ is a group, $\|q_t\|_2 < \infty$, for all $t \in \mathbb{R}$.

Thus, in all cases (1), (2) and (3) we can define

$$U_t^{\text{nat}} := U_{e^{itB}}^{\text{nat}}, \quad (3.36)$$

(see (3.28)). Set

$$\alpha_t(C) := U_t^{\text{nat}} C U_{-t}^{\text{nat}}. \quad (3.37)$$

Clearly, $t \mapsto \alpha_t$ is a 1-parameter group of $*$ -automorphisms satisfying (3.35). The proof will be completed if we show the equivalence of the following statements:

- (i) $t \mapsto U_t^{\text{nat}}$ is strongly continuous at zero;
- (ii) $t \mapsto \alpha_t$ is a C_0^* -group of $*$ -automorphisms;
- (iii) $d_{2,t} := q_t \bar{p}_t^{-1}$ satisfies $\lim_{t \rightarrow 0} \|d_{2,t}\|_2 = 0$;
- (iv) $\lim_{t \rightarrow 0} \|q_t\|_2 = 0$.

(i) \Rightarrow (ii): We easily see that if $t \mapsto U_t^{\text{nat}}$ is strongly continuous at zero and if C is a bounded operator, then $t \mapsto U_t^{\text{nat}} C U_{-t}^{\text{nat}}$ is weakly continuous at zero. This implies that $t \mapsto \alpha_t$ is a C_0^* -group.

(ii) \Rightarrow (iii): Let $|\Omega\rangle\langle\Omega|$ denote the orthogonal projection onto Ω . We have

$$\begin{aligned} \left(\Omega | \alpha(|\Omega\rangle\langle\Omega|) \Omega \right) &= \left| (\Omega | U_t^{\text{nat}} \Omega) \right|^2 \\ &= \left| \det p_t p_t^* \right|^{-\frac{1}{2}} = \det (\mathbb{1} - d_{2,t}^* d_{2,t})^{\frac{1}{2}} \end{aligned} \quad (3.38)$$

$$= \exp \left(\frac{1}{2} \text{Tr} \log (\mathbb{1} - d_{2,t}^* d_{2,t}) \right). \quad (3.39)$$

In (3.38) we used the identity

$$p_t^{\#-1} p_t^{-1} = \mathbb{1} - d_{2,t}^* d_{2,t}. \quad (3.40)$$

(ii) implies that (3.39) goes to 1 for $t \rightarrow 0$. This is equivalent to $\lim_{t \rightarrow 0} \text{Tr} \log(\mathbb{1} - d_{2,t}^* d_{2,t}) = 0$, which is equivalent to $\lim_{t \rightarrow 0} \text{Tr} d_{2,t}^* d_{2,t} = 0$.

(iii) \Rightarrow (i): We have

$$U_t^{\text{nat}} e^{i\hat{\phi}(w, \bar{w})} \Omega = e^{i\hat{\phi}(w_t, \bar{w}_t)} |\det p_t p_t^*|^{-\frac{1}{4}} e^{-\frac{1}{2} a^*(d_{2,t})} \Omega. \quad (3.41)$$

But $t \mapsto e^{i\hat{\phi}(w_t, \bar{w}_t)}$ is strongly continuous. By (3.40), $\lim_{t \rightarrow 0} |\det p_t p_t^*|^{-\frac{1}{4}} = 1$. Besides, (iii) implies that $\lim_{t \rightarrow 0} e^{-\frac{1}{2} a^*(d_{2,t})} \Omega = \Omega$. Therefore, (3.41) is continuous at $t = 0$. But the span of $e^{i\hat{\phi}(w, \bar{w})} \Omega$ is dense and U_t^{nat} is unitary. Hence U_t^{nat} is strongly continuous at $t = 0$.

(iii) \Leftrightarrow (iv) follows from the identities

$$q_t q_t^* = d_{2,t}^* d_{2,t} (\mathbb{1} - d_{2,t}^* d_{2,t})^{-1}, \quad (3.42)$$

$$d_{2,t}^* d_{2,t} = q_t q_t^* (\mathbb{1} + q_t q_t^*)^{-1}. \quad (3.43)$$

□

Below we describe 3 distinguished quantizations.

- (1) If the group $e^{it\hat{H}}$ implementing e^{itB} is contained in $M_{\text{paf}}(\mathcal{Y})$, then \hat{H} will be called *Weyl*. It is easy to see that for a given symplectic generator B , its *Weyl quantization*, if it exists, is unique. We will denote it by \hat{H}_B^{w} . An alternative name for \hat{H}_B^{w} : the *symmetric quantization of B* .
- (2) We say that a quantization \hat{H} of B is *normally ordered*

$$\left. \frac{d}{dt} (\Omega | e^{it\hat{H}} \Omega) \right|_{t=0} = 0. \quad (3.44)$$

Again, a given symplectic generator B possesses at most one *normally ordered quantization*. We will denote it by \hat{H}_B^{n} . An alternative name for \hat{H}_B^{n} : the *Wick quantization of B* .

- (3) If B possess a quantization, which is bounded from below, then all of its quantizations are bounded from below. Then one can introduce the *zero-infimum quantization* \hat{H}_B^{z} fixed by the condition

$$\inf \hat{H}_B^{\text{z}} = 0.$$

Let us stress that there exist B that possess quantizations, but they do not possess \hat{H}_B^{w} , \hat{H}_B^{n} or \hat{H}_B^{z} . We will usually drop the subscript B in the above symbols.

Note that whereas the definitions of \hat{H}^{w} and \hat{H}^{z} are quite obvious, it is less clear how to generalize the concept of normally ordered Bogoliubov Hamiltonian to infinite dimensions. In the following proposition we formulate another condition, which could be considered as another candidate for a definition of \hat{H}^{n} .

Proposition 3.13 *Suppose that B possesses a quantization \hat{H} such that $\Omega \in \text{Dom}(|\hat{H}|^{\frac{1}{2}})$ (the vacuum belongs to the form domain of \hat{H}). Then B possesses the normally ordered quantization.*

Proof. We easily check that

$$\hat{H}^{\text{n}} := \hat{H} - (\Omega | \hat{H} \Omega). \quad (3.45)$$

satisfies (3.44). □

Theorem 3.14 *Consider (3.34).*

(1) The condition

$$\lim_{t \rightarrow 0} \|p_t - \mathbb{1}\|_1 = 0. \quad (3.46)$$

is equivalent to B possessing the Weyl quantization \hat{H}^w . If this is the case, then

$$e^{it\hat{H}^w} = (\det p_t^*)^{-\frac{1}{2}} e^{-\frac{1}{2}\hat{a}^*(d_{2,t})} \Gamma((p_t^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_{1,t})}, \quad (3.47)$$

where the sign of the square root is determined by continuity.

(2) Suppose that there exists a self-adjoint operator h on \mathcal{W} such that

$$\lim_{t \rightarrow 0} \frac{\|e^{-ith} p_t - \mathbb{1}\|_1}{t} = 0. \quad (3.48)$$

Then B possesses the normally ordered quantization \hat{H}^n and

$$e^{it\hat{H}^n} = (\det p_t^* e^{ith})^{-\frac{1}{2}} e^{-\frac{1}{2}\hat{a}^*(d_{2,t})} \Gamma((p_t^*)^{-1}) e^{\frac{1}{2}\hat{a}(d_{1,t})}, \quad (3.49)$$

where the sign of the square root is determined by continuity. The operator h that appears in (3.48) is uniquely defined.

(3) Suppose that the assumptions of (2) hold. In addition, assume that h in (3.48) is trace class. Then B possesses both normally ordered and Weyl quantization, and

$$\hat{H}^n + \frac{1}{2} \text{Tr} h = \hat{H}^w. \quad (3.50)$$

Proof. (1): $\lim_{t \rightarrow 0} \|\mathbb{1} - p_t\|_1 = 0$ implies that $e^{itB} \in Sp_{\text{af}}(\mathcal{Y})$ at least for small t . But $Sp_{\text{af}}(\mathcal{Y})$ is a group. Therefore, $e^{itB} \in Sp_{\text{af}}(\mathcal{Y})$ for all $t \in \mathbb{R}$.

Besides, $t \mapsto e^{itB}$ is continuous in the topology of $Sp_{\text{af}}(\mathcal{Y})$ at zero. By the group property of $Sp_{\text{af}}(\mathcal{Y})$, it is continuous for all $t \in \mathbb{R}$.

Hence, U_t^{met} given by (3.47) is well defined. U_t^{met} obviously is one of the metaplectic implementers of e^{itB} . We have

$$(\Omega | U_t^{\text{met}} \Omega) = (\det p_t^*)^{-\frac{1}{2}} \quad (3.51)$$

depends continuously on t .

Using that $Mp(\mathcal{Y})$ is a group and the continuity of (3.51), we see that U_t^{met} satisfies the group property. Next, repeating the argument of the proof of Thm 3.12, we see that U_t^{met} is continuous on coherent vectors.

Thus U_t^{met} is a strongly continuous group of metaplectic implementers of e^{itB} . Hence B possesses the Weyl quantization.

Conversely, if B possesses the Weyl quantization \hat{H}^w , then $U_t^{\text{met}} = e^{it\hat{H}^w}$. Then (3.51) is true. But $\lim_{t \rightarrow 0} \|\mathbb{1} - p_t\|_1 = 0$ is equivalent to the continuity of the rhs of (3.51).

(2): (3.48) implies

$$\lim_{t \rightarrow 0} \|e^{-ith} p_t - \mathbb{1}\|_1 = 0. \quad (3.52)$$

Therefore, the identity

$$q_t^* q_t = p_t^* p_t - \mathbb{1} = (p_t^* e^{ith} - \mathbb{1}) e^{-ith} p_t + e^{-ith} p_t - \mathbb{1} \quad (3.53)$$

shows that $\lim_{t \rightarrow 0} \|q_t\|_2 = 0$. Therefore, (3.49) is well defined and depends continuously on t . Clearly,

$$|\det p_t^* e^{ith}|^2 = \det p_t^* p_t. \quad (3.54)$$

Hence, (3.49) differs from $U_{e^{itB}}^{\text{nat}}$ by a phase factor. We check by direct calculation that it satisfies the group property [BD].

Using (3.52) and the differentiability of the determinant in the trace norm, for small enough t we have

$$|\det p_t^* e^{ith} - 1| \leq c \|p_t^* e^{ith} - \mathbb{1}\|_1. \quad (3.55)$$

Hence (3.48) implies

$$\lim_{t \rightarrow 0} \frac{\det p_t^* e^{ith} - 1}{t} = 0. \quad (3.56)$$

Therefore, using also (3.52),

$$\lim_{t \rightarrow 0} \frac{(\det p_t^* e^{ith})^{-\frac{1}{2}} - 1}{t} = 0. \quad (3.57)$$

By (3.49),

$$(\Omega | e^{it\hat{H}^n} \Omega) = (\det p_t^* e^{ith})^{-\frac{1}{2}}. \quad (3.58)$$

Hence, (3.44) is true.

Suppose that for h_1, h_2 we have (3.48). Let $w, w' \in \mathcal{W}$ be normalized. Then

$$\frac{1}{t} |(w | e^{ith_1} w') - (w | e^{ith_2} w')| \leq \frac{1}{t} \|e^{ith_1} - e^{ith_2}\| \quad (3.59)$$

$$\leq \frac{1}{t} \|e^{ith_1} - e^{ith_2}\|_1 \quad (3.60)$$

$$\leq \frac{1}{t} \|e^{ith_1} - p_t\|_1 + \frac{1}{t} \|p_t - e^{ith_2}\|_1 \rightarrow 0. \quad (3.61)$$

Hence $h_1 = h_2$ by Lemma A.4.

(3): Using $\|h\|_1 < \infty$, we can write

$$\det p_t = \det e^{ith} \det e^{-ith} p_t = e^{it\text{Tr}h} \det e^{-ith} p_t. \quad (3.62)$$

Thus we see that both (3.47) and (3.49) are well defined and

$$e^{it\hat{H}^w} = e^{it\frac{1}{2}\text{Tr}h} e^{it\hat{H}^n}. \quad (3.63)$$

□

3.8 Criteria for existence of quantizations of classical Hamiltonians

In this subsection we restrict our study to symplectic generators that are bounded perturbations of diagonal symplectic generators.

We will always assume that h is a self-adjoint operator on \mathcal{W} and $g = g^\#$. Besides,

$$B := \begin{bmatrix} h & -g \\ \bar{g} & -\bar{h} \end{bmatrix}, \quad B_0 := \begin{bmatrix} h & 0 \\ 0 & -\bar{h} \end{bmatrix}, \quad (3.64)$$

$$A = BS = \begin{bmatrix} h & g \\ \bar{g} & \bar{h} \end{bmatrix}, \quad A_0 = B_0 S = \begin{bmatrix} h & 0 \\ 0 & \bar{h} \end{bmatrix}, \quad G = \begin{bmatrix} 0 & -g \\ \bar{g} & 0 \end{bmatrix}. \quad (3.65)$$

The following proposition is immediate:

Proposition 3.15 *If g is bounded, then B is a symplectic generator. Besides, $A_B = BS$ is self-adjoint.*

Proof. Clearly, B_0 is a symplectic generator and A_0 is self-adjoint. We can add a bounded perturbation without destroying these properties. \square

The following theorem is a slightly strengthened version of a criterion due to Berezin [Be], see also [BD]. Throughout the subsection we set

$$f(t) := \int_0^t e^{ish} g e^{is\bar{h}} ds. \quad (3.66)$$

- Theorem 3.16** (1) *Suppose that g is bounded and $\lim_{t \rightarrow 0} \|f(t)\|_2 = 0$. Then B possesses quantizations.*
(2) *In addition to assumptions of (1) suppose that $\lim_{t \rightarrow 0} \|\bar{g}f(t)\|_1 = 0$. Then B possesses the normally ordered quantization.*
(3) *In addition to assumptions of (2) suppose that $\|h\|_1 < \infty$. Then B possesses both the Weyl and the normally ordered quantizations, and*

$$\hat{H}^n + \frac{1}{2} \text{Tr}h = \hat{H}^w. \quad (3.67)$$

Proof. (1): Using repeatedly the identity

$$f(2t) = f(t) + e^{it\bar{h}} f(t) e^{it\bar{h}} \quad (3.68)$$

we see that $\|f(t)\|_2$ is finite for all t .

Set

$$V(t) := e^{itB} e^{-itB_0}, \quad (3.69)$$

$$G(t) := e^{itB_0} G e^{-itB_0} = \begin{bmatrix} 0 & -e^{ish} g e^{is\bar{h}} \\ e^{-is\bar{h}} \bar{g} e^{-ish} & 0 \end{bmatrix}, \quad (3.70)$$

$$F(t) := \int_0^t G(s) ds = \begin{bmatrix} 0 & -f(t) \\ f(t) & 0 \end{bmatrix}. \quad (3.71)$$

From

$$V(t) = \mathbb{1} + i \int_0^t V(s) G(s) ds \quad (3.72)$$

and $\|G(t)\| = \|G\|$, we obtain

$$\|V(t)\| \leq e^{|t|\|G\|}. \quad (3.73)$$

Iterating (3.72) gives

$$V(t) = \mathbb{1} + iF(t) - \int_0^t V(s) G(s) e^{isB_0} F(t-s) e^{-isB_0} ds. \quad (3.74)$$

Therefore,

$$\|V(t) - \mathbb{1}\|_2 \leq \|F(t)\|_2 + \int_0^t \|V(s)\| \|G\| \|F(t-s)\|_2 ds. \quad (3.75)$$

But $\|F(t)\|_2 = \sqrt{2}\|f(t)\|_2$, $\|G\| = \|g\|$. Hence $\|V(t) - \mathbb{1}\|_2$ is finite and goes to zero as $t \rightarrow 0$. Arguing as in Proposition 3.9, we obtain

$$16\|q(t)\|_2^2 \leq \|V(t) - \mathbb{1}\|_2. \quad (3.76)$$

Therefore, $\|q(t)\|_2$ is finite and goes to zero as $t \rightarrow 0$. This means that the assumption of Theorem 3.12 (3) is satisfied. Hence B possesses quantizations.

(2): We rewrite (3.74) as

$$\begin{bmatrix} p_t - e^{ith} & q_t + if(t)e^{-it\bar{h}} \\ \bar{q}_t - i\bar{f}(t)e^{ith} & \bar{p}_t - e^{-it\bar{h}} \end{bmatrix} = e^{itB} - e^{itB_0} - iF(t)e^{itB_0} \quad (3.77)$$

$$= - \int_0^t V(s)G(s)e^{isB_0}F(t-s)e^{i(t-s)B_0}ds. \quad (3.78)$$

Therefore, by (A.9),

$$2\|p_t - e^{ith}\|_1 \leq \|e^{itB} - e^{itB_0} - iF(t)e^{itB_0}\|_1 \quad (3.79)$$

$$\leq \int_0^t \|V(s)\| \|GF(t-s)e^{i(t-s)B_0}\|_1 ds. \quad (3.80)$$

Using $\|GF(t)\|_1 = 2\|\bar{g}f(t)\|_1$ and $\lim_{t \rightarrow 0} \|\bar{g}f(t)\|_1 = 0$, we see that (3.80) is $o(t)$. Thus we obtain $\|p_t - e^{ith}\|_1 = o(t)$. This means that the assumption of Theorem 3.14 (2) is satisfied. Hence, B possesses the normally ordered quantization.

(3): We apply Theorem 3.14 (3). \square

The assumptions of Theorem 3.16 are not very convenient to verify. Our next aim is to formulate criteria for the existence of quantizations, which are more convenient to check.

Define

$$\gamma(g) := (h \otimes \mathbb{1} + \mathbb{1} \otimes h)^{-1}g, \quad (3.81)$$

where we use the tensor interpretation of g and assume that $g \in \text{Dom}(h \otimes \mathbb{1} + \mathbb{1} \otimes h)^{-1}$.

Proposition 3.17 *In the operator interpretation, $\gamma(g)$ corresponds to*

$$\gamma(g) = i \lim_{\epsilon \searrow 0} \int_0^\infty e^{-\epsilon t} e^{-ith} g e^{-it\bar{h}} dt \quad (3.82)$$

and satisfies

$$h\gamma(g) + \gamma(g)\bar{h} = g. \quad (3.83)$$

For $h > 0$ we can “Wick rotate” the formula (3.82) and write

$$\gamma(g) = \int_0^\infty e^{-th} g e^{-t\bar{h}} dt. \quad (3.84)$$

Proof. By Prop. 3.5 and $\bar{h} = h^\#$, we can identify the operator $e^{-ith} g e^{-it\bar{h}}$ with the tensor

$$e^{-ith} \otimes e^{-it\bar{h}} g = e^{-it(h \otimes \mathbb{1} + \mathbb{1} \otimes h)} g. \quad (3.85)$$

Clearly,

$$i \int_0^\infty e^{-\epsilon t} e^{-it(h \otimes \mathbb{1} + \mathbb{1} \otimes h)} g = (h \otimes \mathbb{1} + \mathbb{1} \otimes h - i\epsilon)^{-1} g \rightarrow (h \otimes \mathbb{1} + \mathbb{1} \otimes h)^{-1} g, \quad (3.86)$$

where we use the usual Hilbert space convergence, which proves (3.82).

Set

$$\gamma_\epsilon(g) := i \int_0^\infty e^{-\epsilon s} e^{-ish} g e^{-is\bar{h}} ds. \quad (3.87)$$

We compute:

$$e^{-ith}\gamma_\epsilon(g)e^{-it\bar{h}} = i \int_t^\infty e^{-\epsilon(s-t)} e^{-ish} g e^{-is\bar{h}} ds \quad (3.88)$$

$$= -i \int_0^t e^{-ish} g e^{-is\bar{h}} e^{-\epsilon s} ds + e^{\epsilon t} \gamma_\epsilon(g). \quad (3.89)$$

We differentiate wrt t at $t = 0$, obtaining

$$-i(h\gamma_\epsilon(g) + \gamma_\epsilon(g)\bar{h}) = -ig + \epsilon\gamma_\epsilon(g). \quad (3.90)$$

Taking the limit as $\epsilon \searrow 0$, we obtain (3.83).

The proof of (3.84) is almost the same as that of (3.82). \square

We will also write

$$\gamma(G) := \begin{bmatrix} 0 & -\gamma(g) \\ \gamma(g) & 0 \end{bmatrix}. \quad (3.91)$$

Note that in the operator interpretation we have

$$[B_0, \gamma(G)] = G. \quad (3.92)$$

The following criterion is a consequence of Theorem 3.16.

Theorem 3.18 (1) *Suppose that g is bounded and $g = g_1 + g_2$, where $\|g_1\|_2 < \infty$ and $\|\gamma(g_2)\|_2 < \infty$. Then the assumptions of Theorem 3.16 are satisfied, and hence B possesses quantizations.*

(2) *Suppose that $\|g\|_2 < \infty$. Then B possesses the normally ordered quantization.*

(3) *Suppose that $\|h\|_1 < \infty$ and $\|g\|_2 < \infty$. Then B possesses both the Weyl and the normally ordered quantization. Besides,*

$$\hat{H}^w = \hat{H}^n + \text{Tr}h. \quad (3.93)$$

Proof. (1): Set

$$f_i(t) := \int_0^t e^{ish} g_i e^{is\bar{h}} ds. \quad (3.94)$$

It is clear that $\lim_{t \rightarrow 0} \|f_1(t)\|_2 = 0$. The fact that $\lim_{t \rightarrow 0} \|f_2(t)\|_2 = 0$ follows from

$$\int_0^t e^{ish} g_2 e^{is\bar{h}} ds = -i \int_0^t \frac{d}{ds} e^{ish} \gamma(g_2) e^{is\bar{h}} ds = -ie^{ith} \gamma(g_2) e^{it\bar{h}} + i\gamma(g_2). \quad (3.95)$$

Hence assumptions of Thm 3.16 (1) are satisfied.

(2): Clearly, $\|\bar{g}f(t)\|_1 \leq t\|g\|_2^2$. Hence assumptions of Thm 3.16 (2) are satisfied.

Now (3) follows immediately from Thm 3.16 (3). \square

Note that the assumption of Theorem 3.21 implies that of Theorem 3.18:

Proposition 3.19 *$h > 0$ and $\|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\|_2 < \infty$ implies $\|\gamma(g)\|_2 < \infty$.*

Proof. $h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}$ corresponds to $h^{-\frac{1}{2}} \otimes h^{-\frac{1}{2}} g$ in the tensor interpretation. Clearly,

$$2h \otimes h \leq (h \otimes \mathbf{1} + \mathbf{1} \otimes h)^2. \quad (3.96)$$

Hence,

$$h^{-1} \otimes h^{-1} \geq 2(h \otimes \mathbf{1} + \mathbf{1} \otimes h)^{-2}. \quad (3.97)$$

Therefore,

$$\|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\|_2 = \|h^{-\frac{1}{2}} \otimes h^{-\frac{1}{2}} g\| \geq \sqrt{2} \|(h \otimes \mathbf{1} + \mathbf{1} \otimes h)^{-1} g\| = \sqrt{2} \|\gamma(g)\|_2. \quad (3.98)$$

\square

3.9 Positive classical Hamiltonians and their diagonalization

The following theorem is an extension of Theorem 2.3 to arbitrary dimensions. It says that a large class of classical Hamiltonians can be diagonalized by a positive symplectic transformation. This theorem is implicitly contained in [DG] (see Thm 11.20 (3) together with Thm 18.5 (3)). [NNS] contains also a related result about the diagonalizability of classical Hamiltonians. It does not provide, however, a construction of a distinguished diagonalizing operator.

We will use the notation introduced in (3.64) and (3.65). We will assume that $h > 0$. It will not be necessary to assume that g is bounded—we will assume that $g = g^\#$ is a bilinear form with the right domain $\text{Dom}|\bar{h}|^{\frac{1}{2}}$ and the left domain $\text{Dom}|h|^{\frac{1}{2}}$.

Theorem 3.20 *Let h be positive and*

$$\|h^{-\frac{1}{2}}g\bar{h}^{-\frac{1}{2}}\| =: a < 1. \quad (3.99)$$

Then A is a positive self-adjoint operator with the form domain $\text{Dom}A_0^{\frac{1}{2}}$. The corresponding B is a symplectic generator.

Besides,

$$R_0 = SA^{-\frac{1}{2}}(A^{\frac{1}{2}}SASA^{\frac{1}{2}})^{\frac{1}{2}}A^{-\frac{1}{2}}S, \quad (3.100)$$

is a bounded invertible positive symplectic operator. Hence so is

$$R = R_0^{\frac{1}{2}} \quad (3.101)$$

R diagonalizes B and A , that is, for some positive self-adjoint h_{dg}

$$B = R \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & -h_{\text{dg}} \end{bmatrix} R^{-1}, \quad (3.102)$$

$$A = R \begin{bmatrix} h_{\text{dg}} & 0 \\ 0 & h_{\text{dg}} \end{bmatrix} R^*. \quad (3.103)$$

Moreover,

$$\frac{(1-a)^{\frac{1}{4}}}{(1+a)^{\frac{1}{4}}} \leq \|R\| \leq \frac{(1+a)^{\frac{1}{4}}}{(1-a)^{\frac{1}{4}}}. \quad (3.104)$$

Proof. GS is a form bounded perturbation of A_0 :

$$|(v|GSv)| \leq a(v|A_0v), \quad v \in \text{Dom}(A_0^{\frac{1}{2}}).$$

Therefore, A extends to a positive self-adjoint operator by the KLMN Theorem.

A satisfies

$$(1-a)A_0 \leq A \leq (1+a)A_0. \quad (3.105)$$

Similarly,

$$SAS = A_0 - GS$$

extends to a positive operator satisfying

$$(1-a)A_0 \leq SAS \leq (1+a)A_0. \quad (3.106)$$

Therefore

$$A^{\frac{1}{2}}SASA^{\frac{1}{2}} \geq (1-a)A^{\frac{1}{2}}A_0A^{\frac{1}{2}} \geq \frac{(1-a)}{(1+a)}A^2. \quad (3.107)$$

$$A^{\frac{1}{2}}SASA^{\frac{1}{2}} \leq (1+a)A^{\frac{1}{2}}A_0A^{\frac{1}{2}} \leq \frac{(1+a)}{(1-a)}A^2. \quad (3.108)$$

Hence,

$$\frac{\sqrt{1-a}}{\sqrt{1+a}}A \leq (A^{\frac{1}{2}}SASA^{\frac{1}{2}})^{\frac{1}{2}} \leq \frac{\sqrt{1+a}}{\sqrt{1-a}}A. \quad (3.109)$$

Thus R_0 , defined by (3.100), is a well defined bounded invertible positive operator. Hence so is R .

Repeating the arguments of the proof of Theorem 2.3, we obtain (3.102) and (3.103). By (3.102) we have

$$e^{itB} = R \begin{bmatrix} e^{ith_{\text{dg}}} & 0 \\ 0 & e^{-it\overline{h_{\text{dg}}}} \end{bmatrix} R^{-1}. \quad (3.110)$$

(3.110) is clearly symplectic. Hence B is a symplectic generator. \square

For further use we note that we can rewrite (3.100) as follows:

$$R_0 = SA^{-\frac{1}{2}} \left(\int \frac{\tau^2}{(\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}})} \frac{d\tau}{2\pi} \right) A^{-\frac{1}{2}}S.$$

As a side remark note that $h > 0$ and (3.99) not only imply $A > 0$, but the converse implication is ‘‘almost true’’. More precisely, set $\mathcal{W}_0 := (\text{Ker}h)^\perp$. Then $A \geq 0$ is equivalent to the following condition:

- (1) $h \geq 0$,
- (2) $\text{Ker}g \supset \mathcal{W}_0^\perp$ (and hence, since $g = g^\#$, we have $\text{Rang}g \subset \mathcal{W}_0$)
- (3) $\|h^{-\frac{1}{2}}g\overline{h}^{-\frac{1}{2}}\| \leq 1$, in the sense of operators from \mathcal{W}_0 .

3.10 Implementable diagonalizability of positive Hamiltonians

The following theorem is due to [NNS]. The proof that we present below follows closely that of [NNS], with only minor modifications.

Theorem 3.21 *In addition to the assumptions of Thm 3.20, suppose that*

$$\|h^{-\frac{1}{2}}g\overline{h}^{-\frac{1}{2}}\|_2 < \infty. \quad (3.111)$$

Let R be the symplectic operator given by Thm 3.20 and q be given by (3.24). Then

$$\|q\|_2 \leq 2 \frac{1}{(1-a)} \|h^{-\frac{1}{2}}g\overline{h}^{-\frac{1}{2}}\|_2. \quad (3.112)$$

In particular, $R \in Sp_{\text{res}}(\mathcal{Y})$ and hence R is implementable.

Under the assumptions of the above theorem, R possesses a Bogoliubov implementer U . If h_{dg} is given by (3.102), then

$$Ud\Gamma(h_{\text{dg}})U^* \quad (3.113)$$

is the zero-infimum quantization of B . (3.113) possesses a ground state (its infimum is an eigenvalue).

Proof of Thm 3.21. We start from estimating $R^*R - \mathbb{1} = R^2 - \mathbb{1} = R_0 - \mathbb{1}$. We have

$$\begin{aligned}
& S(R_0 - \mathbb{1})S \\
&= A^{-\frac{1}{2}}(A^{\frac{1}{2}}SASA^{\frac{1}{2}})^{\frac{1}{2}}A^{-\frac{1}{2}} - A^{-\frac{1}{2}}(A^2)^{\frac{1}{2}}A^{-\frac{1}{2}} \\
&= \int \frac{d\tau}{2\pi} A^{-\frac{1}{2}} \left(\frac{A^{\frac{1}{2}}SASA^{\frac{1}{2}}}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} - \frac{A^2}{\tau^2 + A^2} \right) A^{-\frac{1}{2}} \\
&= - \int \frac{\tau^2 d\tau}{2\pi} A^{-\frac{1}{2}} \left(\frac{1}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} - \frac{1}{\tau^2 + A^2} \right) A^{-\frac{1}{2}} \\
&= \int \frac{\tau^2 d\tau}{2\pi} A^{\frac{1}{2}} \frac{1}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} A^{-\frac{1}{2}} (SAS - A) \frac{1}{\tau^2 + A^2} \\
&=: \int \frac{\tau^2 d\tau}{2\pi} T(\tau).
\end{aligned}$$

Now, for any $\epsilon > 0$,

$$\begin{aligned}
& \pm 2T(\tau) \\
&= \pm 2A^{-\frac{1}{2}} \frac{1}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} A^{\frac{1}{2}} (SAS - A) \frac{1}{\tau^2 + A^2} \\
&\leq \epsilon^{-1} \frac{1}{\tau^2 + A^2} (SAS - A) A^{\frac{1}{2}} \frac{1}{A^{\frac{1}{2}}SASA^{\frac{1}{2}}} A^{\frac{1}{2}} (SAS - A) \frac{1}{\tau^2 + A^2} \\
&\quad + \epsilon A^{-\frac{1}{2}} \frac{A^{\frac{1}{2}}SASA^{\frac{1}{2}}}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} A^{-\frac{1}{2}} \\
&=: \epsilon^{-1} T_1(\tau) + \epsilon T_2(\tau).
\end{aligned}$$

We deal with the second term:

$$\begin{aligned}
& \int \frac{\tau^2 d\tau}{2\pi} T_2(\tau) \\
&= \int \frac{\tau^2 d\tau}{2\pi} A^{-\frac{1}{2}} \frac{A^{\frac{1}{2}}SASA^{\frac{1}{2}}}{\tau^2 + A^{\frac{1}{2}}SASA^{\frac{1}{2}}} A^{-\frac{1}{2}} \\
&= A^{-\frac{1}{2}} (A^{\frac{1}{2}}SASA^{\frac{1}{2}})^{\frac{1}{2}} A^{-\frac{1}{2}} \\
&= R_0.
\end{aligned}$$

Next we treat the first term:

$$\begin{aligned}
K &:= \int \frac{\tau^2 d\tau}{2\pi} T_1(\tau) \\
&= \int \frac{\tau^2 d\tau}{2\pi} \frac{1}{\tau^2 + A^2} (SAS - A) S \frac{1}{A} S (SAS - A) \frac{1}{\tau^2 + A^2}.
\end{aligned}$$

We have,

$$\text{Tr}K = \int \frac{\tau^2 d\tau}{2\pi} \frac{1}{(\tau^2 + A^2)^2} (SAS - A) S \frac{1}{A} S (SAS - A) \quad (3.114)$$

$$= \text{Tr} \frac{1}{A} (SAS - A) S \frac{1}{A} S (SAS - A) \quad (3.115)$$

$$\leq \frac{1}{(1-a)^2} \text{Tr} \frac{1}{A_0} (SAS - A) \frac{1}{A_0} (SAS - A) \quad (3.116)$$

$$= 8 \frac{1}{(1-a)^2} \text{Tr} \bar{h}^{-1} \bar{g} h^{-1} g. \quad (3.117)$$

Thus we proved that

$$\pm 2S(R_0 - \mathbb{1})S \leq \epsilon^{-1}K + \epsilon SR_0S, \quad (3.118)$$

where K is positive operator with trace bounded by (3.117). We rewrite (3.118) with sign + as

$$(2 - \epsilon)S(R_0 - \mathbb{1})S \leq \epsilon^{-1}K + \epsilon. \quad (3.119)$$

Let $s_n(C)$ denote the n th singular value of an operator C , that means, the n th eigenvalue of $|C| := \sqrt{C^*C}$ in the descending order. We will write for brevity $\lambda_n := s_n(|q|^2)$.

Using U defined in (2.21), we have

$$R_0 - \mathbb{1} = R^2 - \mathbb{1} = 2U \begin{bmatrix} u(|q|^2 + |q|\sqrt{\mathbb{1} + |q|^2})u^* & 0 \\ 0 & |q|^2 - |q|\sqrt{\mathbb{1} + |q|^2} \end{bmatrix} U^*. \quad (3.120)$$

Therefore,

$$s_n(R_0 - \mathbb{1}) = 2(\lambda_n + \sqrt{\lambda_n + \lambda_n^2}). \quad (3.121)$$

Thus,

$$2(2 - \epsilon)(\lambda_n + \sqrt{\lambda_n + \lambda_n^2}) \leq \epsilon^{-1}s_n(K) + \epsilon. \quad (3.122)$$

Let c be an arbitrary positive number. Let

$$\lambda_n \leq c. \quad (3.123)$$

Clearly, (3.122) implies

$$2(2 - \epsilon)\sqrt{\lambda_n} \leq \epsilon^{-1}s_n(K) + \epsilon. \quad (3.124)$$

Taking into account (3.123), we obtain

$$4\sqrt{\lambda_n} \leq \epsilon^{-1}s_n(K) + \epsilon(1 + 2\sqrt{c}). \quad (3.125)$$

Optimizing wrt ϵ , we obtain

$$4\sqrt{\lambda_n} \leq 2\sqrt{s_n(K)}\sqrt{1 + 2\sqrt{c}}. \quad (3.126)$$

Hence,

$$\lambda_n \leq s_n(K) \frac{1 + 2\sqrt{c}}{4}. \quad (3.127)$$

Let

$$c \leq \lambda_n. \quad (3.128)$$

Clearly, (3.122) implies

$$4(2 - \epsilon)\lambda_n \leq \epsilon^{-1}s_n(K) + \epsilon. \quad (3.129)$$

Taking into account (3.128), we obtain

$$\lambda_n \leq \frac{1}{\epsilon(8 - \epsilon(4 + c^{-1}))} s_n(K). \quad (3.130)$$

Optimizing wrt ϵ , we obtain

$$\lambda_n \leq \frac{4 + c^{-1}}{16} s_n(K). \quad (3.131)$$

Setting $c = \frac{1}{4}$ in (3.127) and (3.131), we obtain

$$\lambda_n \leq \frac{1}{2} s_n(K). \quad (3.132)$$

Hence,

$$\|q\|_2^2 = \sum_{n=1}^{\infty} \lambda_n \leq \frac{1}{2} \sum_{n=1}^{\infty} s_n(K) = \frac{1}{2} \text{Tr} K. \quad (3.133)$$

This together with (3.117) yields (3.112). \square

3.11 Normally ordered Hamiltonian

In this subsection we give conditions on B that guarantee the existence of a bounded from below normally ordered quantization. We follow [NNS], whose approach is based on quadratic forms. Similar results were contained in [BD]. They were however weaker, since only operator bounded perturbations were used in [BD].

Suppose that $\Phi, \Psi \in \Gamma_s(\mathcal{W})$. Define the *reduced 1-body density operator* $\gamma_{\Psi, \Phi}$ and the *pairing operator* $\alpha_{\Psi, \Phi}$ as follows:

$$\begin{aligned} (w_1 | \gamma_{\Psi, \Phi} w_2) &:= (\Phi | \hat{a}^*(w_2) \hat{a}(w_1) \Psi), \\ (\alpha_{\Phi, \Psi} \bar{w}_2 | w_1) = (\alpha_{\Phi, \Psi} | w_1 \otimes w_2) &:= (\Phi | \hat{a}^*(w_2) \hat{a}^*(w_1) \Psi), \quad w_1, w_2 \in \mathcal{W}. \end{aligned}$$

(Note that, as usual for similar objects, $\alpha_{\Psi, \Phi}$ has two interpretations: as a symmetric operator from $\overline{\mathcal{W}}$ to \mathcal{W} , or as an element of the Hilbert space $\otimes_s^2 \mathcal{W}$. We will treat the former interpretation as the standard one).

We will write

$$\gamma_{\Phi} := \gamma_{\Phi, \Phi}, \quad \alpha_{\Phi} := \alpha_{\Phi, \Phi}.$$

Note that

$$\alpha_{\Phi, \Psi}^{\#} = \alpha_{\Phi, \Psi}, \tag{3.134}$$

$$\begin{bmatrix} \gamma_{\Phi} & \alpha_{\Phi} \\ \bar{\alpha}_{\Phi} & \mathbb{1} + \bar{\gamma}_{\Phi} \end{bmatrix} \geq 0. \tag{3.135}$$

For further use note that (3.135) is equivalent to

$$\gamma_{\Phi} \geq 0, \quad \gamma_{\Phi} \geq \alpha_{\Phi} (\mathbb{1} + \bar{\gamma}_{\Phi})^{-1} \bar{\alpha}_{\Phi}. \tag{3.136}$$

Clearly, if h is an operator on \mathcal{W} and $g \in \otimes_s^2 \mathcal{W}$, then

$$(\Phi | d\Gamma(h) \Psi) = \text{Tr} \gamma_{\Phi, \Psi} h, \tag{3.137}$$

$$(\Phi | \hat{a}^*(g) \Psi) = \text{Tr} \alpha_{\Phi, \Psi}^* g, \tag{3.138}$$

$$(\Phi | \hat{a}(g) \Psi) = \text{Tr} \alpha_{\Psi, \Phi} g^*. \tag{3.139}$$

Note that (3.138) and (3.139) are still true if g is an unbounded functional on $\otimes_s^2 \mathcal{W}$ with domain $\otimes_s^{\text{al}^2} \mathcal{D}$, provided that $\Psi, \Phi \in \Gamma_s^{\text{al}}(\mathcal{D})$ where $\mathcal{D} = \text{Dom} h^{-\frac{1}{2}}$, as discussed in Remark 3.4.

The following proposition provides a key estimate for the construction of normally ordered Bogoliubov Hamiltonians:

Proposition 3.22 *Assume that $\|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\| \leq 1$ and $\text{Tr} g^* h^{-1} g < \infty$. Let $\|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\| \leq c$. Then for $\Phi \in \Gamma_s(\mathcal{W})$ with $\|\Phi\| = 1$,*

$$(\Phi | \hat{a}^*(g) \Phi) \leq c (\Phi | d\Gamma(h) \Phi) + \frac{1}{2c} \text{Tr}(g^* h^{-1} g). \tag{3.140}$$

Proof.

$$\begin{aligned}
& (\Phi|\hat{a}^*(g)\Phi) = |\mathrm{Tr}\overline{\alpha_\Phi}g| \\
& = \left| \mathrm{Tr}(\mathbb{1} + \overline{\gamma_\Phi})^{-\frac{1}{2}} \overline{\alpha_\Phi} h^{\frac{1}{2}} h^{-\frac{1}{2}} g (\mathbb{1} + \overline{\gamma_\Phi})^{\frac{1}{2}} \right| \\
& \leq \left(\mathrm{Tr} h^{\frac{1}{2}} \alpha_\Phi (\mathbb{1} + \overline{\gamma_\Phi})^{-1} \alpha_\Phi^* h^{\frac{1}{2}} \right)^{\frac{1}{2}} \\
& \quad \times \left(\mathrm{Tr} h^{-\frac{1}{2}} g (\mathbb{1} + \overline{\gamma_\Phi}) g^* h^{-\frac{1}{2}} \right)^{\frac{1}{2}} \\
& \leq \left(\mathrm{Tr} h^{\frac{1}{2}} \gamma_\Phi h^{\frac{1}{2}} \right)^{\frac{1}{2}} \left(\mathrm{Tr} h^{-\frac{1}{2}} g g^* h^{-\frac{1}{2}} + \|h^{-\frac{1}{2}} g h^{-\frac{1}{2}}\|^2 \mathrm{Tr} h^{\frac{1}{2}} \overline{\gamma_\Phi} h^{\frac{1}{2}} \right)^{\frac{1}{2}} \\
& = \left((\Phi|\mathrm{d}\Gamma(h)\Phi) \right)^{\frac{1}{2}} \left(\mathrm{Tr} g^* h^{-1} g + \|h^{-\frac{1}{2}} g h^{-\frac{1}{2}}\|^2 (\Phi|\mathrm{d}\Gamma(h)\Phi) \right)^{\frac{1}{2}}.
\end{aligned}$$

Then we use the inequality

$$\sqrt{x(y + c_0^2 x)} \leq cx + \frac{y}{2c} \quad (3.141)$$

valid for $x, y \geq 0$, $c > c_0$. \square

Theorem 3.23 *Assume that $\|h^{-\frac{1}{2}} g h^{-\frac{1}{2}}\| < 1$ and $\mathrm{Tr} g^* h^{-1} g < \infty$. Then the quadratic form*

$$\mathrm{d}\Gamma(h) + \frac{1}{2} \hat{a}^*(g) + \frac{1}{2} \hat{a}(g) \quad (3.142)$$

defined on the form domain of $\mathrm{d}\Gamma(h)$ is closed and bounded from below by $-\frac{1}{2} \mathrm{Tr}(g^ h^{-1} g)$. Hence it defines a self-adjoint operator, which we temporarily denote by C . It satisfies*

$$(1 + \mathrm{d}\Gamma(h))^{\frac{1}{2}} (\mathfrak{i} + C)^{-1} (1 + \mathrm{d}\Gamma(h))^{\frac{1}{2}} \text{ is bounded.} \quad (3.143)$$

Proof. By Proposition 3.22,

$$\frac{1}{2} |(\Phi|(\hat{a}^*(g) + \hat{a}(g))\Phi)| \leq c (\Phi|\mathrm{d}\Gamma(h)\Phi) + \frac{1}{2c} \mathrm{Tr} g^* h^{-1} g \|\Phi\|^2. \quad (3.144)$$

Setting $c := \|h^{-\frac{1}{2}} g h^{-\frac{1}{2}}\| < 1$ and using the KLMN Theorem, we see that the form (3.142) is closed and bounded from below, and hence defines a bounded from below self-adjoint operator C . Setting $c = 1$, we see that

$$-\frac{1}{2} \mathrm{Tr}(g^* h^{-1} g) < C. \quad (3.145)$$

(3.143) is also a consequence of the KLMN Theorem. \square

Theorem 3.24 *The operator defined in Thm 3.23 is the normally ordered quantization of B . In other words, following the notation introduced in Subsection 3.7, $C = \hat{H}_B^n$.*

On a formal level the above theorem is essentially obvious. However, there are technical difficulties, for which we will need a few technical lemmas. In these lemmas we use $h \geq 0$ and $\|h^{-\frac{1}{2}} g h^{-\frac{1}{2}}\| < 1$. Note that under this assumption, $\mathfrak{i}\tau$ belongs to the resolvent set of B for $\tau \neq 0$.

Lemma 3.25 *For $\tau \neq 0$, $B(\tau^2 + B^2)^{-1}$ has a dense range.*

Proof. We write

$$B(\tau^2 + B^2)^{-1} = (i\tau + B)^{-1}(-i\tau + B)^{-1}B. \quad (3.146)$$

We will show that (3.146) has a dense range when restricted to $\text{Dom}B$.

First note that $B = AS$ where A is self-adjoint and S is unitary. Hence $\text{Dom}B = S\text{Dom}A$ and $B\text{Dom}B = A\text{Dom}A$. This shows that $B\text{Dom}B$ is dense.

Then we apply twice Lemma A.3 to the bounded operators with dense range $(i\tau + B)^{-1}$ and $(-i\tau + B)^{-1}$. \square

Lemma 3.26 *For $\tau \neq 0$, the operator $A_0^{-\frac{1}{2}}B(\tau^2 + B^2)^{-1}$ is bounded.*

Proof. First note that

$$\|(i\tau S + A_0)^{-\frac{1}{2}}GS(i\tau S + A_0)^{-\frac{1}{2}}\| = \|h^{\frac{1}{2}}g\bar{h}^{\frac{1}{2}}\| < 1. \quad (3.147)$$

Next we check that all the terms on the right of the following identity are bounded:

$$A_0^{-\frac{1}{2}}B(i\tau + B)^{-1} = (1 + A_0^{-\frac{1}{2}}GSA_0^{-\frac{1}{2}}) \quad (3.148)$$

$$\times A_0^{\frac{1}{2}}(i\tau S + A_0)^{-\frac{1}{2}} \quad (3.149)$$

$$\times (1 + (i\tau S + A_0)^{-\frac{1}{2}}GS(i\tau S + A_0)^{-\frac{1}{2}})^{-1} \quad (3.150)$$

$$\times (i\tau S + A_0)^{-\frac{1}{2}}. \quad (3.151)$$

(To see that (3.150) is well defined we use (3.147)). Therefore, $A_0^{-\frac{1}{2}}B(i\tau + B)^{-1}$ is bounded, which obviously implies the boundedness of $A_0^{-\frac{1}{2}}B(\tau^2 + B^2)^{-1}$. \square

Proof of Thm 3.24. Consider $w \in \text{Ran}B(B^2 + 1)^{-1}$. By Lemma 3.25, such w are dense in \mathcal{W} .

Set

$$\begin{bmatrix} w_t \\ \bar{w}_t \end{bmatrix} := e^{itB} \begin{bmatrix} w \\ \bar{w} \end{bmatrix}. \quad (3.152)$$

By Lemma 3.26, $\|h^{-\frac{1}{2}}w_t\|$ is uniformly bounded. Therefore, by Prop. 3.3,

$$(1 + d\Gamma(h))^{-\frac{1}{2}}\hat{\phi}(w_t, \bar{w}_t)(1 + d\Gamma(h))^{-\frac{1}{2}} \quad (3.153)$$

is uniformly bounded. Hence, by (3.143), so is

$$k(t) := (C + i)^{-1}e^{-itC}\hat{\phi}(w_t, \bar{w}_t)e^{itC}(C + i)^{-1}. \quad (3.154)$$

We know that $(w_t, \bar{w}_t) \in \text{Dom}(B)$. But this does not necessarily imply that $w_t \in \text{Dom}h$. It only implies $w_t \in \text{Dom}h^{\frac{1}{2}}$. Therefore, strictly speaking, we cannot write

$$\frac{d}{dt}w_t = ihw_t - ig\bar{w}_t, \quad (3.155)$$

but only

$$\frac{d}{dt}h^{-\frac{1}{2}}w_t = ih^{-\frac{1}{2}}hw_t - ih^{-\frac{1}{2}}g\bar{w}_t. \quad (3.156)$$

However, using the boundedness of $(i + C)^{-1}(i + d\Gamma(h))^{\frac{1}{2}}$ and Prop. 3.3, it is sufficient to compute:

$$(i + C)^{-1}i[C, \hat{\phi}(w_t, \bar{w}_t)](i + C)^{-1} \quad (3.157)$$

$$= (i + C)^{-1}\left(\hat{a}^*(hw_t) + \hat{a}(g\bar{w}_t) - \hat{a}(hw_t) - \hat{a}^*(g\bar{w}_t)\right)(i + C)^{-1} \quad (3.158)$$

$$= (i + C)^{-1}\hat{\phi}\left(\frac{d}{dt}w_t, \frac{d}{dt}\bar{w}_t\right)(i + C)^{-1}. \quad (3.159)$$

Therefore,

$$\frac{d}{dt}k(t) = (C + i)^{-1}e^{-itC} \left(-i[C, (\hat{\phi}(w_t, \bar{w}_t))] \right) \quad (3.160)$$

$$+ \hat{\phi} \left(\frac{d}{dt}w_t, \frac{d}{dt}\bar{w}_t \right) e^{itC} (C + i)^{-1} = 0. \quad (3.161)$$

This shows that $k(t)$ does not depend on t . Therefore,

$$(C + i)^{-1}e^{itC} \hat{\phi}(w, \bar{w}) e^{-itC} (C + i)^{-1} = (C + i)^{-1} \hat{\phi}(w_t, \bar{w}_t) (C + i)^{-1}. \quad (3.162)$$

This proves that e^{itC} implements e^{itB} .

Clearly, $\gamma_\Omega = 0$, $\alpha_\Omega = 0$, and $\Omega \in \text{Dom}(d\Gamma(h)^{\frac{1}{2}}) = \text{Dom}(|C|^{\frac{1}{2}})$. Therefore,

$$(\Omega|C\Omega) = 0. \quad (3.163)$$

Thus, by Proposition 3.13, the operator temporarily denoted C is the normally ordered quantization of B . \square

3.12 Infimum of normally ordered Hamiltonians

In Subsection 2.10, in the finite dimensional context, we defined E_B^n as the infimum of the normally ordered Hamiltonian \hat{H}_B^n . In infinite dimension it is useful to define E_B^n independently of whether \hat{H}_B^n exists or not.

As a basic condition on the symplectic generator B we assume that $h > 0$, $\|h^{-\frac{1}{2}}g\bar{h}^{-\frac{1}{2}}\| < 1$. As in (2.72), for $\sigma \in \mathbb{R}$ we set

$$A_\sigma := A_0 + \sigma GS = \begin{bmatrix} h & \sigma g \\ \sigma \bar{g} & \bar{h} \end{bmatrix}, \quad (3.164)$$

so that $A = A_1$.

Out of the formulas for E^n listed in (2.65)–(2.70) valid in finite dimensions, the most suitable for infinite dimensions seems to be (2.69), which we choose as the definition of E^n :

$$E_B^n := \frac{1}{8} \int_0^1 d\sigma \text{Tr} A_\sigma^{\frac{1}{2}} (A_\sigma^{\frac{1}{2}} S A_\sigma S A_\sigma^{\frac{1}{2}})^{-\frac{1}{2}} A_\sigma^{\frac{1}{2}} G S, \quad (3.165)$$

provided that the above integral is well defined.

(2.70) is another formula for E^n useful in infinite dimension:

Proposition 3.27 *We have*

$$E_B^n := \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1 - \sigma) \text{Tr} \frac{1}{(A_\sigma + i\tau S)} S G \frac{1}{(A_\sigma + i\tau S)} S G. \quad (3.166)$$

More precisely, if (3.166) is well defined as a convergent integral, then it coincides with (3.165).

Below we list a few criteria for the existence of E_B^n .

Theorem 3.28 (1) *Let $\|g\|_1 < \infty$. Then E_B^n is well defined by (3.165).*

(2) *Let $s_- < \frac{1}{2} < s_+$. Suppose that $\text{Tr} g \bar{h}^{-s_-} g^* h^{-s_-} < \infty$ and $\text{Tr} g \bar{h}^{-s_+} g^* h^{-s_+} < \infty$. Then E_B^n is well defined by (3.165) or (3.166).*

(3) *Suppose that $\text{Tr} g h^{-1} g^* < \infty$. Then E_B^n is well defined by (3.165) or (3.166).*

Proof. (1): We apply Inequality (A.7) to the operator $Y := A_\sigma^{\frac{1}{2}}(A_\sigma^{\frac{1}{2}}SA_\sigma SA_\sigma^{\frac{1}{2}})^{-\frac{1}{2}}A_\sigma^{\frac{1}{2}}$ and $X := GS$. Note that Y is uniformly bounded for $\sigma \in [0, 1]$. We obtain

$$|E^n| \leq \frac{1}{8} \int_0^1 d\sigma |\text{Tr} A_\sigma^{\frac{1}{2}}(A_\sigma^{\frac{1}{2}}SA_\sigma SA_\sigma^{\frac{1}{2}})^{-\frac{1}{2}}A_\sigma^{\frac{1}{2}}GS| \quad (3.167)$$

$$\leq \frac{1}{8} \int_0^1 d\sigma \|A_\sigma^{\frac{1}{2}}(A_\sigma^{\frac{1}{2}}SA_\sigma SA_\sigma^{\frac{1}{2}})^{-\frac{1}{2}}A_\sigma^{\frac{1}{2}}\| \text{Tr}\sqrt{G^2} \leq c \text{Tr}\sqrt{G^2}. \quad (3.168)$$

But $\text{Tr}\sqrt{G^2} = 2\text{Tr}\sqrt{g\bar{g}}$. This proves (1).

(2): First note that for $0 \leq s \leq 1$

$$\|A_0^{\frac{s}{2}}(A_0 + i\tau S)^{-\frac{1}{2}}\| \leq \tau^{-\frac{1}{2} + \frac{s}{2}}. \quad (3.169)$$

Indeed,

$$A_0^{\frac{s}{2}}(A_0 + i\tau S)^{-\frac{1}{2}} = A_0^{\frac{s}{2}}(A_0 + i\tau S)^{-\frac{s}{2}} \times (A_0 + i\tau S)^{-\frac{1}{2} + \frac{s}{2}}, \quad (3.170)$$

where the first term is bounded by 1 and the second is bounded by $\tau^{-\frac{1}{2} + \frac{s}{2}}$.

Moreover,

$$\|(A_0 + i\tau S)^{-\frac{1}{2}}GS(A_0 + i\tau S)^{-\frac{1}{2}}\| \leq \|A_0^{-\frac{1}{2}}GSA_0^{-\frac{1}{2}}\| = a < 1. \quad (3.171)$$

Therefore, we can write

$$(A_\sigma + i\tau S)^{-1} = (A_0 + i\tau S)^{-\frac{1}{2}}(\mathbb{1} - \sigma(A_0 + i\tau S)^{-\frac{1}{2}}GS(A_0 + i\tau S)^{-\frac{1}{2}})^{-1}(A_0 + i\tau S)^{-\frac{1}{2}}. \quad (3.172)$$

(3.172) together with (3.169) and (3.171) yield

$$\|A_0^{\frac{s}{2}}(A_\sigma + i\tau S)^{-1}A_0^{\frac{s}{2}}\| \leq (1 - a)^{-1}\tau^{-1+s}. \quad (3.173)$$

Now,

$$\left| \text{Tr} \frac{1}{(A_\sigma + i\tau S)} GS \frac{1}{(A_\sigma + i\tau S)} G \right| \quad (3.174)$$

$$\leq \left\| A_0^{\frac{s}{2}}(A_\sigma + i\tau S)^{-1}A_0^{\frac{s}{2}} \right\|^2 \text{Tr}GA_0^{-s}G^*A_0^{-s}, \quad (3.175)$$

$$\leq (1 - a)^{-2}\tau^{2s-2}\text{Tr}GA_0^{-s}G^*A_0^{-s}, \quad (3.176)$$

where we first used Inequality (A.8) with $Y = Z := A_0^{\frac{s}{2}}(A_\sigma + i\tau S)^{-1}A_0^{\frac{s}{2}}$ and $X := A_0^{-\frac{s}{2}}GA_0^{-\frac{s}{2}}$, and then we applied (3.173). Thus

$$|E^n| \leq \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1 - \sigma) \left| \text{Tr} \frac{1}{(A_\sigma + i\tau S)} GS \frac{1}{(A_\sigma + i\tau S)} G \right| \quad (3.177)$$

$$\leq c \text{Tr}GA_0^{-s+}G^*A_0^{-s+} \int_0^1 \tau^{2s+2}d\tau + c \text{Tr}GA_0^{-s-}G^*A_0^{-s-} \int_1^\infty \tau^{2s-2}d\tau. \quad (3.178)$$

But $\text{Tr}GA_0^{-s\pm}GA_0^{-s\pm} = 2\text{Tr}\bar{g}h^{-s\pm}g\bar{h}^{-s\pm}$. This proves (2).

(3): Applying (A.8) to

$$Y = Z := (\mathbb{1} - \sigma(A_0 + i\tau S)^{-\frac{1}{2}}GS(A_0 + i\tau S)^{-\frac{1}{2}})^{-1}, \quad X := (A_0 + i\tau S)^{-\frac{1}{2}}GS(A_0 + i\tau S)^{-\frac{1}{2}},$$

and using (3.172), (3.171), we obtain

$$\left| \text{Tr} \frac{1}{(A_\sigma + i\tau S)} G S \frac{1}{(A_\sigma + i\tau S)} G S \right| \quad (3.179)$$

$$= \text{Tr} X Y X Y \leq \|Y\|^2 \text{Tr} X X^* \quad (3.180)$$

$$\leq \frac{1}{(1-\sigma a)^2} \text{Tr} \frac{1}{(A_0^2 + \tau^2)^{\frac{1}{2}}} G \frac{1}{(A_0^2 + \tau^2)^{\frac{1}{2}}} G^* \quad (3.181)$$

$$\leq \frac{1}{(1-\sigma a)^2} \text{Tr} G \frac{1}{(A_0^2 + \tau^2)} G^*. \quad (3.182)$$

Therefore,

$$|E^n| \leq \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1-\sigma) \left| \text{Tr} \frac{1}{(A_\sigma + i\tau S)} G S \frac{1}{(A_\sigma + i\tau S)} G S \right| \quad (3.183)$$

$$\leq \frac{1}{8} \int_0^1 d\sigma \frac{(1-\sigma)}{(1-a\sigma)^2} \int \frac{d\tau}{2\pi} \text{Tr} G \frac{1}{(A_0^2 + \tau^2)} G^* \quad (3.184)$$

$$= \frac{(-\log(1-a) - a)}{8a^2} \text{Tr} G \frac{1}{A_0} G^*. \quad (3.185)$$

But $\text{Tr} G \frac{1}{A_0} G = 2\text{Tr} g h^{-1} g^*$. This proves (3) \square

Theorem 3.29 *Suppose that $\text{Tr} g h^{-1} g^* < \infty$, as in Thm 3.28 (3). Let \hat{H}_B^n be defined as in Subsect. 3.11. Let E_B^n be defined as in (3.165). Then*

$$E_B^n = \inf \hat{H}_B^n. \quad (3.186)$$

If \mathcal{W} is finite dimensional, then (3.186) was proven in Thm 2.6. In our proof, we will reduce the full problem to this case. The proof will be divided into several steps.

Step 1. *Suppose that there exists a finite dimensional \mathcal{W}_0 such that $\text{Rang} \subset \mathcal{W}_0$ and h preserves \mathcal{W}_0 . Then (3.186) is true.*

Proof. Set $\mathcal{W}_1 := \mathcal{W}_0^\perp$. Note that $g = g^\#$ implies that $\overline{\mathcal{W}}_1 \subset \text{Kerg}$. Let h_0, g_0 denote the restrictions of g, h to \mathcal{W}_0 . Let h_1 denotes the restriction of h to \mathcal{W}_1 . Consider the symplectic generator on \mathcal{W}_0

$$B_0 := \begin{bmatrix} h_0 & -g_0 \\ \bar{g}_0 & -\bar{h}_0 \end{bmatrix}, \quad (3.187)$$

and the corresponding normally ordered Bogoliubov Hamiltonian

$$\hat{H}_0^n := \hat{H}_{B_0}^n = d\Gamma(h_0) + \frac{1}{2}(\hat{a}^*(g_0) + \hat{a}(g_0)). \quad (3.188)$$

We will write E_0^n , resp. E^n for $E_{B_0}^n$, resp. E_B^n .

We have the decomposition

$$\Gamma_s(\mathcal{W}) \simeq \Gamma_s(\mathcal{W}_0) \otimes \Gamma_s(\mathcal{W}_1) \quad (3.189)$$

The operator \hat{H}^n can be decomposed as

$$\hat{H}^n \simeq \hat{H}_0^n \otimes \mathbb{1} + \mathbb{1} \otimes d\Gamma(h_1). \quad (3.190)$$

We have

$$\inf \hat{H}^n = \inf \hat{H}_0^n = E_0^n = E^n, \quad (3.191)$$

where in the middle step we used the finite dimension of \mathcal{W}_0 . \square

Step 2. Suppose that g is finite dimensional and $\mathbb{1}_{[\delta, \delta-1]}(h)g = g$. Then (3.186) is true.

Proof. Let $\epsilon > 0$. Let us set

$$\pi_{\epsilon, n} := \mathbb{1}_{[(1+\epsilon)^n, (1+\epsilon)^{n+1]}(h), \quad (3.192)$$

$$h_\epsilon := \sum_{n=-\infty}^{\infty} (1+\epsilon)^{n+1} \pi_{\epsilon, n}. \quad (3.193)$$

Note that

$$(1+\epsilon)^{-1} h_\epsilon \leq h \leq h_\epsilon. \quad (3.194)$$

Hence,

$$d\Gamma((1+\epsilon)^{-1} h_\epsilon) \leq d\Gamma(h) \leq d\Gamma(h_\epsilon). \quad (3.195)$$

Now

$$\hat{H}_{\epsilon, -}^n := d\Gamma((1+\epsilon)^{-1} h_\epsilon) + \frac{1}{2}(\hat{a}^*(g) + \hat{a}(g)) \leq \hat{H}^n \leq d\Gamma(h_\epsilon) + \frac{1}{2}(\hat{a}^*(g) + \hat{a}(g)) =: H_{\epsilon, +}^n.$$

Let $\mathcal{W}_{\epsilon, 0}$ be the smallest subspace of \mathcal{W} containing Rang and left invariant by h_ϵ . In other words,

$$\mathcal{W}_{\epsilon, 0} := \text{Span}\{\pi_{\epsilon, n} w : w \in \text{Rang}\}. \quad (3.196)$$

Note that $\pi_{\epsilon, n} \text{Rang} = 0$ for $|n|$ large enough. Therefore, $\mathcal{W}_{\epsilon, 0}$ is finite dimensional.

Thus $\hat{H}_{\epsilon, \pm}^n$ satisfy the conditions of Step 1, and so

$$\inf \hat{H}_{\epsilon, \pm}^n = E_{\epsilon, \pm}^n, \quad (3.197)$$

in the obvious notation. Using Lemma 3.30 we show that

$$\lim_{\epsilon \rightarrow 0} E_{\epsilon, \pm}^n = E^n, \quad (3.198)$$

Besides,

$$\inf \hat{H}_{\epsilon, -}^n \leq \inf H^n \leq \inf \hat{H}_{\epsilon, +}^n. \quad (3.199)$$

\square

Step 3. Suppose that for some $\delta > 0$ we have $\mathbb{1}_{[\delta, \delta-1]}(h)g = g$. Then (3.186) is true.

Proof. We know that $h^{-\frac{1}{2}}g$ is Hilbert-Schmidt. Finite dimensional operators are dense in Hilbert-Schmidt operators. Therefore, given $\epsilon > 0$, we can find a finite dimensional g_ϵ such that $g_\epsilon = g_\epsilon^\#$, $\mathbb{1}_{[\delta, \delta-1]}(h)g = g$ and

$$\|h^{-\frac{1}{2}}g - h^{-\frac{1}{2}}g_\epsilon\|_2 = \sqrt{\text{Tr}(\bar{g} - \bar{g}'_\delta)h^{-1}(g - g'_\delta)} < \epsilon^2. \quad (3.200)$$

Now, the Hilbert-Schmidt norm dominates the operator norm. Hence, (3.200) implies

$$\|h^{-\frac{1}{2}}(g - g_\epsilon)\| \leq \epsilon. \quad (3.201)$$

As a consequence,

$$\|h^{-\frac{1}{2}}(g - g_\epsilon)\bar{h}^{-\frac{1}{2}}\| \leq \epsilon\delta^{-\frac{1}{2}}. \quad (3.202)$$

Therefore, by (3.202),

$$\|h^{-\frac{1}{2}}g_\epsilon\bar{h}^{-\frac{1}{2}}\| \leq \|h^{-\frac{1}{2}}g\bar{h}^{-\frac{1}{2}}\| + \|h^{-\frac{1}{2}}(g-g_\epsilon)\bar{h}^{-\frac{1}{2}}\| \quad (3.203)$$

$$\leq a + \epsilon\delta^{-\frac{1}{2}} =: a_1. \quad (3.204)$$

By choosing ϵ small enough we can guarantee that $a_1 < 1$.

Now

$$\hat{H}^n = (1-\nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\epsilon) + \hat{a}(g_\epsilon)) \quad (3.205)$$

$$+\nu d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g-g_\epsilon) + \hat{a}(g-g_\epsilon)) \quad (3.206)$$

$$\geq (1-\nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\epsilon) + \hat{a}(g_\epsilon)) - \frac{\epsilon^2}{\nu}, \quad (3.207)$$

$$\hat{H}^n = (1+\nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\epsilon) + \hat{a}(g_\epsilon)) \quad (3.208)$$

$$-\nu d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g-g_\epsilon) + \hat{a}(g-g_\epsilon)) \quad (3.209)$$

$$\leq (1+\nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\epsilon) + \hat{a}(g_\epsilon)) + \frac{\epsilon^2}{\nu}. \quad (3.210)$$

The Hamiltonians $(1 \pm \nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\epsilon) + \hat{a}(g_\epsilon))$ satisfy the assumptions of Step 2. \square

Step 4. (3.186) is true without additional assumptions.

Proof. $h^{-\frac{1}{2}}g$ is Hilbert-Schmidt and $s\text{-}\lim_{\delta \rightarrow 0} \mathbb{1}_{[\delta, \delta-1]}(h) = \mathbb{1}$. Hence, for any $\epsilon > 0$ we can find $1 \geq \delta > 0$ such that if we set

$$g_\delta := \mathbb{1}_{[\delta, \delta-1]}(h)g\mathbb{1}_{[\delta, \delta-1]}(\bar{h}), \quad (3.211)$$

then

$$\|h^{-\frac{1}{2}}g - h^{-\frac{1}{2}}g'_\delta\|_2 = \sqrt{\text{Tr}(\bar{g} - \bar{g}_\delta)h^{-1}(g - g_\delta)} < \epsilon. \quad (3.212)$$

Note that

$$\|h^{-\frac{1}{2}}g_\delta\bar{h}^{-\frac{1}{2}}\| \leq \|h^{-\frac{1}{2}}g\bar{h}^{-\frac{1}{2}}\| = a. \quad (3.213)$$

Then we estimate similarly as at the end of Step 3. We argue, that we need only estimates about the Hamiltonians $(1 \pm \nu)d\Gamma(h) + \frac{1}{2}(\hat{a}^*(g_\delta) + \hat{a}(g_\delta))$, which satisfy the assumptions of Step 3. \square

A result about the continuity of E_B^n with respect to h , which we needed in the above proof, is described below.

Lemma 3.30 *Let*

$$B = \begin{bmatrix} h & -g \\ \bar{g} & -\bar{h} \end{bmatrix}, \quad B' = \begin{bmatrix} h' & -g' \\ \bar{g}' & -\bar{h}' \end{bmatrix}. \quad (3.214)$$

with

$$\|h^{-\frac{1}{2}}g\bar{h}^{-\frac{1}{2}}\|, \|h'^{-\frac{1}{2}}g'\bar{h}'^{-\frac{1}{2}}\| \leq a, \quad \text{Tr}gh^{-1}g^*, \text{Tr}g'h'^{-1}g'^* \leq a_1$$

Then

$$\|E_B^n - E_{B'}^n\| \leq c\|h^{-\frac{1}{2}}(h-h')h'^{-\frac{1}{2}}\|, \quad (3.215)$$

where c depends only on a and a_1 .

Proof.

$$\begin{aligned}
& E_B^n - E_{B'}^n \\
&= \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1-\sigma) \operatorname{Tr} \frac{1}{(A_\sigma + i\tau S)} (A'_0 - A_0) \frac{1}{(A'_\sigma + i\tau S)} SG \frac{1}{(A_\sigma + i\tau S)} SG \\
&+ \frac{1}{8} \int_0^1 d\sigma \int \frac{d\tau}{2\pi} (1-\sigma) \operatorname{Tr} \frac{1}{(A'_\sigma + i\tau S)} SG \frac{1}{(A_\sigma + i\tau S)} (A'_0 - A_0) \frac{1}{(A'_\sigma + i\tau S)} SG.
\end{aligned}$$

Then we argue similarly as in the proof of Theorem 3.28 (3). \square

3.13 Weyl Bogoliubov Hamiltonian

Weyl Bogoliubov Hamiltonians play a central role in the theory of Bogoliubov Hamiltonians, providing the simplest algebraic formulas. Unfortunately, in infinite dimensions they are usually ill-defined.

If $A_B \geq 0$, then we can define

$$E_B^w := \frac{1}{4} \operatorname{Tr} \sqrt{A_B^{\frac{1}{2}} S A_B S A_B^{\frac{1}{2}}}, \quad (3.216)$$

which is a nonnegative number, often infinite. Recall that in finite dimension it coincides with the infimum of \hat{H}_B^w .

The following theorem gives (rather restrictive) conditions when we can define the Weyl quantization in any dimension.

Theorem 3.31 *Assume that $h > 0$, $\|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\| =: a < 1$ and $\operatorname{Tr} h < \infty$. Then the following holds:*

- (1) $\operatorname{Tr} g^* h^{-1} g < \infty$.
- (2) $\|g\|_1 < \infty$.
- (3) *By (1), we can define \hat{H}_B^n as in Subsect. 3.11, E_B^n is well defined as in Subsect 3.12, and by Thm 3.29,*

$$\inf H_B^n = E_B^n. \quad (3.217)$$

- (4) *By Thm 3.18 (3), H_B^w is well defined. We have*

$$\hat{H}_B^w := \hat{H}_B^n + \frac{1}{2} \operatorname{Tr} h. \quad (3.218)$$

- (5) $\sqrt{A_B^{\frac{1}{2}} S A_B S A_B^{\frac{1}{2}}}$ *is trace class, so that E_B^w is finite. We have*

$$E_B^w = E_B^n + \frac{1}{2} \operatorname{Tr} h \quad (3.219)$$

$$E_B^w = \inf \hat{H}_B^w. \quad (3.220)$$

Proof. We have

$$h^{-\frac{1}{2}} g g^* h^{-\frac{1}{2}} = h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}} \bar{h} h^{-\frac{1}{2}} g^* h^{-\frac{1}{2}}. \quad (3.221)$$

But $h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}$ and $\bar{h}^{-\frac{1}{2}} g^* h^{-\frac{1}{2}}$ are bounded and \bar{h} is trace class. Therefore, (3.221) is trace class. Hence (1) is true.

$$\|g\|_1 = \|h^{\frac{1}{2}} h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}} \bar{h}^{\frac{1}{2}}\|_1 \leq \|h^{\frac{1}{2}}\|_2 \|h^{-\frac{1}{2}} g \bar{h}^{-\frac{1}{2}}\| \| \bar{h}^{\frac{1}{2}} \|_2 \leq a \|h\|_1 \quad (3.222)$$

proves (2).

We know that h is trace class. Besides, $\|g\|_1 < \infty$ implies $\|g\|_2 < \infty$. Therefore, the assumptions of Thm 3.18 (3) are satisfied. Therefore, \hat{H}_B^w is well defined.

By (3.107), (3.108)

$$\sqrt{A^{\frac{1}{2}}SASA^{\frac{1}{2}}} \leq \frac{(1+a)^{\frac{1}{2}}}{(1-a)^{\frac{1}{2}}}A \leq \frac{(1+a)^{\frac{3}{2}}}{(1-a)^{\frac{1}{2}}}A_0. \quad (3.223)$$

But A_0 is trace class. Hence so is the left hand side of (3.223).

By repeating the arguments of Thm 2.6 we see that (3.219) is true. By Thm 3.18 (3) we have (3.218). Combining (3.217), (3.219) and (3.218), we obtain (3.220). \square

3.14 Infimum of the renormalized Hamiltonian

Recall that in Subsections 2.12 and 2.13 we discussed the renormalized Hamiltonians \hat{H}_B^{ren} and its infimum E_B^{ren} in the context of finite dimension. These objects are of course especially interesting in infinite dimensions.

Note that it may happen that E_B^{ren} is well defined and \hat{H}_B^{ren} is not. In this subsection we discuss only E_B^{ren} , without asking whether \hat{H}_B^{ren} exists.

We will use the framework of Subsection 2.13 with $\lambda = 1$. That means, we assume that $h_0 > 0$, $h = h_0 + h_1$ and $h_1^2 = g\bar{g}$, $h_1g = g\bar{h}_1$. Recall that we have

$$B_0 = \begin{bmatrix} h_0 & 0 \\ 0 & -\bar{h}_0 \end{bmatrix}, \quad (3.224)$$

$$A_0 = B_0S = \begin{bmatrix} h_0 & 0 \\ 0 & \bar{h}_0 \end{bmatrix}, \quad (3.225)$$

$$B_0^2 = A_0^2 = \begin{bmatrix} h_0^2 & 0 \\ 0 & \bar{h}_0^2 \end{bmatrix}, \quad (3.226)$$

$$V = B^2 - B_0^2 = \begin{bmatrix} h_0h_1 + h_1h_0 & -h_0g + g\bar{h}_0 \\ \bar{g}h_0 - \bar{h}_0\bar{g} & \bar{h}_0\bar{h}_1 + \bar{h}_0\bar{h}_1 \end{bmatrix}. \quad (3.227)$$

We use (2.109) to define E_B^{ren} :

$$E^{\text{ren}} := -\frac{1}{4} \int \text{Tr} \frac{1}{B_0^2 + \tau^2} V \frac{1}{B^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^2 \tau^2 \frac{d\tau}{2\pi}, \quad (3.228)$$

provided that the above integral is well defined. Below we give a simple criterion for the well definedness of E^{ren} .

Theorem 3.32 *Suppose that*

$$\|B_0^{-1}VB_0^{-1}\| =: a_1 < 1, \quad (3.229)$$

and

$$L_3 := \frac{1}{4 \cdot 6} \int \text{Tr} \left(V \frac{1}{B_0^2 + \tau^2} \right)^3 \frac{d\tau}{2\pi}, \quad (3.230)$$

$$L_4 := -\frac{1}{4 \cdot 8} \int \text{Tr} \left(V \frac{1}{B_0^2 + \tau^2} \right)^4 \frac{d\tau}{2\pi} \quad (3.231)$$

are finite. (In the case of L_4 the meaning of the assumption is clear, since the integrand is always positive. This does not need to be the case of L_3 —here we assume that the integrand is integrable). Then E^{ren} is well defined.

Proof. Assumption (3.229) is equivalent to

$$a_1 B_0^2 \leq V \leq a_1 B_0^2. \quad (3.232)$$

Therefore,

$$B^2 = B_0^2 + V \geq (1 - a_1) B_0^2. \quad (3.233)$$

Hence, with $c := \frac{1}{1-a_1} > 0$,

$$\frac{1}{(B^2 + \tau^2)} \leq c \frac{1}{(B_0^2 + \tau^2)}. \quad (3.234)$$

Besides,

$$E^{\text{ren}} := \frac{1}{4 \cdot 6} \int \text{Tr} \left(V \frac{1}{B_0^2 + \tau^2} \right)^3 \frac{d\tau}{2\pi} \quad (3.235)$$

$$- \frac{1}{4} \int \text{Tr} \left(\frac{1}{B_0^2 + \tau^2} V \right)^2 \frac{1}{B^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^2 \tau^2 \frac{d\tau}{2\pi}. \quad (3.236)$$

The first term is precisely L_3 . The second term is controlled by L_4 , because

$$0 \leq \frac{1}{4} \int \text{Tr} \left(\frac{1}{B_0^2 + \tau^2} V \right)^2 \frac{1}{B^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^2 \tau^2 \frac{d\tau}{2\pi} \quad (3.237)$$

$$\leq \frac{c}{4} \int \text{Tr} \left(\frac{1}{B_0^2 + \tau^2} V \right)^2 \frac{1}{B_0^2 + \tau^2} \left(V \frac{1}{B_0^2 + \tau^2} \right)^2 \tau^2 \frac{d\tau}{2\pi} \quad (3.238)$$

$$= \frac{c}{4 \cdot 8} \int \text{Tr} \left(V \frac{1}{B_0^2 + \tau^2} \right)^4 \frac{d\tau}{2\pi} = -cL_4. \quad (3.239)$$

To pass from (3.238) to (3.239) we use identity (2.93), which involves integration by parts. \square

A Appendix

A.1 Complex conjugate space

This appendix is a side remark about complex conjugate spaces. This well known but abstract and somewhat confusing concept appears naturally in the context of Bogoliubov Hamiltonians. We follow [DG].

Let \mathcal{W} be a Hilbert space. By definition, a space *complex conjugate to* \mathcal{W} is another Hilbert space $\overline{\mathcal{W}}$ equipped with a fixed anti-unitary map $\chi : \mathcal{W} \rightarrow \overline{\mathcal{W}}$.

In the literature, various authors use several concrete realizations of χ and $\overline{\mathcal{W}}$.

- (1) We can assume that $\mathcal{W} = \overline{\mathcal{W}}$ and χ is antiunitary on \mathcal{W} satisfying $\chi^2 = \mathbb{1}$. Suppose that we choose a basis fixed by χ (which is always possible). Then χ amounts to conjugating the components of vectors in this basis.
- (2) We can also identify $\overline{\mathcal{W}}$ with the space of continuous linear functionals on \mathcal{W} . We then define χ to be the *Riesz isomorphism*, that is,

$$\langle \chi z | w \rangle := (z | w), \quad (\text{A.1})$$

(where $(\cdot | \cdot)$ denotes the scalar product and $\langle \cdot | \cdot \rangle$ the action of a linear functional). If we choose an orthonormal basis in \mathcal{W} and the dual basis in $\overline{\mathcal{W}}$, then again χ amounts to conjugating the components of vectors in this basis.

- (3) Finally, one can set $\overline{\mathcal{W}} = \mathcal{W}$ as the real vector space, changing only the complex structure to the opposite one and the scalar product to the complex conjugate of the original scalar product. χ is defined to be the identity operator. If we fix any basis, then similarly as before, χ is conjugating the components of vectors.

The interpretation (1) is the most naive one. It is often natural—especially if \mathcal{W} is L^2 of some measure space. It is used eg. in Sect. 1 or in [HS]. The interpretation (2) is used in [NNS]. The interpretation (3) is probably the most “orthodox” option—it does not invoke anything besides the vector space structure. In particular, it does not involve the Hilbert space structure of \mathcal{W} .

Note that for all three interpretations, in typical bases, the action of χ is equivalent to conjugating components of vectors. Similarly $\chi p \chi^{-1}$ and $\chi q \chi$ amounts to conjugating matrix elements of p and q .

A.2 *-automorphisms of the algebra of bounded operators

Let \mathcal{H} be a Hilbert space. A bijective linear map α on $B(\mathcal{H})$ is a **-automorphism* if

$$\alpha(BC) = \alpha(B)\alpha(C), \quad \alpha(C^*) = \alpha(C)^*, \quad B, C \in B(\mathcal{H}). \quad (\text{A.2})$$

Proposition A.1 (Example 3.2.14, [BR1]). *The following statements are equivalent:*

- (1) α is a *-automorphism of $B(\mathcal{H})$.
- (2) There exists a unitary $U \in B(\mathcal{H})$ such that

$$\alpha(C) = UCU^*, \quad C \in B(\mathcal{H}), \quad (\text{A.3})$$

If (1),(2) hold, then U is determined uniquely up to a phase factor.

Let $\mathbb{R} \ni t \mapsto \alpha_t$ be a 1-parameter group of *-automorphisms of $B(\mathcal{H})$. We say that it is a C_0^* -group if $t \mapsto \alpha_t(C)$ for any $C \in B(\mathcal{H})$ is weakly continuous.

Proposition A.2 (Example 3.2.35, [BR1]). *The following statements are equivalent:*

- (1) $t \mapsto \alpha_t$ is a C_0^* -group of *-automorphisms of $B(\mathcal{H})$.
- (2) There exists self-adjoint H on \mathcal{H} such that

$$\alpha_t(C) = e^{itH} C e^{-itH}, \quad C \in B(\mathcal{H}). \quad (\text{A.4})$$

If (1), (2) hold, then H is defined uniquely up to an additive real constant.

A.3 Useful identities and inequalities

$$\int \frac{d\tau}{2\pi} \frac{1}{(C^2 + \tau^2)} = \frac{1}{\sqrt{C^2}}. \quad (\text{A.5})$$

$$\int \frac{\tau^2 d\tau}{2\pi} \frac{1}{(C^2 + \tau^2)^2} = \frac{1}{4\sqrt{C^2}}. \quad (\text{A.6})$$

$$|\text{Tr}XY| \leq \|Y\| \text{Tr}\sqrt{XX^*}. \quad (\text{A.7})$$

$$|\text{Tr}XYXZ| \leq \sqrt{\text{Tr}XY^*X^*} \sqrt{\text{Tr}Z^*ZX} \leq \|Y\| \|Z\| \text{Tr}XX^*. \quad (\text{A.8})$$

$$\|a_{11}\|_1 + \|a_{22}\|_1 \leq \left\| \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \right\|_1. \quad (\text{A.9})$$

A.4 Useful lemmas

Lemma A.3 *Let C be a bounded operator on \mathcal{W} with a dense range. Let \mathcal{D} be a dense subspace of \mathcal{W} . Then $C\mathcal{D}$ is dense.*

Proof. Let $w \in \mathcal{W}$. We can find $v_1 \in \mathcal{W}$ such that $\|w - Cv_1\| < \frac{\epsilon}{2}$. We can find $v_2 \in \mathcal{D}$ such that $\|v_1 - v_2\| < \frac{\epsilon}{2\|C\|}$. Now

$$\|w - Cv_2\| < \|w - Cv_1\| + \|Cv_1 - Cv_2\| < \epsilon. \quad (\text{A.10})$$

□

Lemma A.4 *Suppose that h_1, h_2 are self-adjoint operators on \mathcal{W} such that for any $w, w' \in \mathcal{W}$*

$$\lim_{t \rightarrow 0} \frac{1}{t} \left((w|e^{ith_1}w') - (w|e^{ith_2}w') \right) = 0. \quad (\text{A.11})$$

Then $h_1 = h_2$.

Proof. Let $w \in \text{Dom}h_1$ and $w' \in \text{Dom}h_2$. Then (A.11) implies

$$(h_1w|w') = (w|h_2w'). \quad (\text{A.12})$$

This means that $h_1 \subset h_2^* = h_2$ and $h_2 \subset h_1 = h_1^*$. Hence $h_1 = h_2$. □

References

- [BB] V. Bach and J.-B. Bru, Diagonalizing quadratic bosonic operators by non- autonomous flow equation, *Memoirs Amer. Math. Soc.*, (2015).
- [Be] Berezin E. A., *The method of second quantization*, Academic Press, 1966.
- [Bo] N. N. Bogoliubov, On the theory of superfluidity, *J. Phys. (USSR)*, 11 (1947), p. 23.
- [BR1] Brattelli, O., Robinson D. W.: *Operator Algebras and Quantum Statistical Mechanics, Volume 1*, Springer-Verlag, Berlin, second edition 1987.
- [BR2] Brattelli, O., Robinson D. W.: *Operator Algebras and Quantum Statistical Mechanics, Volume 2*, Springer-Verlag, Berlin, second edition 1996.
- [BD] L. Bruneau and J. Dereziński, Bogoliubov hamiltonians and one-parameter groups of Bogoliubov transformations, *J. Math. Phys.*, 48 (2007), p. 022101.
- [DG] Dereziński, J., Gérard, C.: *Mathematics of Quantization and Quantum Fields*, Cambridge Monographs in Mathematical Physics, Cambridge University Press 2013
- [De] Dereziński, J.: Quantum fields with classical perturbations, *Journ. Math. Phys.* 55 (2014) 075201
- [DDMS] Deckert, D.-A., Dürr, D., Merkl, F., Schottenloher, M., 2010: Time Evolution of the External Field Problem in QED *J. Math. Phys.* 51
- [F1] Friedrichs K. O. *Mathematical aspects of the quantum theory of field*, New York, 1953
- [IZ] Itzykson, C., Zuber, J.-B. 1980: *Quantum Field Theory*, McGraw Hill

- [HS] Hainzl, C., Seiringer, R.: The BCS functional of superconductivity and its mathematical properties, *Journal of Mathematical Physics* 57, *Journal of Mathematical Physics* 57(2016) 021101
- [Ho1] L. Hörmander, *The analysis of linear partial differential equations III*, Springer, 1985
- [Ho] L. Hörmander, Symplectic classification of quadratic forms, and general Mehler formulas, *Math. Z.*, 219 (1995), pp. 413–449.
- [NNS] Nam, P.T., Napiórkowski, M., Solovej, J.P.: Diagonalization of bosonic quadratic Hamiltonians by Bogoliubov transformations, *Journal of Functional Analysis* 270 (2016) 4340-4368
- [Ru1] Ruijsenaars S. N. M., On Bogoliubov transformations for systems of relativistic charged particles, *J. Math. Phys.* **18**, 517-526 (1976).
- [Ru2] Ruijsenaars S. N. M., On Bogoliubov transforms II. The general case., *Ann. Phys.* **116**, 105-134 (1978).
- [Sh] D. Shale, Linear symmetries of free boson fields, *Trans. Am. Math. Soc.*, 103 (1962), pp. 149–167.
- [Wi] J. Williamson, On the algebraic problem concerning the normal forms of linear dynamical systems, *Amer. J. of Math.*, 58 (1963), pp. 141–163.