

# HOMOGENEOUS SCHRÖDINGER OPERATORS ON HALFLINE

JAN DEREZIŃSKI

Dep. Math. Meth. in Phys., Faculty of Physics, University of Warsaw

1. **TOY MODEL OF RENORMALIZATION GROUP**
2. **HOMOGENEOUS SCHRÖDINGER OPERATORS**  
(in collaboration with LAURENT BRUNEAU  
and VLADIMIR GEORGESCU)
3. **ALMOST HOMOGENEOUS SCHRÖDINGER OPERATORS**  
(in collaboration with SERGE RICHARD)

## TOY MODEL OF RENORMALIZATION GROUP

Consider the Hilbert space  $\mathcal{H} = L^2(\mathbb{R}_+)$  and operator  $X$

$$Xf(x) := xf(x).$$

Let  $m \in \mathbb{C}$ ,  $\lambda \in \mathbb{C} \cup \{\infty\}$ . We consider a family of operators formally given by

$$H_{m,\lambda} := X + \lambda |x^{\frac{m}{2}}\rangle \langle x^{\frac{m}{2}}|.$$

Let  $U_\tau$  be the **group of dilations** on  $L^2[0, \infty[$ , that is

$$(U_\tau f)(x) = e^{\tau/2} f(e^\tau x).$$

We say that  $B$  is **homogeneous of degree  $\nu$**  if

$$U_\tau B U_\tau^{-1} = e^{\nu\tau} B.$$

E.g.  $X$  is homogeneous of degree 1 and  $|x^{\frac{m}{2}}\rangle\langle x^{\frac{m}{2}}|$  is homogeneous of degree  $1 + m$ .

If  $-1 < \operatorname{Re} m < 0$ , the perturbation  $|x^{\frac{m}{2}}\rangle\langle x^{\frac{m}{2}}|$  is **form bounded** relatively to  $X$  and then  $H_{m,\lambda}$  can be defined.

The perturbation is formally rank one. Therefore,

$$\begin{aligned}
 (z - H_{m,\lambda})^{-1} &= (z - X)^{-1} \\
 &+ \sum_{n=0}^{\infty} (z - X)^{-1} |x^{\frac{m}{2}}\rangle (-\lambda)^{n+1} \langle x^{\frac{m}{2}}| (z - X)^{-1} |x^{\frac{m}{2}}\rangle^n \langle x^{\frac{m}{2}}| (z - X)^{-1} \\
 &= (z - X)^{-1} \\
 &+ \left( \lambda^{-1} - \langle x^{\frac{m}{2}}| (z - X)^{-1} |x^{\frac{m}{2}}\rangle \right)^{-1} (z - X)^{-1} |x^{\frac{m}{2}}\rangle \langle x^{\frac{m}{2}}| (z - X)^{-1}.
 \end{aligned}$$

By straightforward complex analysis methods we obtain

$$\begin{aligned} & \langle x^{\frac{m}{2}} | (z - X)^{-1} | x^{\frac{m}{2}} \rangle \\ &= \int_0^{\infty} x^m (z - x)^{-1} dx = (-z)^m \frac{\pi}{\sin \pi m}. \end{aligned}$$

Therefore, the resolvent of  $H_{m,\lambda}$  can be given in a closed form:

$$\begin{aligned} & (z - H_{m,\lambda})^{-1} = (z - X)^{-1} \\ & + \left( \lambda^{-1} - (-z)^m \frac{\pi}{\sin \pi m} \right)^{-1} (z - X)^{-1} | x^{\frac{m}{2}} \rangle \langle x^{\frac{m}{2}} | (z - X)^{-1}. \end{aligned}$$

The above formula defines a resolvent of a closed operator for all  $-1 < \operatorname{Re} m < 1$  and  $\lambda \in \mathbb{C} \cup \{\infty\}$ . This allows us to define a **holomorphic family of closed operators**  $H_{m,\lambda}$ .

Note that  $H_{m,0} = X$ .

$m = 0$  is special:  $H_{0,\lambda} = X$  for all  $\lambda$ .

We introduce  $H_0^\rho$  for any  $\rho \in \mathbb{C} \cup \{\infty\}$  by

$$(z - H_0^\rho)^{-1} = (z - X)^{-1} - (\rho + \ln(-z))^{-1} (z - X)^{-1} |x^0\rangle \langle x^0| (z - X)^{-1}.$$

In particular,  $H_0^\infty = X$ .

The group of dilations (“the renormalization group”) acts on our operators in a simple way:

$$U_\tau H_{m,\lambda} U_\tau^{-1} = e^\tau H_{m,e^{\tau m} \lambda},$$
$$U_\tau H_0^\rho U_\tau^{-1} = e^\tau H_0^{\rho+\tau}.$$



The essential spectrum of  $H_{m,\lambda}$  and  $H_0^\nu$  is  $[0, \infty[$ .

### **Theorem 1 .**

*1.  $z \in \mathbb{C} \setminus [0, \infty[$  belongs to the point spectrum of  $H_{m,\lambda}$  iff*

$$(-z)^{-m} = \lambda \frac{\pi}{\sin \pi m}.$$

*2.  $H_0^\rho$  possesses an eigenvalue iff  $-\pi < \text{Im } \rho < \pi$ , and then it is  $z = -e^\rho$ .*

For a given  $m, \lambda$  all eigenvalues form a **geometric sequence** that lies on a **logarithmic spiral**, which should be viewed as a curve on the Riemann surface of the logarithm. Only its **“physical sheet”** gives rise to eigenvalues.

For  $m$  which are not purely imaginary, only a finite piece of the spiral is on the “physical sheet” and therefore the number of eigenvalues is finite.

If  $m$  is purely imaginary, this spiral degenerates to a half-line starting at the origin.

If  $m$  is real, the spiral degenerates to a circle. But then the operator has at most one eigenvalue.

**Theorem 2 .** Let  $m = m_r + im_i \in \mathbb{C}^\times$  with  $|m_r| < 1$ .


(i) Let  $m_r = 0$ .

(a) If  $\frac{\ln(|\varsigma|)}{m_i} \in ] - \pi, \pi[$ , then  $\#\sigma_p(H_{m,\lambda}) = \infty$ ,

(a) if  $\frac{\ln(|\lambda \frac{\pi}{\sin \pi m}|)}{m_i} \notin ] - \pi, \pi[$  then  $\#\sigma_p(H_{m,\lambda}) = 0$ .

(ii) If  $m_r \neq 0$  and if  $N \in \mathbb{N}$  satisfies  $N < \frac{m_r^2 + m_i^2}{|m_r|} \leq N + 1$ ,  
then

$$\#\sigma_p(H_{m,\lambda}) \in \{N, N + 1\}.$$



# HOMOGENEOUS SCHRÖDINGER OPERATORS

(in collaboration with LAURENT BRUNEAU  
and VLADIMIR GEORGESCU)

Consider the differential expression

$$L_\alpha = -\partial_x^2 + \left(-\frac{1}{4} + \alpha\right)\frac{1}{x^2}.$$

$L_\alpha$  is is **homogeneous of degree  $-2$** .

How to interpret  $L_\alpha$  as a closed operator on  $L^2[0, \infty[$  homogeneous of degree  $-2$ ?

$L_\alpha$ , and closely related operators  $H_m$ , which we introduce shortly, are interesting for many reasons.

- They appear as the **radial** part of the Laplacian in all dimensions, in the decomposition of the **Aharonov-Bohm Hamiltonian**, in the membranes with **conical singularities**, in the theory of **many body systems with contact interactions** and in the **Efimov effect**.
- They have rather subtle and rich properties illustrating various concepts of the operator theory in Hilbert spaces (eg. the **Friedrichs and Krein extensions**, **holomorphic families of closed operators**).

- Essentially all basic objects related to  $H_m$ , such as their **resolvents, spectral projections, wave and scattering operators**, can be explicitly computed.
- A number of nontrivial identities involving special functions, especially from the **Bessel family**, find an appealing operator-theoretical interpretation in terms of the operators  $H_m$ . Eg. the **Barnes identity** leads to the formula for wave operators.

Two naive interpretations of  $L_\alpha$ :

1. The **minimal** operator  $L_\alpha^{\min}$ : We start from  $L_\alpha$  on  $C_c^\infty]0, \infty[$ , and then we take its closure.
2. The **maximal** operator  $L_\alpha^{\max}$ : We consider the domain consisting of all  $f \in L^2]0, \infty[$  such that  $L_\alpha f \in L^2]0, \infty[$ .



We will see that it is often natural to write  $\alpha = m^2$

### **Theorem 3 .**

- 1. For  $1 \leq \operatorname{Re} m$ ,  $L_{m^2}^{\min} = L_{m^2}^{\max}$ .*
- 2. For  $-1 < \operatorname{Re} m < 1$ ,  $L_{m^2}^{\min} \subsetneq L_{m^2}^{\max}$ , and the codimension of their domains is 2.*
- 3.  $(L_{\alpha}^{\min})^* = L_{\bar{\alpha}}^{\max}$ . Hence, for  $\alpha \in \mathbb{R}$ ,  $L_{\alpha}^{\min}$  is Hermitian.*
- 4.  $L_{\alpha}^{\min}$  and  $L_{\alpha}^{\max}$  are homogeneous of degree  $-2$ .*

Notice that

$$Lx^{\frac{1}{2}\pm m} = 0.$$

Let  $\xi$  be a compactly supported cutoff equal 1 around 0.

Let  $-1 < \operatorname{Re} m$ . Note that  $x^{\frac{1}{2}+m}\xi$  belongs to  $\operatorname{Dom}L_{m^2}^{\max}$ .

This suggests to define the operator  $H_m$  to be the restriction of  $L_{m^2}^{\max}$  to

$$\operatorname{Dom}L_{m^2}^{\min} + \mathbb{C}x^{\frac{1}{2}+m}\xi.$$

## **Theorem 4 .**

1. *For  $1 \leq \operatorname{Re} m$ ,  $L_{m^2}^{\min} = H_m = L_{m^2}^{\max}$ .*
2. *For  $-1 < \operatorname{Re} m < 1$ ,  $L_{m^2}^{\min} \subsetneq H_m \subsetneq L_{m^2}^{\max}$  and the codimension of the domains is 1.*
3.  *$H_m^* = H_{\bar{m}}$ . Hence, for  $m \in ]-1, \infty[$ ,  $H_m$  is self-adjoint.*
4.  *$H_m$  is homogeneous of degree  $-2$ .*
5.  *$\operatorname{sp} H_m = [0, \infty[$ .*
6.  *$\{\operatorname{Re} m > -1\} \ni m \mapsto H_m$  is a holomorphic family of closed operators.*

## Theorem 5 .

1. For  $\alpha \geq 1$ ,  $L_{\alpha}^{\min} = H_{\sqrt{\alpha}}$  is *essentially self-adjoint* on  $C_c^{\infty}[0, \infty[$ .
2. For  $\alpha < 1$ ,  $L_{\alpha}^{\min}$  is not essentially self-adjoint on  $C_c^{\infty}[0, \infty[$ .
3. For  $0 \leq \alpha < 1$ , the operator  $H_{\sqrt{\alpha}}$  is the *Friedrichs extension* and  $H_{-\sqrt{\alpha}}$  is the *Krein extension* of  $L_{\alpha}^{\min}$ .
4.  $H_{\frac{1}{2}}$  is the *Dirichlet Laplacian* and  $H_{-\frac{1}{2}}$  is the *Neumann Laplacian* on halfline.
5. For  $\alpha < 0$ ,  $L_{\alpha}^{\min}$  has no homogeneous selfadjoint extensions.

It is easy to see that

$$\begin{aligned} & x^{-\frac{1}{2}} \left( -\partial_x^2 + \left( -\frac{1}{4} + \alpha \right) \frac{1}{x^2} \pm 1 \right) x^{\frac{1}{2}} \\ &= -\partial_x^2 - \frac{1}{x} \partial_x + \left( -\frac{1}{4} + \alpha \right) \frac{1}{x^2} \pm 1, \end{aligned}$$

which is the **(modified) Bessel operator**.

Therefore, it is not surprising that various objects related to  $H_m$  can be computed with help of functions from the Bessel family.

**Theorem 6 .** *If  $R_m(\lambda; x, y)$  is the integral kernel of the operator  $(\lambda - H_m)^{-1}$ , then for  $\operatorname{Re} k > 0$  we have*

$$R_m(-k^2; x, y) = \begin{cases} \sqrt{xy} I_m(kx) K_m(ky) & \text{if } x < y, \\ \sqrt{xy} I_m(ky) K_m(kx) & \text{if } x > y, \end{cases}$$

where  $I_m$  is the **modified Bessel function** and  $K_m$  is the **MacDonald function**.

**Proposition 7 .** For  $0 < a < b < \infty$ , the integral kernel of  $\mathbb{1}_{[a,b]}(H_m)$  is

$$\mathbb{1}_{[a,b]}(H_m)(x, y) = \int_{\sqrt{a}}^{\sqrt{b}} \sqrt{xy} J_m(kx) J_m(ky) k dk,$$

where  $J_m$  is the **Bessel function**.

Let  $\mathcal{F}_m$  be the operator on  $L^2(0, \infty)$  given by

$$\mathcal{F}_m : f(x) \mapsto \int_0^\infty J_m(kx) \sqrt{kx} f(x) dx$$

$\mathcal{F}_m$  is the so-called **Hankel transformation**.

**Theorem 8.**  $\mathcal{F}_m$  is a bounded invertible involution on  $L^2(0, \infty)$  diagonalizing  $H_m$ , more precisely

$$\mathcal{F}_m H_m \mathcal{F}_m^{-1} = X^2.$$

It satisfies  $\mathcal{F}_m A = -A \mathcal{F}_m$ , where

$$A = \frac{1}{2i} (x \partial_x + \partial_x x)$$

is the self-adjoint **generator of dilations**.



## Theorem 9 Set

$$\mathcal{I}f(x) = x^{-1}f(x^{-1}), \quad \Xi_m(t) = e^{i \ln(2)t} \frac{\Gamma(\frac{m+1+it}{2})}{\Gamma(\frac{m+1-it}{2})}.$$

*Then*

$$\mathcal{F}_m = \Xi_m(A)\mathcal{I}.$$

*Therefore, we have the identity*

$$H_m := \Xi_m^{-1}(A)X^{-2}\Xi_m(A)$$

(Result obtained independently by Bruneau, Georgescu, D, and by Richard and Pankrashkin).

**Theorem 10 .** The *wave operators* associated to the pair  $H_m, H_k$  exist and

$$\begin{aligned}
 \Omega_{m,k}^{\pm} &:= \lim_{t \rightarrow \pm\infty} e^{itH_m} e^{-itH_k} \\
 &= e^{\pm i(m-k)\pi/2} \mathcal{F}_m \mathcal{F}_k \\
 &= e^{\pm i(m-k)\pi/2} \frac{\Xi_k(A)}{\Xi_m(A)}.
 \end{aligned}$$

The formula

$$H_m := \Xi_m^{-1}(A)X^{-2}\Xi_m(A) \quad (1)$$

valid for  $\operatorname{Re} m > -1$ . can be used as an alternative definition of the family  $H_m$  also beyond this domain. It defines a family of closed operators for the parameter  $m$  that belongs to

$$\{m \mid \operatorname{Re} m \neq -1, -2, \dots\} \cup \mathbb{R}. \quad (2)$$

Its spectrum is always equal to  $[0, \infty[$  and it is analytic in the interior of (2).

In fact,  $\Xi_m(A)$  is a unitary operator for all real values of  $m$ . Therefore, for  $m \in \mathbb{R}$ , (1) is well-defined and self-adjoint.

$\Xi_m(A)$  is bounded and invertible also for all  $m$  such that  $\operatorname{Re} m \neq -1, -2, \dots$ . Therefore, the formula (1) defines an operator for all such  $m$ .

One can then pose various questions:

- What happens with this operator along the lines  $\operatorname{Re} m = -1, -2, \dots$ ?
- What is the meaning of the operator to the left of  $\operatorname{Re} = -1$ ? (It is not a differential operator!)

The definition (or actually a number of equivalent definitions) of a **holomorphic family of bounded operators** is quite obvious and does not need to be recalled. In the case of unbounded operators the situation is more subtle.

Suppose that  $\Theta$  is an open subset of  $\mathbb{C}$ ,  $\mathcal{H}$  is a Banach space, and  $\Theta \ni z \mapsto H(z)$  is a function whose values are closed operators on  $\mathcal{H}$ . We say that this is a **holomorphic family of closed operators** if for each  $z_0 \in \Theta$  there exists a neighborhood  $\Theta_0$  of  $z_0$ , a Banach space  $\mathcal{K}$  and a holomorphic family of injective bounded operators  $\Theta_0 \ni z \mapsto B(z) \in B(\mathcal{K}, \mathcal{H})$  such that  $\text{Ran } B(z) = \mathcal{D}(H(z))$  and

$$\Theta_0 \ni z \mapsto H(z)B(z) \in B(\mathcal{K}, \mathcal{H})$$

is a holomorphic family of bounded operators.

We have the following practical criterion:

**Theorem 11** . *Suppose that  $\{H(z)\}_{z \in \Theta}$  is a function whose values are closed operators on  $\mathcal{H}$ . Suppose in addition that for any  $z \in \Theta$  the resolvent set of  $H(z)$  is nonempty. Then  $z \mapsto H(z)$  is a **holomorphic family of closed operators** if and only if for any  $z_0 \in \Theta$  there exists  $\lambda \in \mathbb{C}$  and a neighborhood  $\Theta_0$  of  $z_0$  such that  $\lambda$  belongs to the resolvent set of  $H(z)$  for  $z \in \Theta_0$  and  $z \mapsto (H(z) - \lambda)^{-1} \in B(\mathcal{H})$  is holomorphic on  $\Theta_0$ .*



The above theorem indicates that it is more difficult to study holomorphic families of closed operators that for some values of the complex parameter have an **empty resolvent set**.

**Conjecture 12** . *It is impossible to extend*

$$\{\operatorname{Re} m > -1\} \ni m \mapsto H_m$$

*to a holomorphic family of closed operators on a larger connected open subset of  $\mathbb{C}$ .*

# ALMOST HOMOGENEOUS SCHRÖDINGER OPERATORS (in collaboration with SERGE RICHARD)

For any  $\kappa \in \mathbb{C} \cup \{\infty\}$  let  $H_{m,\kappa}$  be the restriction of  $L_{m^2}^{\max}$  to the domain

$$\text{Dom}(H_{m,\kappa}) = \left\{ f \in \text{Dom}(L_{m^2}^{\max}) \mid \text{for some } c \in \mathbb{C}, \right. \\ \left. f(x) - c(x^{1/2-m} + \kappa x^{1/2+m}) \in \text{Dom}(L_{m^2}^{\min}) \right. \\ \left. \text{around } 0 \right\}, \quad \kappa \neq \infty;$$

$$\text{Dom}(H_{m,\infty}) = \left\{ f \in \text{Dom}(L_{m^2}^{\max}) \mid \text{for some } c \in \mathbb{C}, \right. \\ \left. f(x) - cx^{1/2+m} \in \text{Dom}(L_{m^2}^{\min}) \text{ around } 0 \right\}.$$

For  $\nu \in \mathbb{C} \cup \{\infty\}$ , let  $H_0^\nu$  be the restriction of  $L_0^{\max}$  to

$$\text{Dom}(H_0^\nu) = \left\{ f \in \text{Dom}(L_0^{\max}) \mid \text{for some } c \in \mathbb{C}, \right. \\ \left. f(x) - c(x^{1/2} \ln x + \nu x^{1/2}) \in \text{Dom}(L_0^{\min}) \right. \\ \left. \text{around } 0 \right\}, \quad \nu \neq \infty;$$

$$\text{Dom}(H_0^\infty) = \left\{ f \in \text{Dom}(L_0^{\max}) \mid \text{for some } c \in \mathbb{C}, \right. \\ \left. f(x) - cx^{1/2} \in \text{Dom}(L_0^{\min}) \text{ around } 0 \right\}.$$

## Proposition 13 .

1. For any  $|\operatorname{Re}(m)| < 1$ ,  $\kappa \in \mathbb{C} \cup \{\infty\}$

$$H_{m,\kappa} = H_{-m,\kappa^{-1}}.$$

2.  $H_{0,\kappa}$  does not depend on  $\kappa$ , and these operators coincide with  $H_0^\infty$ .

**Proposition 14 .** *For any  $m$  with  $|\operatorname{Re}(m)| < 1$  and any  $\kappa, \nu \in \mathbb{C} \cup \{\infty\}$ , we have*

$$U_\tau H_{m,\kappa} U_{-\tau} = e^{-2\tau} H_{m, e^{-2\tau m \kappa}},$$
$$U_\tau H_0^\nu U_{-\tau} = e^{-2\tau} H_0^{\nu+\tau}.$$

*In particular, only*

$$H_{m,0} = H_{-m},$$

$$H_{m,\infty} = H_m,$$

$$H_0^\infty = H_0$$

*are homogeneous.*

## Proposition 15 .

$$H_{m,\kappa}^* = H_{\overline{m},\overline{\kappa}} \quad \text{and} \quad H_0^{\nu*} = H_0^{\overline{\nu}}.$$


*In particular,*

- (i)  $H_{m,\kappa}$  is self-adjoint for  $m \in ] - 1, 1[$  and  $\kappa \in \mathbb{R} \cup \{\infty\}$ ,  
and for  $m \in i\mathbb{R}$  and  $|\kappa| = 1$ .*
- (ii)  $H_0^\nu$  is self-adjoint for  $\nu \in \mathbb{R} \cup \{\infty\}$ .*

## Self-adjoint extensions of the Hermitian operator

$$L_\alpha = -\partial_x^2 + \left(-\frac{1}{4} + \alpha\right)\frac{1}{x^2}.$$

K—Krein, F—Friedrichs, dashed line—single bound state, dotted line—infinite sequence of bound states.



Define the unitary operator

$$(If)(x) := x^{-\frac{1}{4}} f(2\sqrt{x}).$$

Its inverse is

$$(I^{-1}f)(x) := \left(\frac{y}{2}\right)^{\frac{1}{2}} f\left(\frac{y^2}{4}\right).$$

Note that

$$I^{-1}XI = \frac{X^2}{4},$$
$$I^{-1}AI = \frac{A}{2}.$$



We change slightly notation: the operators  $H_m$ ,  $H_{m,\kappa}$  and  $H_0^\nu$  will be denoted  $\tilde{H}_m$ ,  $\tilde{H}_{m,\kappa}$  and  $\tilde{H}_0^\nu$

Recall that we introduced the Hankel transformation  $\mathcal{F}_m$ , which is a bounded invertible involution satisfying

$$\begin{aligned}\mathcal{F}_m \tilde{H}_m \mathcal{F}_m^{-1} &= X^2, \\ \mathcal{F}_m A \mathcal{F}_m^{-1} &= -A.\end{aligned}$$

## Theorem 16 .

1.

$$\mathcal{F}_m^{-1} I^{-1} H_{m,\lambda} I \mathcal{F}_m = \frac{1}{4} \tilde{H}_{m,\kappa},$$

where

$$\lambda \frac{\pi}{\sin(\pi m)} = \kappa \frac{\Gamma(m)}{\Gamma(-m)},$$

2.

$$\mathcal{F}_m^{-1} I^{-1} H_0^\rho I \mathcal{F}_m = \frac{1}{4} \tilde{H}_0^\nu,$$

where  $\rho = -2\nu$ .