Non-vacuum twisting type N metrics *

Paweł Nurowski †
Instytut Fizyki Teoretycznej, Uniwersytet Warszawski
ul. Hoża 69, 00-618 Warszawa, Poland

Jerzy F. Plebański ‡
Departamento de Fisica, CINVESTAV
Apdo postal 14-740, 07000 Mexico, DF

July 10, 2000

Abstract

A maximally reduced system of equations corresponding to the twisting type N Einstein metrics is given. When the cosmological constant $\lambda \to 0$ they reduce to the standard equations for the vacuum twisting type N's. All the metrics which are conformally equivalent to the twisting type N metrics and which admit 3-dimensional conformal group of symmetries are presented. In the Feferman class of metrics an example is given of a twisting type N metric which satisfies Bach's equations but is not Einstein.

*Research supported by Komitet Badań Naukowych (Grant nr 2 P03B 060 17)
†`e-mail: nurowski@fuw.edu.pl`
‡`e-mail: Jerzy.Plebanski@fis.cinvestav.mx`

1 Introduction

A way of obtaining the maximally reduced system of equations corresponding to the metrics with the energy momentum tensor of the form $T_{\mu\nu} = k_\mu k_\nu$, where k_μ is a quadruple principal null direction, was presented by one of us (JFP) in an unpublished paper [16]. The method[1] of Ref. [16] can be also applied to the vacuum type N equations

$$R_{\mu\nu} = \lambda g_{\mu\nu} \tag{1}$$

with cosmological constant λ. It turns out that any type N metric satisfying the Einstein equations (1) is generated by a single complex function $L = L(u, z, \bar{z})$ of variables u (real) and z (complex), subjected to the following equations

$$\begin{aligned} \bar{\mathcal{D}}^2\mathcal{D}L - \mathcal{D}^2\bar{\mathcal{D}}\bar{L} &= \frac{\lambda}{3}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)^3 \\ \bar{\mathcal{D}}\partial_u\mathcal{D}L &= \frac{\lambda}{2}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)[\mathcal{D}^2\bar{L} - \bar{\mathcal{D}}\mathcal{D}L]. \end{aligned} \tag{2}$$

Here, $\mathcal{D} = \partial_z - L\partial_u$. In terms of L the metric reads

$$\begin{aligned} g &= 2(\theta^1\theta^2 - \theta^3\theta^4) \\ \theta^3 &= \overline{\theta^3} = \mathrm{d}u + L\mathrm{d}z + \bar{L}\mathrm{d}\bar{z} \\ \theta^1 &= \overline{\theta^2} = [r + \tfrac{1}{2}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)]\mathrm{d}z + \bar{L}_u\theta^3 \end{aligned} \tag{3}$$

$$\begin{aligned} \theta^4 = \overline{\theta^4} = \mathrm{d}r &+ [L_u(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L) - \tfrac{1}{2}\mathcal{D}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)]\mathrm{d}z + [\bar{L}_u(\bar{\mathcal{D}}L - \mathcal{D}\bar{L}) - \tfrac{1}{2}\bar{\mathcal{D}}(\bar{\mathcal{D}}L - \mathcal{D}\bar{L})]\mathrm{d}\bar{z} + \\ &[\tfrac{\lambda}{6}(-r^2 + \tfrac{5}{4}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)^2) - \tfrac{1}{2}\partial_u(\mathcal{D}\bar{L} + \bar{\mathcal{D}}L)]\theta^3, \end{aligned}$$

and the 4-dimensional space-time is parametrized by $(r, u, z, \bar{z})$, r beeing a real coordinate. The only nonvanishing coefficient of the Weyl tensor is

$$\Psi_4 = \frac{\mathcal{D}_u^2\bar{\mathcal{D}}\bar{L} + \frac{\lambda}{6}(3\bar{L}_u - \bar{\mathcal{D}})(\bar{\mathcal{D}}^2L - \mathcal{D}\bar{\mathcal{D}}\bar{L})}{r - \frac{1}{2}(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L)}. \tag{4}$$

The metric (3) has a quadruple principall null direction $k = \partial_r$. It is twisting iff and only iff

$$\mathcal{D}\bar{L} - \bar{\mathcal{D}}L \neq 0. \tag{5}$$

If k is non-twisting all the metrics corresponding to solutions of (2) are known [5]. On the other hand, if condition (5) is satsified no explicit solution to the equations (2) is avaliable. Even the Hauser explicit solution [6] is not easily expressible in terms of function L only.

2 The Feferman class

In this paper we relax the Einstein condition and search for twisting type N metrics, which do not satisfy any additional curvature conditions. One class of such metrics is given by the Feferman conformal class [4], which in the context of GR was first studied by Sparling [17]. To describe the Feferman metrics one needs the notion of a Cauchy-Riemann structure.

[1] The results of Ref. [16] are described in [8] pp. 240-242.

Definition 1 *A Cauchy-Riemann (CR) structure* $(\mathcal{N},[(\Omega,\Omega_1)])$ *is a 3-dimensional manifold* $\mathcal{N}$ *equipped with a class of pairs of 1-forms* $[(\Omega,\Omega_1)]$ *such that* Ω *is real- and* Ω_1 *is complex-valued,* $\Omega\wedge\Omega_1\wedge\bar\Omega_1\neq 0$ *at each point of* $\mathcal{N}$*, two pairs* (Ω,Ω_1) *and* (Ω',Ω_1') *are equivalent iff there existsnonvanishing functions* f *(real) and* h *(complex) and a complex function* p *on* $\mathcal{N}$ *such that*

$$\Omega'=f\Omega\qquad\qquad \Omega_1'=h\Omega_1+p\Omega. \tag{6}$$

A Cauchy-Riemann structure is nondegenerate iff $\mathrm{d}\Omega\wedge\Omega\neq 0$.

Given a nondegenerate CR structure $(\mathcal{N},[(\Omega,\Omega_1)])$ one can always choose a representative (Ω,Ω_1) from the class $[(\Omega,\Omega_1)]$ such that [3, 14]

$$\mathrm{d}\Omega=i\Omega_1\wedge\bar\Omega_1. \tag{7}$$

Since $(\Omega,\Omega_1,\bar\Omega_1)$ constitutes a basis of 1-forms on $\mathcal{N}$ then the differential of Ω_1 uniquely defines functions α,θ (complex) and β (real) such that

$$\mathrm{d}\Omega_1=\bar\alpha\Omega_1\wedge\bar\Omega_1+i\beta\Omega\wedge\Omega_1-\theta\Omega\wedge\bar\Omega_1. \tag{8}$$

Let $(\partial_0,\partial,\bar\partial)$ be a basis of vector fields on $\mathcal{N}$ dual to $(\Omega,\Omega_1,\bar\Omega_1)$, respectively. Then the equation $\mathrm{d}^2\Omega_1\equiv 0$ implies the following identity

$$\partial_0\alpha-i\partial\beta+\bar\partial\bar\theta+i\beta\alpha-\bar\alpha\bar\theta=0. \tag{9}$$

It is convenient to introduce the following operators

$$\triangle=\partial-\alpha\qquad\qquad \delta=\partial_0+i\beta. \tag{10}$$

Condition (6) does not fix the forms (Ω,Ω_1) totally. They still can be transformed by means of the following transformations

$$\Omega\to t\bar t\Omega\qquad\qquad \Omega_1\to t[\Omega_1+i\bar\partial\log(t\bar t)\Omega], \tag{11}$$

where t is a nonvanishing complex function on $\mathcal{N}$. The corresponding transformations of functions α,β,θ are

$$\begin{gathered}\alpha\to\tfrac{1}{t}(\alpha-\partial\log(t\bar t^2))\\ \theta\to\tfrac{1}{\bar t^2}(\theta+\bar\alpha z+\bar\partial z+iz^2)\\ \beta\to\tfrac{1}{|t|^2}[\beta-\bar\partial\partial\log t-\partial\bar\partial\log\bar t-\alpha\bar\partial\log t-\bar\alpha\partial\log\bar t2(\bar\partial\log t\partial+\partial\log\bar t\bar\partial)\log|t|],\end{gathered} \tag{12}$$

where $z=i\bar\partial\log(t\bar t)$.

Let $(\mathcal{N},[(\Omega,\Omega_1)])$ be a CR-structure with (Ω,Ω_1) satisfying (7). One defines a manifold

$$\mathcal{M}=\mathbf{R}\times\mathcal{N} \tag{13}$$

with a canonical projection $\pi:\mathcal{M}\to\mathcal{N}$ and pull-backs the forms (Ω,Ω_1) to $\mathcal{M}$ by means of π. Then, using the same letters to denote the pull-backs, one equipps $\mathcal{M}$ with a class of Lorentzian metrics of the form

$$g=\mathrm{e}^{2\phi}[\Omega_1\bar\Omega_1-\Omega(\mathrm{d}r+W\Omega_1+\bar W\bar\Omega_1+H\Omega)]. \tag{14}$$

Here ϕ,H (real) and W (complex) are arbitrary functions on $\mathcal{N}$ and r is a real coordinate along the factor $\mathbf{R}$ in $\mathcal{M}=\mathbf{R}\times\mathcal{N}$.

It follows that the vector field $k=\partial_r$ is null geodesic and shear-free in any metric (14). It generates a congruence of shear-free and null geodesics in $\mathcal{M}$ which is always twisting due to condition (7). The converse is also true. Any space-time admitting a twisting congruence of shear-free and null geodesics can be obtained in this way [18]. This, in particular, means that any such sace-time defines its corresponding CR-structure - the 3-dimensional manifold of the lines of the congruence.

In the context of the present paper it is interesting to ask when the metrics (14) are of type N with k beeing a quadruple principal null direction. The answer is given by the following theorem [13, 11].

Theorem 1 *The metric (14) has $k=\partial_r$ as a quadruple principal null direction if and only if*

$$\begin{aligned} W &= 2ai\mathrm{e}^{ir}+b \\ H &= \mathrm{e}^{ir}(\bar\triangle - i\bar b)a + \mathrm{e}^{-ir}(\triangle + ib)\bar a + h \\ h &= i(\triangle\bar b - \bar\triangle b) - 6a\bar a - \tfrac{1}{2}(\triangle\bar\alpha + \bar\triangle\alpha + \beta) \end{aligned} \tag{15}$$

where the complex functions $a:\mathcal{N}\to\mathbf{C}$ and $b:\mathcal{N}\to\mathbf{C}$ satisfy

$$2iha - 2\delta a - i\partial\bar\triangle a - \partial(\bar b a) - b\bar\triangle a + ib\bar b a = 0 \tag{16}$$

$$4\bar\alpha\bar\theta - 2\bar\partial\bar\theta + 3i(\delta b - \bar\theta\bar b) + \partial(\bar\triangle b - \triangle\bar b - 4ih) + 8i(a\triangle\bar a - \partial(a\bar a) + iba\bar a) = 0. \tag{17}$$

Take

$$a=0, \qquad b=\frac{2}{3}i\alpha. \tag{18}$$

Then equation (16) is automatically satisfied and equation (17) becomes the identity (9). Then, applying Theorem 1 we have the following Corollary.

Corollary 1 *Let $(\mathcal{N},[(\Omega,\Omega_1)])$ be a CR-structure generated by forms (Ω,Ω_1) satisfying condition (7). Then the metric*

$$g=\mathrm{e}^{2\phi}\{\Omega_1\bar\Omega_1 - \Omega[\mathrm{d}r + \frac{2}{3}i\alpha\Omega_1 - \frac{2}{3}i\bar\alpha\bar\Omega_1 + \frac{1}{6}(\triangle\bar\alpha + \bar\triangle\alpha - 3\beta)\Omega]\} \tag{19}$$

on manifold $\mathcal{M}=\mathbf{R}\times\mathcal{N}$ is of type N with twisting shear-free null geodesics generated by the quadruple principal null dierction $k=\partial_r$.

The metrics (19) are called the Feferman metrics [4]. Their main property is presented in the following theorem.

Theorem 2
Let a pair (Ω,Ω_1) satisfying (7) undergoes transformation (11). Then the metric g of (19) transforms according to[2] $g\to t\bar t g$.

Thus, any nondegenerate CR-structure defines a conformal class of Feferman metrics (19). Each of the metrics in the class is of type N and its quadruple principal null direction defines a congruence of shear-free and null geodesics with twisting rays. We stress here that Corollary 1 provides an effective method of evaluating Feferman twisting type N metrics for each nondegenrate CR-structure. For example, if the CR-structure $\mathcal{N}$ is embeddable in $\mathbf{C}^2$ (cf. [19], p. 499 for definition of embeddability) it may be parametrized by coordinates $(u,z,\bar z)$, u - real, z - complex, and generated by a free complex function $L=L(u,z,\bar z)$ such that $\mathcal{D}\bar L - \bar{\mathcal{D}}L \neq 0$. The 1-forms (Ω,Ω_1) satisfying (7) may be choosen to be

$$\Omega = i\frac{\mathrm{d}u + L\mathrm{d}z + \bar L\mathrm{d}\bar z}{\mathcal{D}\bar L - \bar{\mathcal{D}}L} \tag{20}$$

[2]To get this result one has to redefine the r coordinate according to $r\to r+\frac{i}{3}\log\frac{t}{\bar t}$

$$\Omega_1 = \mathrm{d}z - i\bar{w}\Omega,$$

where

$$w = L_u + \mathcal{D}\log(\mathcal{D}\bar{L} - \bar{\mathcal{D}}L). \tag{21}$$

Then, the Feferman metric (19) is

$$g = \mathrm{e}^{2\phi}\{\mathrm{d}z\mathrm{d}\bar{z} - \Omega[\mathrm{d}r - \frac{i}{3}w\mathrm{d}z + \frac{i}{3}\bar{w}\mathrm{d}\bar{z} - \frac{1}{6}\mathcal{D}\bar{w}\Omega]\} \tag{22}$$

Inserting (2)-(21) into (22) gives an explicit form of the Feferman metric for each embeddable CR-structure.

Another characterization of the Feferman class of metrics is given by the following theorem [17, 10].

Theorem 3 *Feferman metrics g are the only metrics satisfying the following three conditions:*

(a) g are of type N with a quadruple principal null direction generated by a vector filed k,

(b) k is geodesic, shear-free and twisting,

(c) k is a conformal Killing vector field.

A bit disapointing property of the Feferman class is given below [9].

Theorem 4 *None of the Feferman metrics satisfies Einstein equations $R_{\mu\nu} = \lambda g_{\mu\nu}$.*

3 Twisting type N metrics with 3-dimensional group of conformal symmetries

In this section we find a local form of all Lorentzian metrics g which satisfy the following assumptions

(i) g are of type N with k beeing a quadruple principal null direction,

(ii) k is geodesic, shear-free and twisting,

(iii) g is not conformally equivalent to any of the Feferman metrics and not conformally flat,

(iv) g admit at least 3 conformal Killing vector fields.

The following theorem is implicit in Sections 4-6 of Ref. [11].

Theorem 5 *All the metrics satisfying assumptions (i)-(iv) can locally be represnted by*

$$g = \mathrm{e}^{2\phi}[\Omega_1\bar{\Omega}_1 - \Omega(\mathrm{d}r + W\Omega_1 + \bar{W}\bar{\Omega}_1 + H\Omega)], \tag{23}$$

where

$$W = 2ai\mathrm{e}^{ir} + b, \qquad H = -a[\mathrm{e}^{ir}(\bar{\alpha} + i\bar{b}) + \mathrm{e}^{-ir}(\alpha - ib)] + i(\bar{\alpha}b - \alpha\bar{b}) - 6a^2 - \frac{1}{2}\beta + \alpha\bar{\alpha} \tag{24}$$

and the functions $a > 0, b, \alpha, \beta, \theta$ are constants [3]*, satisfying*

$$-12ia^2 - \bar{\alpha}b + ib\bar{b} + 2\alpha(i\bar{\alpha} + \bar{b}) - 3i\beta = 0 \tag{25}$$

[3] If $a \equiv 0$ then the corresponding metric is in the Feferman class.

$$-8ia^2(\alpha-ib)-3b\beta+4\bar{\alpha}\bar{\theta}-3i\bar{b}\bar{\theta}=0. \tag{26}$$

Ω and Ω_1 are related to α,β,θ by (8) and satisfy (7). The space-time is locally a cartesian product $\mathcal{M}=\mathbf{R}\times\mathcal{N}$, with $(\mathcal{N},(\Omega,\Omega_1))$ beeing a three dimensional nondegenerate CR-structure. The coordinate r is choosen so that the orbits of the three conformal symmetries X_i, $i=1,2,3$ are given by r=const and $X_i(r)=0$. The three symmetries X_i are such that

$$\mathcal{L}_{X_i}\Omega=\mathcal{L}_{X_i}\Omega_1=0 \qquad \forall i=1,2,3, \tag{27}$$

so that they constitute also three symmetries of the CR-structure $(\mathcal{N},(\Omega,\Omega_1))$.

It follows from this theorem that all the metrics satisfying (i)-(iv) can be obtained by inspecting the list [3, 15, 11] of all nondegenerate CR-structures admitting three symmetries. Such structures are classified according to the Bianchi type of the corresponding symmetries. For each Bianchi type the forms Ω,Ω_1 and the constants α,β,θ are presented in Ref. [11]. Using this list, one has to check whether a given Bianchi type represented by constants α,β,θ is admitted by the type N equations (25)-(26). If it is, one finds the corresponding a and b.

It turns out that only CR-structures with symmetry groups of Bianchi types VI_h and $VIII$ are admitted by equations (25)-(26). Below we describe the corresponding solutions.

3.1 Solutions for Bianchi type $VIII$

In this case one has a 1-parameter family of nonequivalent CR-structures, parametrized by $k\geq 0$, $k\neq 1$. The manifold $\mathcal{N}$ of such CR-structures can be coordinatized by $(u,z,\bar{z})$, (u-real, z-complex) and the forms Ω,Ω_1 and the constants α,β,θ of Theorem 5 can be choosen so that

$$\Omega=\tfrac{2}{k^2-1}\lambda_0 \qquad \Omega_1=\mu_0-\tfrac{k}{k^2-1}\lambda_0$$

$$\mu_0=\tfrac{2e^{iu}}{z\bar{z}-1}\mathrm{d}z \qquad \lambda_0=\mathrm{d}u+\tfrac{ke^{iu}-i\bar{z}}{z\bar{z}-1}\mathrm{d}z+\tfrac{ke^{-iu}+iz}{z\bar{z}-1}\mathrm{d}\bar{z} \tag{28}$$

$$\alpha=0 \qquad \beta=\tfrac{1}{4}(k^2-2) \qquad \theta=-\tfrac{i}{4}k^2.$$

Inserting the above α,β,θ to the type N equations (25)-(26) one finds two branches of solutions for a and b. The corresponding metrics are

$$g=e^{2\phi}[\Omega_1\bar{\Omega}_1-\Omega\nu] \tag{29}$$

where the forms Ω and Ω_1 are given by (28) and the real 1-form ν is given below for each branch (a) and (b) separately.

(a) If $0\leq k\leq\frac{\sqrt{2}}{2}$ then

$$\nu=\mathrm{d}r+\tfrac{i}{2}\sqrt{3(1-k^2)}(e^{ir}\pm\sqrt{\tfrac{1-2k^2}{1-k^2}})\Omega_1-\tfrac{i}{2}\sqrt{3(1-k^2)}(e^{-ir}\pm\sqrt{\tfrac{1-2k^2}{1-k^2}})\bar{\Omega}_1+$$
$$(\mp\tfrac{3}{8}\sqrt{(1-2k^2)(1-k^2)}(e^{ir}+e^{-ir})+k^2-\tfrac{7}{8})\Omega.$$

Solutions are not conformally flat iff $k\neq\frac{\sqrt{2}}{2}$.

(b) If $k\geq 0$, $k\neq 1$, $k\neq\sqrt{3}$ then

$$\nu=\mathrm{d}r+\tfrac{\sqrt{3}}{2}(ie^{ir}\pm\sqrt{1+k^2})\Omega_1+\tfrac{\sqrt{3}}{2}(-ie^{-ir}\pm\sqrt{1+k^2})\bar{\Omega}_1+$$
$$\tfrac{1}{8}(\mp 3i\sqrt{1+k^2}(e^{ir}-e^{-ir})-k^2-7)\Omega.$$

Solutions are not conformally flat iff $k\neq 1$, $k\neq\sqrt{3}$.

3.2 Solutions for Bianchi type VI_h

In this case one has only one CR-structures for each value of the real parameter $h = -(\frac{1-d}{1+d})^2$, $-1 < d \leq 1$. For each value of d the CR-manifold $\mathcal{N}$ can be coordinatized by real (u, x, y), and the forms Ω, Ω_1 and the constants α, β, θ of Theorem 5 can be choosen so that

$$\Omega = -\tfrac{2}{d+1}\lambda_0 \qquad \Omega_1 = \mu_0 + \tfrac{d}{d+1}\lambda_0$$

$$\mu_0 = y^{-1}\mathrm{d}(x+iy) \qquad \lambda_0 = y^d\mathrm{d}u - y^{-1}\mathrm{d}x$$
$$\alpha = \tfrac{i}{2}(d-1) \qquad \beta = -\tfrac{1}{4}d \qquad \theta = -\tfrac{i}{4}d.$$

For the above α, β, θ the type N equations (25)-(26) imply that the constant b is real and satsifies

$$8b^3 + 8(d-1)b^2 - 2(1+4d+d^2)b - 2d^3 - 3d^2 + 3d + 2 = 0. \tag{30}$$

Note that equation (30) always admits at least one real solution. Once the real solution $b = b(d)$ of this equation is known one has to check, whether the quantity

$$A = \frac{1}{48}[4b^2 + 6(d-1)b + 2d^2 - d + 2]$$

is positive. If it is positive, the constant

$$a = \sqrt{A}. \tag{31}$$

Otherwise there is no solution to (25)-(26) corresponding to $b = b(d)$.
To describe an example of explicit solutions to equations (30)-(31) we choose

$$d = \frac{1}{2}. \tag{32}$$

Then, there are three different solutions for a and b:

$$b_1 = -\frac{1}{4}(\sqrt{6}+1) \qquad a_1 = \frac{1}{8}\sqrt{6 + \frac{5}{3}\sqrt{6}}, \tag{33}$$

$$b_2 = 1 \qquad a_2 = \frac{1}{4}, \tag{34}$$

$$b_3 = \frac{1}{4}(\sqrt{6}-1) \qquad a_3 = \frac{1}{8}\sqrt{6 - \frac{5}{3}\sqrt{6}}. \tag{35}$$

The corresponding metrics are given by

$$g = \mathrm{e}^{2\phi}[\Omega_1\bar{\Omega}_1 - \Omega\nu] \tag{36}$$

where the forms Ω and Ω_1 are given by (30) with $d = \frac{1}{2}$, and the real 1-form ν is given below for each of the three above solutions by

$$\nu_1 = \mathrm{d}r + \tfrac{1}{4}[i\sqrt{6 + \tfrac{5}{6}\sqrt{6}}\mathrm{e}^{ir} - \sqrt{6} - 1]\Omega_1 + \tfrac{1}{4}[-i\sqrt{6 + \tfrac{5}{6}\sqrt{6}}\mathrm{e}^{-ir} - \sqrt{6} - 1]\bar{\Omega}_1 +$$
$$\tfrac{1}{32}[i\sqrt{36 + 10\sqrt{6}}(\mathrm{e}^{ir} - \mathrm{e}^{-ir}) - \sqrt{6} - 10]\Omega$$

$$\nu_2 = \mathrm{d}r + [\tfrac{i}{2}\mathrm{e}^{ir} + 1]\Omega_1 + [-\tfrac{i}{2}\mathrm{e}^{-ir} + 1]\bar{\Omega}_1 +$$
$$[\tfrac{5}{16}i(\mathrm{e}^{-ir} - \mathrm{e}^{ir}) - \tfrac{3}{4}]\Omega$$

$$\nu_3 = \mathrm{d}r + \tfrac{1}{4}[i\sqrt{6-\tfrac{5}{6}\sqrt{6}}\mathrm{e}^{ir} + \sqrt{6} - 1]\Omega_1 + \tfrac{1}{4}[-i\sqrt{6-\tfrac{5}{6}\sqrt{6}}\mathrm{e}^{-ir} + \sqrt{6} - 1]\bar{\Omega}_1 + \\ \tfrac{1}{32}[i\sqrt{36-10\sqrt{6}}(\mathrm{e}^{-ir} - \mathrm{e}^{ir}) + \sqrt{6} - 10]\Omega$$

Each of these metrics is conformally non-flat.

Although equation (30) can be solved explicitely, the formula for b is not very useful in obtaining the explicit forms of the metrics. It is more convenient to solve equation (30) for particular values of d as we did above for $d = \frac{1}{2}$. Instead of giving further examples we present the qualitative description of the solutions, which was obtained by numerical analysis of (30)- (31). We have three possible branches of solutions, corresponding to three different roots $b_1(d), b_2(d), b_3(d)$ of (30). These branches are as follows.

$b_1(d)$ The solutions corresponding to the first root $b_1(d)$ are only possible if $d \geq -0.511878$. It turns out then, that for each value of $d \geq -0.511878$ there exists a type N metric, which is not conformally flat iff $d \neq -\frac{1}{2}$ and $d \neq 0$.

$b_2(d)$ The second root $b_2(d)$ admits solutions for each value of the parameter $d \geq 0$. For each such d there exists precisely one type N metric, which is non conformally flat iff $d \neq 0$.

$b_3(d)$ The third root $b_3(d)$ admits solutions only for $-0.511878 \leq d \leq -0.220789$ or $d \geq 0$. For each such d there exists precisely one type N metric which is conformally non-flat iff $d \neq -0.333347$, $d \neq -0.220789$, $d \neq 0$, $d \neq 1$.

We close this section with a remark that the metrics satisfying assumptions (i)-(iv) are not conformally equivalent to the Ricci flat metrics. This result follows directly from the analysis performed in Ref. [11].

4 Examples of twisting type N metrics admitting two conformal symmetries.

In this section we present examples of type N metrics admitting two conformal symmetries. We additionally assume that the metrics are not conformally flat and that they do not belong to the Feferman class. The general solution for such a problem is rather hopeless to obtain but the following two examples can be given.

Case A.

Consider a 3-dimensional manifold parametrized by the real coordinates (u, x, y). The CR-structure $(\mathcal{N}, [(\Omega, \Omega_1)])$ is generated on $\mathcal{N}$ by the forms

$$\Omega = \mathrm{d}u - y\mathrm{d}x \qquad \Omega_1 = \frac{\sqrt{2}}{2}(\mathrm{d}x + i\mathrm{d}y). \tag{37}$$

The forms (37) satisfy (7). On $\mathcal{M} = \mathbf{R} \times \mathcal{N}$ introduce a coordinate r along the $\mathbf{R}$ factor and consider the metric

$$g = \mathrm{e}^{2\phi}[\Omega_1\bar{\Omega}_1 - \Omega\nu] \tag{38}$$

with the 1-form ν defined by

$$\nu = \mathrm{d}r + y^{-1}[(\tfrac{\sqrt{2}}{2} - \sqrt{3} + (\tfrac{3}{2})^{\frac{1}{4}}i\mathrm{e}^{ir})\Omega_1 + (\tfrac{\sqrt{2}}{2} - \sqrt{3} - (\tfrac{3}{2})^{\frac{1}{4}}i\mathrm{e}^{-ir})\bar{\Omega}_1] + \\ y^{-2}[\tfrac{i6^{\frac{1}{4}}}{4}(\sqrt{6} - 2)(\mathrm{e}^{ir} - \mathrm{e}^{-ir}) + \tfrac{\sqrt{6}}{4} - 1]\Omega \tag{39}$$

It is a matter of straigthforward calculation to see that the so defined metric is of type N, admits a congruence of twisting shear-free and null geodesics aligned with the principal null direction and is never conformally flat. Moreover it has only two conformal symmetries $X_1 = \partial_u$, $X_2 = \partial_x$.

Case B

Now the CR-structure $(\mathcal{N}, [(\Omega, \Omega_1)])$ is generated by the forms

$$\Omega = \mathrm{d}u - y^2\mathrm{d}x \qquad \Omega_1 = \sqrt{y}(\mathrm{d}x + i\mathrm{d}y). \tag{40}$$

The forms (40) satisfy (7). The type N metric is defined on $\mathcal{M} = \mathbf{R} \times \mathcal{N}$ by

$$g = \mathrm{e}^{2\phi}[\Omega_1\bar{\Omega}_1 - \Omega\nu] \tag{41}$$

with the 1-form ν defined by

$$\begin{aligned}\nu = \mathrm{d}r + \tfrac{1}{2y^{3/2}}[(i\sqrt{5+2\sqrt{19}}\mathrm{e}^{ir} - \sqrt{19})\Omega_1 + (-i\sqrt{5+2\sqrt{19}}\mathrm{e}^{-ir} - \sqrt{19})\bar{\Omega}_1] + \\ \tfrac{i}{8y^3}[\sqrt{5+2\sqrt{19}}(\sqrt{19}-1)(\mathrm{e}^{ir} - \mathrm{e}^{-ir}) - 16 - 2\sqrt{19}]\Omega.\end{aligned} \tag{42}$$

Here r is a coordinate r along the factor $\mathbf{R}$ in $\mathcal{M}$.

The above metric is of type N, admits a congruence of twisting shear-free and null geodesics aligned with the principal null direction and is never conformally flat. It has only two conformal symmetries $X_1 = \partial_u$, $X_2 = \partial_x$.

5 Example of a type N metric with vanishing Bach tensor and not conformal to an Einstein metric

It is interesting to ask whether metrics (38)-(39), (41)-(42) are conformally equivalent to Einstein metrics. It is known that a neccessary condition for a metric to be conformal to an Einstein metric is that its Bach tensor

$$B_{\mu\nu} = C_{\mu\rho\nu\sigma}{}^{;\sigma\rho} + \frac{1}{2}C_{\mu\rho\nu\sigma}R^{\rho\sigma} \tag{43}$$

identically vanishes [2, 7]. (We denoted the Weyl conformal curvature by $C_{\mu\rho\nu\sigma}$.)

With a help of an extremely powerful symbolic algebra package GRTensor [12] we calculated the Bach tensor for metrics (38)-(39) and (41)-(42). In both cases it is never vanishing, so we conclude that the metrics (38)-(39) and (41)-(42) are not conformally equivalent to any Einstein metric.

Surprisingly within the Feferman class we found a metric which has vanishing Bach tensor. This metric reads as follows:

$$g = \mathrm{e}^{2\phi}[\mathrm{d}x^2 + \mathrm{d}y^2 - \frac{2}{3}(\mathrm{d}x + y^3\mathrm{d}u)(y\mathrm{d}r + \frac{11}{9}\mathrm{d}x - \frac{1}{9}y^3\mathrm{d}u)]. \tag{44}$$

This metric, like any other from the Feferman class, is of type N, admits a congruence of twisting shear-free and null geodesics aligned with the principal null direction and is not conformally flat. It is interesting that it satisfies the Bach equations and, beeing in the Feferman class, is not conformal to any Einstein metrics. To the best of our knowledge this is the only known example of a Lorentzain metric having this last property [1].

References

[1] Bailey T N, Eastwood M G (1995) "Self-dual manifolds need not be locally conformal to Einstein" in *Further advances in twistor theory. Pitman Research Notes in Mathematics Series* **232**, Hughston L P, Kobak P Z, Mason L J (eds), 132-134

[2] Baston R J, Mason L J (1995) "The conformal Einstein equations" in *Further advances in twistor theory. Pitman Research Notes in Mathematics Series* **232**, Hughston L P, Kobak P Z, Mason L J (eds), 131-132

[3] Cartan E (1932) "Sur la geometrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes" Part I: *Ann. Math. Pura Appl.* **11**, 17-90, Part II: *Ann. Scuola Norm. Sup. Pisa* **1**, 333-354

[4] Feferman C L (1976) "Monge-Ampere equations, the Bergman kernel, and geometry of pseudoconvex domains" *Ann. of Math* **103**, 395-416, and correction ibid. **104**, 393-394

[5] Garcia Diaz A, Plebański J F (1981) "All nontwisting N's with cosmological constant" *Journ. Math. Phys.* **22**, 2655-2658

[6] Hauser I (1974) "Type N gravitational fields with twist" *Phys. Rev. Lett.* **33** 1112-1113

[7] Kozameh C N, Newman E T, Tod K P (1985) "Conformal Einstein spaces" *Gen. Rel. Grav.* **17**, 343-352

[8] Kramer D, Stephani H, MacCallum M, Herlt E (1980) *Exact solutions of Einstein's field equations* Deutscher, Berlin

[9] Lewandowski J (1988) "On the Feferman class of metrics associated with 3-dimensional CR space" *Lett. Math. Phys.* **15**, 129-135

[10] Lewandowski J (1989) *Gravitational fields and CR geometry*, PhD Thesis (in Polish), Department of Physics, Warsaw University

[11] Lewandowski J, Nurowski P (1990) "Algebraically special twisting gravitational fields and 3-dimensional CR structures", *Class. Q. Grav.* **7**, 309-328

[12] Musgrave P, Pollney D, Lake K, (1996-1998) "GRTensor. Symbolic computation package for doing GR on computers" http://grtensor.phy.queensu.ca/

[13] Nurowski P (1991) *Gravitational and electromagnetic fields associated with shear-free congruences of null geodesics and CR structures*, MSc Thesis, SISSA

[14] Nurowski P (1993) *Einstein equations and Cauchy-Riemann geometry*, PhD Thesis, SISSA

[15] Nurowski P, Tafel J (1988) "Symmetries of CR spaces", *Lett. Math. Phys.* **15**, 31-38 (1988)

[16] Plebański J F (1978) "Solutions of Einstein equations $G_{\mu\nu} = -\rho k_\mu k_\nu$ determined by the condition that k_μ is a quadruple Debever-Penrose vector" (unpublished)

[17] Sparling G A J (1985) "Twistor theory and characterization of Feferman's conformal structures", (unpublished)

[18] Trautman A (1984) "Deformations of the Hodge map and optical geometry" *Journ. Geom. Phys.* **1** 85-95

[19] Trautman A (1998) "On complex structures in physics" in *On Einstein's path. Essays in honor of Engelbert Schuking*, Harvey A, ed., Springer, 487-501