

# Phenomenology of Heterotic Orbifolds

Saúl Ramos-Sánchez


String Pheno '09

June 16th, 2009

Based on collaborations with:

K.S. Choi, O. Lebedev, H.P. Nilles, S. Raby, M. Ratz, P. Vaudrevange & A. Wingerter

hep-th/0611095, arXiv:0812.2120, arXiv:0902.3070



Dixon, Harvey, Vafa, Witten (1985-86)  
Ibáñez, Nilles, Quevedo (1987)  
Font, Ibáñez, Quevedo, Sierra (1990)  
Katsuki, Kawamura, Kobayashi, Ohtsubo, Ono, Tanioka (1990)  
Kobayashi, Raby, Zhang (2004)  
Förste, Nilles, Vaudrevange, Wingerter (2004)  
Buchmüller, Hamaguchi, Lebedev, Ratz (2004-06)  
Kobayashi, Nilles, Plöger, Raby, Ratz (2006)  
Faraggi, Förste, Timirgaziu (2006)  
Förste, Kobayashi, Ohki, Takahashi (2006)  
Kim, Kye (2006-07)  
Choi, Kim (2006-08)

...


# What we know so far...

## From the heterotic Mini-Landscape

cf. talk by Peter Nilles

*Factorizable* orbifolds:

$$\mathcal{O} = T^2 \times T^2 \times T^2 / \mathbb{Z}_6 - \Pi$$



$$E_8 \times E_8 \xrightarrow{\mathcal{O}} \mathcal{G}_{4D} = \mathcal{G}_{SM} \times \mathcal{G}_{\text{hidden}}$$

Lebedev, Nilles, Raby, R-S, Ratz, Vaudrevange, Wingerter (2006-08)

## From the heterotic Mini-Landscape

cf. talks by Rolf Kappl, Peter Nilles

- $\sim 300$  MSSM-like orbifolds

- $\mathcal{G}_{4D} = \mathcal{G}_{SM} \times \mathcal{G}_{\text{hidden}}$
- $\mathcal{N} = 1$  SUSY vacua
- 3 SM generations + Higgses + **no** exotics
- see-saw neutrino masses
- gauge coupling unification
- $y_t \sim g$  @  $M_{GUT}$  Hosteins, Kappl, Ratz, Schmidt-Hoberg (2009)
  - & suppression  $y_t < g$  @ low energy

- Most models

- intermediate scale SUSY favored 😊
- endowed with local GUTs 😊

Nilles, R-S, Ratz, Vaudrevange (2008)

## Ingredients:

- *only one pair* of *untwisted* Higgs doublets
- $\mathcal{W}$  preserves an *accidental*  $U(1)_R$  @ order  $N$
- vacuum such that  $-F_i^\dagger = \frac{\partial \mathcal{W}}{\partial s_i} = 0$  (for all singlets  $s_i$ )

## Consequences:

- $H_u H_d \supset \mathbf{1}$  (gauge & string symm.)  
 $\Rightarrow \mu \sim \mathcal{W}(s_i)$
- $\langle \mathcal{W} \rangle = 0$  @ order  $N$   
 $\Rightarrow \mu = 0$  &  $D\mathcal{W} = 0$
- $U(1)_R$  *explicitly* broken @ order  $M > N$   
 $\Rightarrow \langle \mathcal{W} \rangle \sim \mu \sim \langle s_i \rangle^M \ll 1, \quad \langle s_i \rangle \sim 0.1$

If there were CP-violation in QCD

⇒ electron dipole moment of neutron... **unobserved!**

**BUT**  $-\theta F^{\mu\nu} \tilde{F}_{\mu\nu} \dots$  **allowed!**

Implying  $\theta \leq 10^{-11}$  ... **why?**

Baker *et al.* (2006)

An elegant solution:  $U(1)_{PQ}$

- (QCD-) anomalous & global symmetry
- $U(1)_{PQ}$  spontaneously broken
- $\theta = \langle a(x) \rangle \sim 0$ , axion:  $a(x)$
- axion decay constant  $F_a$   
$$10^9 \text{ GeV} \leq F_a \leq 10^{12} \text{ GeV}$$
- highly suppressed  $m_a \sim 0$

Acci😊ns

(no, it isn't a typo)

Choi, Nilles, R-S, Vaudrevange (2009)

In string theory:

- No **exact** global  $U(1)$
- If  $U(1)$  anomalous, then **gauged**
- Natural scale close to  $M_{\text{Pl}}$

$\Rightarrow$  QCD-axions are elusive in string theory

Svrček, Witten (2006)



In string theory:

- No **exact** global  $U(1)$
- If  $U(1)$  anomalous, then **gauged**
- Natural scale close to  $M_{\text{Pl}}$

⇒ QCD-axions are elusive in string theory


Svrček, Witten (2006)

Addressing the strong CP problem amounts to

In heterotic orbifolds:

- 1 finding  $U(1)_{\text{PQ}}$
- 2 getting  $F_a < M_{\text{Pl}}$
- 3 guaranteeing large suppression of  $m_a$

stringy discrete symmetries  $\rightarrow U(1)$  accidental symmetries of  $\mathcal{W}$



- compute  $\mathcal{W}$  @ order  $N$
- identify accidental  $U(1)$ s of  $\mathcal{W}$
- verify whether they have  $SU(3)$ - $SU(3)$ - $U(1)_{PQ}$  anomalies
- identify SM singlets charged under  $U(1)_{PQ}$ ,  
$$s_i = v_i e^{ia_i/v_i} \quad a_i: \text{accions}$$
- consistent vacuum (preserving e.g. SUSY)

Choi, Kim, Kim (2006)

## Accions: ② Lowering $F_a$

A single  $U(1)_{PQ}$ :

- $\langle \varphi \rangle = v \neq 0 \quad \Rightarrow \quad F_a \sim q_{PQ} v / \mathcal{A}$
- $\langle \varphi_i \rangle = v_i \neq 0 \quad \Rightarrow \quad F_a \sim \sum q_{PQ}^i v_i / \mathcal{A}$

At least one  $v_i \sim M_{Pl}$  😞

## Accions: ② Lowering $F_a$

A single  $U(1)_{PQ}$ :

- $\langle \varphi \rangle = v \neq 0 \quad \Rightarrow F_a \sim q_{PQ} v / \mathcal{A}$
- $\langle \varphi_i \rangle = v_i \neq 0 \quad \Rightarrow F_a \sim \sum q_{PQ}^i v_i / \mathcal{A}$

At least one  $v_i \sim M_{Pl}$  ☹️

Alternative:  $U(1)_P \times U(1)_Q$

- $\langle \varphi_{1,2} \rangle = v_{1,2} \neq 0$   
 $\Rightarrow F_a = \frac{v_1 v_2 (q_P^1 q_Q^2 - q_Q^1 q_P^2)}{\sqrt{(q_f^1 v_1)^2 + (q_f^2 v_2)^2}} \approx v_2 \frac{q_P^1 q_Q^2 - q_Q^1 q_P^2}{q_f^1} \quad (*)$

At least one  $v_i \sim M_{Pl} \quad v_1 > v_2 \quad \text{☺️}$

For  $\langle \varphi_i \rangle = v_i \neq 0$  &  $v_1 > v_2 > v_3 > \dots$ ,  $(*)$  still holds!! ☺️

## Accions: ② Lowering $F_a$

A single  $U(1)_{PQ}$ :

- $\langle \varphi \rangle = v \neq 0 \quad \Rightarrow \quad F_a \sim q_{PQ} v / \mathcal{A}$
- $\langle \varphi_i \rangle = v_i \neq 0 \quad \Rightarrow \quad F_a \sim \sum q_{PQ}^i v_i / \mathcal{A}$

At least one  $v_i \sim M_{Pl}$  ☹️

In general:  $U(1)_{PQ}^n$

$$F_a \propto v_n \times \text{charges}$$

At least one  $v_i \sim M_{Pl} \quad v_1 > v_2 > \dots > v_n > \dots$

intermediate scale  $v_n$  possible! 😊

## Accions: ③ Verifying small $m_a$ ?

No exact global U(1) in string theory

$\Rightarrow$  U(1)<sub>PQ</sub> broken *explicitly* @ order  $M > N$

The explicit breaking induces an axion mass  $m_a$

Barr, Seckel (1992)

Neutrino-dipole-moment constraint:  $m_a \lesssim 3 \times 10^{-7} \text{ GeV}^2 / F_a$

Kim, Carosi (2008)

Sufficient condition for suppressed  $m_a$ :  $M \geq 12$

Kamionkowski, March-Russell (1992)

Also achieved if first **mass contribution** appears at order

$$\mathcal{O} \geq 12, \quad \mathcal{O} \geq M > N$$

Choi, Nilles, R-S, Vaudrevange (2009)

# Example

Choi, Nilles, R-S., Vaudrevange (2009)

# Minilandscape Example with QCD *Accions*

- $\mathcal{G}_{4D} = \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y \times [\text{SU}(4) \times \text{SU}(2)]$

3	$(\mathbf{3}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{1/6}$	$q_i$					
9	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{-1/2}$	$\ell_i$	6	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{1/2}$	$\bar{\ell}_i$	4	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_0$ $m_i$
3	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_1$	$\bar{e}_i$				2	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{2})_0$ $n_i$
3	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{-2/3}$	$\bar{u}_i$				65	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_0$ $s_i^0$
8	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{1/3}$	$\bar{d}_i$	5	$(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{-1/3}$	$d_i$	12	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_0$ $h_i$
12	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{1/2}$	$s_i^+$	12	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{-1/2}$	$s_i^-$	10	$(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_0$ $w_i$
2	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{1/2}$	$x_i^+$	2	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{-1/2}$	$x_i^-$	10	$(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_0$ $\bar{w}_i$

- 1 accidental  $\text{U}(1)_{\text{PQ}}$  + gauge anomalous  $\text{U}(1)_A$
- Break  $\text{U}(1)_A$ , preserve  $\mathcal{N} = 1$  SUSY & unbroken  $\text{U}(1)_{\text{PQ}}$

$$v_1 : \langle s_{19}^0 \rangle \sim \langle s_{49}^0 \rangle \sim \langle s_{52}^0 \rangle \sim \langle s_{57}^0 \rangle \sim \langle s_{59}^0 \rangle \sim M_{\text{Pl}}$$

- Vacuum that breaks  $\text{U}(1)_{\text{PQ}}$  at lower energies:

$$v_2 : \langle s_1^0 \rangle \sim \langle s_{36}^0 \rangle \sim \langle s_{37}^0 \rangle \sim 10^{12} \text{ GeV}$$

In this model

$$F_a \propto v_2 \Rightarrow m_a \sim 10^{-9} \text{ GeV}^2 / F_a \approx 10^{-21} \text{ GeV} \quad \text{☺}$$



In promising  $\mathbb{Z}_6$ -II heterotic orbifolds:

QCD axions

- *accidental* U(1)s with QCD-anomalies ✓
- spontaneous breaking  $\Rightarrow$  axions ✓
- two or more U(1)<sub>PQ</sub> broken ✓
  - $\Rightarrow F_a$  fits bounds ☺
- explicit breaking of U(1)<sub>PQ</sub> @ high orders ✓
  - $\Rightarrow$  suppressed  $m_a \sim 0$  ☺