

Advanced Graduate Quantum Mechanics

somer term 2021-22

Krzysztof Byczuk

Institute of Theoretical Physics, Faculty of Physics,
University of Warsaw
byczuk@fuw.edu.pl
www.fuw.edu.pl/ byczuk
17-06-2022

Rules

- Lectures are on Tuesdays at 14:15-16:00 in B2.38Biof room.
- Tutorials are on Wednesday at 11:15-13:00 in 2.25 room.
- Home problems will be offered but not be checked. Some of these problems or similar ones might occur during a colloquium or an exam.
- Standard way of passing the course
 - Colloquium, 2/5 pts.
 - Written exam, 3/5 pts.
 - Oral exam (in uncertain cases)
- Resit exam to pass the course
 - Written exam, 100% pt.
 - Oral exam (in uncertain cases)

Final grade is based on total score points normalized to 100 and determined as follows:

5+ for 99-100 pt.
5 for 90-98 pt.
4+ for 81-89 pt.
4 for 72-80 pt.
3+ for 62-71 pt.
3 for 50-61 pt.
2 for 0-49 pt.

Warning: points from the first colloquium and exam and from the second exams do not sum up.

Dates of exams:

colloquium, 26/04/22, 14:15-16:00, B2.38Biof room

written exam I, 20/06/22, 9:15-13:00, B2.38Biof room

oral exam I, on e-mail note

written exam II, in September 2022

oral exam II, on e-mail note

1 Week I, 28/02-06/03/2022

1.1 Lecture

1.2 Tutorial

1. *Conservation of momentum in classical physics* - Consider a single particle moving in a homogeneous space. Within the Lagrangian formalism show that the momentum of the particle is conserved in time.
2. *Conservation of energy in classical physics* - Consider a single particle moving in space in a time independent potential. Within the Lagrangian formalism show that the energy of the particle is conserved in time.
3. *Conservation of angular momentum in classical physics* - Consider a particle moving in an isotropic space. Within the Lagrangian formalism show that the angular momentum of the particle is conserved in time.
4. *Conserved quantity for a charge classical particle in a homogeneous electric field* - Derive a conservation law and find a conserved quantity for a classical particle with charge q and mass m moving in a homogeneous electric field with an intensity \mathbf{E} .

1.3 Homework problems

1. *Conserved quantity for a charge classical particle in a homogeneous magnetic field* - Derive a conservation law and find a conserved quantity for a classical particle with charge q and mass m moving in a homogeneous magnetic field with an induction \mathbf{B} .
2. *Angular momenta in different reference frames* -
 - (a) What is the connection between the angular momenta in two reference systems which are at rest relative to each other and whose origins are separated by the distance vector \mathbf{a} ?
 - (b) What is the relation between the angular momenta in two inertial reference systems which move with velocity \mathbf{V} relative to each other?
3. *Runge-Lenz-Laplace vector in the Kepler-Coulomb problem* - Consider a single particle moving in a central potential $V(\mathbf{r}) = -\alpha r/r^3$. Introduce a vector $\mathbf{J} = \mathbf{p} \times \mathbf{L} - \alpha \mathbf{r}/r$, where \mathbf{p} and \mathbf{L} are momentum and angular momentum, respectively. Check that $\mathbf{J} \cdot \mathbf{L} = 0$ and $J^2 = 2EL^2 + \beta^2$, where E is the energy (Hamiltonian) and $\beta = \alpha/m$, with m as a mass of the particle. Prove that

$$\frac{d}{dt} \left(\dot{\mathbf{r}} \times \mathbf{L} - \alpha \frac{\mathbf{r}}{r} \right) = 0,$$

so \mathbf{J} is invariant in time. How many components of \mathbf{J} are in fact independent? Conclude why \mathbf{J} and \mathbf{r} are in the plane perpendicular to \mathbf{L} and how the motion of a particle is constrained. In the polar coordinate system parametrize \mathbf{J} and \mathbf{r} and write

$\mathbf{J} \cdot \mathbf{r} = Jr \cos(\phi - \phi_0)$, where ϕ and ϕ_0 are angles between horizontal axis and the vectors \mathbf{r} and \mathbf{J} , respectively. Derive that the shape of the particle's trajectory is expressed by

$$r(\phi) = \frac{p}{1 + e \cos(\phi - \phi_0)},$$

where $p = L^2/m\alpha$ and $e = J/\alpha = \sqrt{1 + 2EL^2/m\alpha^2}$. What are interpretations of these parameters? Think about the role of the vector \mathbf{J} in this solution.

2 Week II, 07-13/03/2022

2.1 Lecture

I - Symmetries in Quantum Mechanics

&1. *Axioms of quantum mechanics* - Postulates of quantum mechanics, Ehrenfest theorem, conservation laws.

&2. *Symmetry transformations* - definition of a symmetry transformation in quantum mechanics, Wigner's theorem, conservation laws obtained from a symmetry, infinitesimal symmetry transformations, symmetry generators as observables, symmetry and degeneracy, classification of different symmetry transformations: continuous (space translations, time translations, rotations) and discrete (periodic translation in space, periodic translation in time, parity, time reversal).

&3. *Continuous symmetry transformations* - active and passive view on space and time transformations, translation in space, infinitesimal translation and its generator, symmetry operator of arbitrary translation, homogeneity of space and conservation of momentum, translation in time, infinitesimal translation and its generator, symmetry operator of arbitrary translation, homogeneity of time and conservation of energy, rotation in space, infinitesimal rotation and its generator, symmetry operator of arbitrary rotation, isotropy of space and conservation of angular momentum.

2.2 Tutorial

1. *Ehrenfest theorem* - Prove the Ehrenfest equation.
2. *Translational symmetry operator* - Construct the unitary translation operator $\hat{U}(\mathbf{a}) = e^{-i\mathbf{a} \cdot \hat{\mathbf{p}}/\hbar}$ for an arbitrary translation vector \mathbf{a} .
3. *Translational symmetry operator* - Let $\hat{U}(\mathbf{a})$ is a translation operator and $\hat{O}(\mathbf{a})$ is an observable operator. Show that $\hat{U}(\mathbf{a})^\dagger \hat{O}(\mathbf{a}) \hat{U}(\mathbf{a}') = \hat{O}(\mathbf{a} + \mathbf{a}')$, using that $\hat{U}(\mathbf{a})\psi(\mathbf{r}) = \psi(\mathbf{r} - \mathbf{a})$ and $\hat{U}(\mathbf{a}')^\dagger \psi(\mathbf{r}) = \psi(\mathbf{r} + \mathbf{a}')$, where $\psi(\mathbf{r})$ is a wave function.
4. *Conservation law in a uniform external electric field* - Derive a quantum mechanical generator for the translational symmetry of a charged particle in a homogeneous electric field with the intensity \mathbf{E} .
5. *Antiunitary operators* - Let \hat{A} is an antiunitary operator and $|\tilde{u}\rangle = \hat{A}|u\rangle$ and $|\tilde{v}\rangle = \hat{A}|v\rangle$. Show that $\langle \tilde{u} | \tilde{v} \rangle = \langle v | u \rangle = \langle u | v \rangle^*$.

2.3 Homework problems

1. *Algebraic relations for translation operators* - Show that

$$\left(\frac{i}{\hbar}\hat{p}\right)^n \hat{B}(x) = \sum_{\nu=0}^n \binom{n}{\nu} \frac{\partial^\nu \hat{B}}{\partial x^\nu} \left(\frac{i}{\hbar}\hat{p}\right)^{n-\nu},$$

where \hat{p} is a momentum operator and $\hat{B}(x)$ is every differentiable operator. Next, using the result above, calculate $\hat{U}(a)^\dagger \hat{A}(x) \hat{U}(a)$ where $\hat{U}(a) = e^{-ia\hat{p}/\hbar}$.

2. *Conservation law in a uniform external magnetic field* - Derive a quantum mechanical generator for the translational symmetry of a charged particle in a homogeneous magnetic field with the induction \mathbf{B} .

3 Week III, 14-20/03/2022

3.1 Lecture

&4 *Discrete symmetry transformations* - Discrete translational symmetry in space, periodic potential and primitive translational vectors of a crystal structure (lattice), Bloch theorem and simultaneous eigenstates of symmetry operator of discrete translations and a periodic Hamiltonian, Bloch wave function, quasimomentum in crystals, discrete translations in time, time dependent periodic Hamiltonian, properties of the evolution operators for periodic Hamiltonians, Floquet Hamiltonian, Floquet theorem, Floquet eigenstates in time periodic systems, parity transformation, parity transformation in classical physics, polar and axial vectors and examples, role of parity transformation in quantum mechanics, transformation of different operators under the parity, conservation of parity for parity symmetric Hamiltonians, classification of energy eigenstates under their parity symmetry, even and odd states, time reversal transformation, reversal of time in classical physics, Newton law, transformation of position, velocity, momentum, force, Maxwell equations, transformation of current, electric intensity, magnetic induction, to be continued ...

3.2 Tutorial

1. *Antiunitary operator acting on bra* - How to define an action of the antiunitary operator \hat{B} on a bra vector (to the left)

$$\langle v | \hat{B} = ?$$

2. *Adjoint to antiunitary operator* - How to define an adjoint operator \hat{B}^\dagger to an antiunitary operator \hat{B} ?
3. *Product of unitary and antiunitary operators* - Let $\hat{\Psi} = \hat{\Theta}\hat{U}$, where \hat{U} is a unitary operator and $\hat{\Theta}$ is an antiunitary operator. Show that $\hat{\Psi}$ is antiunitary.

4. *Base dependence in representing an antiunitary operator* - Let \hat{K} is a complex conjugation operator. Show that \hat{K} is not independent of the phase of the basis vectors in terms of which it is defined.
5. *Rotation of spin one particle wave function* - Find a transformation operator for a three-component vector wave function (field). Conclude that it describes a spin one particle.
6. *Derivation of Pauli equation* - Consider an invariant Hamiltonian

$$\hat{H} = \frac{(\vec{\sigma} \cdot \hat{\mathbf{p}})^2}{2m},$$

where $\vec{\sigma}$ is a three component vector made of 2×2 matrices. Show that if these matrices obey an algebra of Pauli matrices

$$\sigma_i \sigma_j + \sigma_j \sigma_i = 2\delta_{ij},$$

$$[\sigma_i, \sigma_j] = 2i\epsilon_{ijk}\sigma_k,$$

then the Hamiltonian is equivalent to the one for free particles. For this you need to show

$$(\vec{\sigma} \cdot \mathbf{a})(\vec{\sigma} \cdot \mathbf{b}) = \mathbf{a} \cdot \mathbf{b} + i\vec{\sigma} \cdot (\mathbf{a} \times \mathbf{b}).$$

Next, introducing a magnetic field via the vector potential and the minimal coupling procedure $\hat{\mathbf{p}} \rightarrow \hat{\mathbf{p}} - q\mathbf{A}$ derive the Pauli Hamiltonian for a spin 1/2 particles in an external magnetic field \mathbf{B} . This problem follows an article in Am. J. Phys. **49**, 645 (1981).

3.3 Homework problems

1. *Product of two antiunitary operators* - Let $\hat{\Psi}$ and $\hat{\Theta}$ are antiunitary operators. Show that $\hat{U} = \hat{\Psi}\hat{\Theta}$ is unitary.
2. *Rotation of spin one-half particle wave function* - Find how the two-component spinor wave function is transformed under rotations. Show that such a wave function describes a spin one-half particle. Hint: to find the transformation rules for the bispinor wave function you need to discuss an invariance of the probability density and the Pauli equation, cf. W. Greiner's book.

4 Week IV, 21-27/03/2022

4.1 Lecture

... continued, problem with a unitary time reversal operator in quantum mechanics, antiunitary time reversal operator, classification of operators regarding time reversal operation, transformation of a scalar wave function under reversing a time, transformation of spin under time reversing, Kramers degeneracy.

II - Spontaneous Symmetry Breaking in Quantum Mechanics

&1. *Methaphysic* - Theory of almost everything (around us), symmetries of general many-body non-relativistic quantum Hamiltonian, existence of phases without a symmetry of their Hamiltonians, some examples, thermodynamic limit, emergency principle.

&2. *Spontaneous symmetry breaking* - Definition of spontaneous symmetry breaking and a Bogoliubov method of its detection, concept of an order parameter, singular (non-commuting) limits in perturbation and number of particles, mathematical and mechanical examples of singular limiting procedure. For more reading: SciPost Phys. Lect. Notes **11**, (2019).

4.2 Tutorial

1. *Dynamical symmetry in hydrogen atom, Runge-Lenz vector* - For the Hamiltonian

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} - \frac{\kappa}{r}$$

we introduce the Runge-Lenz vector

$$\hat{\mathbf{M}} = \frac{1}{2m}(\hat{\mathbf{p}} \times \hat{\mathbf{L}} - \hat{\mathbf{L}} \times \hat{\mathbf{p}}) - \kappa \frac{\mathbf{r}}{r}.$$

One can show that

$$[\hat{\mathbf{M}}, \hat{H}] = 0$$

$$\hat{\mathbf{L}} \cdot \hat{\mathbf{M}} = \hat{\mathbf{M}} \cdot \hat{\mathbf{L}} = 0$$

$$\hat{\mathbf{M}}^2 = \frac{2}{m}\hat{H}(\hat{\mathbf{L}}^2 + \hbar^2) + \kappa^2,$$

and

$$[\hat{L}_i, \hat{L}_j] = i\hbar\epsilon_{ijk}\hat{L}_k$$

$$[\hat{M}_i, \hat{L}_j] = i\hbar\epsilon_{ijk}\hat{M}_k$$

$$[\hat{M}_i, \hat{M}_j] = -2\frac{\hat{H}}{m}i\hbar\epsilon_{ijk}\hat{L}_k.$$

We consider bound states subspace of the Hilbert space with $E < 0$ and perform the rescaling of the Runge vector

$$\hat{\mathbf{M}}' = \sqrt{-\frac{m}{2\hat{H}}}\hat{\mathbf{M}}$$

and obtain the SO(4) commutator algebra

$$[\hat{L}_i, \hat{L}_j] = i\hbar\epsilon_{ijk}\hat{L}_k$$

$$[\hat{M}'_i, \hat{L}_j] = i\hbar\epsilon_{ijk}\hat{M}'_k$$

$$[\hat{M}'_i, \hat{M}'_j] = i\hbar\epsilon_{ijk}\hat{L}_k.$$

We introduce new operators

$$\hat{\mathbf{I}} = \frac{1}{2}(\hat{\mathbf{L}} + \hat{\mathbf{M}})$$

$$\hat{\mathbf{K}} = \frac{1}{2}(\hat{\mathbf{L}} - \hat{\mathbf{M}})$$

and we show that

$$[\hat{I}_i, \hat{I}_j] = i\hbar\epsilon_{ijk}\hat{I}_k$$

$$[\hat{K}_i, \hat{K}_j] = i\hbar\epsilon_{ijk}\hat{K}_k$$

$$[\hat{L}_i, \hat{K}_j] = 0.$$

The eigenstates of these operators and the Hamiltonian are $\{|im_i km_k\rangle\}$ with eigenvalues

$$\hat{\mathbf{I}}^2 \rightarrow \hbar^2 i(i+1)$$

$$\hat{I}_z \rightarrow \hbar m_i$$

$$\hat{\mathbf{K}}^2 \rightarrow \hbar^2 k(k+1)$$

$$\hat{K}_z \rightarrow \hbar m_k$$

and

$$\hat{H} \rightarrow E_n = -\frac{m\kappa^2}{2\hbar^2 n^2},$$

where $n = 2i+1 \in N$ and $i = k$ was used from the orthogonality constraint. Discussion of symmetry and degeneracy of the hydrogen eigenstates.

4.3 Homework problems

1. *Identity with $\hat{\mathbf{M}}$* - Show that

$$\hat{\mathbf{M}}^2 = \frac{2}{m} \hat{H}(\hat{\mathbf{L}}^2 + \hbar^2) + \kappa^2.$$

2. *One of the commutator involving $\hat{\mathbf{M}}$ and $\hat{\mathbf{L}}$* - Show that

$$[\hat{M}_1, \hat{L}_2] = i\hbar \hat{M}_3.$$

3. *One of the commutator involving $\hat{\mathbf{M}}$ and $\hat{\mathbf{L}}$* - Show that

$$[\hat{M}_1, \hat{L}_1] = 0.$$

4. *For ambitious students* - Prove by yourself all equations presented in the tutorial problem.

5 Week V, 28/03/-03/04/2022

5.1 Lecture

&3. *Spontaneous symmetry breaking in quantum mechanical model - harmonic crystal* - Model of a harmonic crystal in one-dimension, creation and annihilation operator, diagonalization of the Hamiltonian via Fourier transform and Bogoliubov transform, analysis of the excitation spectrum, emergence of new quasiparticles - phonons - Goldstone modes, separate treatment of the zero momentum part of the Hamiltonian, total momentum of the system, existence of *thin spectrum*, its role in thermodynamics and in translationally symmetry breaking in the thermodynamic limit, adding a small translationally symmetry breaking perturbation, non-commutative limits. Based on: Am. J. Phys. **75**, 636 (2007).

&4. *Ferromagnet - a prominent exception* - The Heisenberg model, rotational symmetry of the Heisenberg model, exact ferromagnetic ground state without rotational symmetry, co be continued ...

5.2 Tutorial

1. *Harmonic oscillator with driven time-dependent force* - Find the exact solution of the problem with one-dimensional quantum harmonic oscillator in the presence of a driving force and described by the following Schrödinger equation

$$i\hbar \frac{\partial \Psi(x,t)}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x,t)}{\partial x^2} + \frac{1}{2} m \omega^2 x^2 \Psi(x,t) - xF(t)\Psi(x,t),$$

where $F(t)$ is a time dependent force. Next, discuss an explicit solution for a periodic driving force $F(t) = A \sin(\Omega t)$ and check the validity of the Floquet theorem. Based on P. Hänggi, *Quantum transport and dissipation*, chapt. 5.

2. *Parity and spherical harmonics* - Discuss how spherical harmonics are transformed under the parity operation.
3. *Time reversal and spherical harmonics* - Discuss how spherical harmonics are transformed under the time reversal operation.

5.3 Homework problems

1. *A quantum particle with a time dependent potential* - Find the exact solution for a problem of a one-dimensional quantum particle described by the following Schrödinger equation

$$i\hbar \frac{\partial \Psi(x,t)}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x,t)}{\partial x^2} - V(t)\Psi(x,t),$$

where $V(t)$ is a time dependent potential, constant in space. Find a solution for time-periodic potential $V(t) = V_0 \sin(\Omega t + \theta)$. Check the validity of the Floquet theorem.

2. *A quantum particle in a gravity field with a time dependent force* - Find the exact solution for a problem of a one-dimensional quantum particle in a gravity field described by the following Schrödinger equation

$$i\hbar \frac{\partial \Psi(x,t)}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x,t)}{\partial x^2} + mgx\Psi(x,t) - xF(t)\Psi(x,t),$$

where $F(t)$ is a time dependent force, constant in space, and $x \geq 0$. Find a solution for time-periodic force $F(t) = A \sin(\Omega t)$. Check the validity of the Floquet theorem. Based on arXiv:2202.01213.

6 Week VI, 04-10/04/2022

6.1 Lecture

...cont. definition of orientation vector and its classical behavior in the thermodynamic limit, orthogonality of different oriented ferromagnetic states in the thermodynamic limit, symmetry breaking and classical

behavior, low energy excitations of ferromagnets, quadratic dispersion of magnons, spin-waves.

III. Galilean and gauge transformations in quantum mechanics

&1. *Landé-Lévy-Lebonds pseudoparadox* - Discussion of de Broglie $p = h/\lambda$ and Einstein $E = h\nu$ hypothesis in laboratory and moving frames, Galilean transformation in classical wave physics and classical Doppler effect, based on: Am. J. Phys. **44**, 1130 (1974).

&2. *Galilean transformation in quantum mechanics* - Wigner symmetry operator for Galilean shift of position and momentum operators, explicit derivation of this operator

$$\hat{U}(\mathbf{v}, t) = e^{\frac{i}{\hbar}(\hat{\mathbf{p}}t - m\hat{\mathbf{r}})\mathbf{v}},$$

wave function transformation between two moving frames

$$\Psi_T(\mathbf{r}_T, t) = e^{\frac{i}{\hbar}(\frac{m\mathbf{v}^2}{2}t - m\mathbf{v}\mathbf{r})}\Psi(\mathbf{r}, t),$$

role of the local phase change in the wave function, resolution of the Landé's pseudoparadox, quantum Doppler effect

$$\frac{1}{\lambda'} = \frac{1}{\lambda} - \frac{mv}{h},$$

$$v' = v - \frac{v}{\lambda} + \frac{mv^2}{2h}.$$

&3. *Non-equivalent representations of momentum operators in quantum mechanics* - Position and momentum operators in quantum mechanics and their commutator, explicit form of these operators in position representation, to be continued

6.2 Tutorial

1. *Rotation of a spin around one of the coordinate axis* - For the spin rotation operator with the angle π around the axis y ,

$$\hat{U} = e^{-i\frac{\pi}{\hbar}\hat{S}_y},$$

show that $\hat{U}\hat{S}_x = -\hat{S}_x\hat{U}$ and $\hat{U}\hat{S}_z = -\hat{S}_z\hat{U}$.

2. *Rotation operator for fermions* - Find an explicit form of the spin rotation operator $\hat{U}(\phi \mathbf{n})$ for spin 1/2 particles. Show that $\hat{U}(2\pi \mathbf{n}) = -1$. ϕ is the rotation angle around the \mathbf{n} direction.
3. *Tight binding model* - Consider an infinite one-dimensional system with a periodic potential $V(x \pm a) = V(x)$. Let $|\psi_n\rangle$ be a ground state vector describing a particle localized in the n -th cell of the crystal. The ground state energy is E_0 . Find an eigenstate vector both of the Hamiltonian \hat{H} and of the discrete translation operator $\hat{U}(a)$. Assume that,

$$\langle \psi_n | \psi_m \rangle = \delta_{nm},$$

$$\langle \psi_n | \hat{H} | \psi_n \rangle = E_0,$$

$$\langle \psi_n | \hat{H} | \psi_{n\pm 1} \rangle = -\Delta \ll 0,$$

and other amplitudes vanish. Discuss the Bloch theorem and the dispersion relation of particles.

6.3 Homework problems

1. *Chain molecule - Tight binding model* - Consider a chain molecule of N atoms. Find the eigenstates and eigenenergies of such a system. Hints: Take a one-particle localized base $\{|j\rangle\}$ and expand any state

$$|\psi\rangle = \sum_{j=1}^N c_j |j\rangle.$$

Solve the Schrodinger equation

$$\hat{H}|\psi\rangle = E|\psi\rangle,$$

assuming that $\langle j | \hat{H} | j \rangle = \alpha$ and $\langle j | \hat{H} | k \rangle = \beta$ for j and k nearest neighbors, and zero otherwise. Prove that $E = \alpha + 2\beta \cos(m\pi/(N+1))$ and $c_j^m = \sqrt{2/(N+1)} \sin(mj\pi/(N+1))$. Discuss a transition from a single atom to an infinite system and appearance of the continuum band.

2. *Ring molecule - Tight binding model* - Consider a ring molecule of N atoms. Find the eigenstates and eigenenergies of such a system. Hints: Take a one-particle localized base $\{|j\rangle\}$ and expand any state

$$|\psi\rangle = \sum_{j=1}^N c_j |j\rangle.$$

Solve the Schrodinger equation

$$\hat{H}|\psi\rangle = E|\psi\rangle,$$

assuming that $\langle j | \hat{H} | j \rangle = \alpha$ and $\langle j | \hat{H} | k \rangle = \beta$ for j and k nearest neighbors, and zero otherwise. Impose the periodic boundary conditions and show that $E = \alpha + 2\beta \cos(2\pi m/N)$ and $c_j^m = \exp(i2\pi jm/N)/\sqrt{N}$. Discuss a transition from few atoms to the thermodynamic limit. Discuss the Bloch theorem in the finite and in the infinite systems.

7 Week VII, 11-17/04/2022

7.1 Lecture

... cont., momentum operator defined with respect to an arbitrary function, $\hat{x} = x$, $\hat{p} = -i\hbar d/dx + \chi(x)$, invariance of canonical commutation relations, transformation of the base $\{|x\rangle\}$ by multiplying by the phase factor $\exp(ig(x)/\hbar)$, where $g(x) = \int^x \chi(y)dy$, unitary change of the base, higher dimensional generalization, path independence of the phase $g(\mathbf{r}) = \int^{\mathbf{r}} \chi(\mathbf{r}')d\mathbf{r}'$, rotationless of the function $\nabla \times \chi(\mathbf{r}) = 0$, non-unique method in non-compact spaces.

&4. *Electromagnetic gauge representations in quantum mechanics* - reminder of scalar and vector potentials in classical electrodynamics, gauge freedom in classical electrodynamics, some popular gauges, incorporation of electromagnetic field in quantum mechanics, minimal coupling prescription, mechanical momentum vs. dynamical momentum, change of the wave

function and states by adding the vector potential, electromagnetic gauge transformations in quantum mechanics.

&5. *Aharonov-Bohm effect* - model of infinite solenoid, absence of magnetic induction outside the solenoid tube, motion of a charge quantum particle outside the tube, to be continued....

7.2 Tutorial

1. *Lattice (discrete) Fourier transform* - discussion of the discrete Fourier transform for periodic sequence and lattice sums.
2. *Model of quantum harmonic crystal* - Detailed solution of the quantum harmonic crystal model in one dimension with periodic boundary condition: rising operators in lattice space, lattice Fourier transformation, Bogoliubov-Valatin diagonalization.
3. *Model of quantum harmonic crystal* - Solve this model by Fourier transform and rising operators in momentum space.

7.3 Homework problems

1. *Model of quantum harmonic crystal with two atoms unit cell* - Consider one dimensional model of harmonic crystal with two different ions of masses m_1 and m_2 placed in an alternative manner. Distances between atoms are a and the interaction is of harmonic (quadratic) type between nearest neighbor ions. Find the excitation spectrum of this system. Check that there are two branches of excitations one gapless (Goldstone) and the other gapped, with finite energy at $k = 0$. Check the limit $m_1 = m_2$.

8 Week VIII, 18-24/04/2022

8.1 Lecture

Easter holiday, no lecture.

8.2 Tutorial

1. *Poor men's version of Hohenberg-Mermin-Wagner theorem, instability of $d = 1$ crystal* - Calculate the mean square deviation of the n -th ion from its equilibrium position at $T = 0$, i.e. $\langle 0 | \hat{x}_n^2 | 0 \rangle$, within the model of quantum harmonic crystal.
2. *Spin-wave (magnon) excitation spectrum* - Find a dispersion relation of the spin-wave excitations around the ferromagnetic ground state of $d = 1$ Heisenberg model.
3. *Einstein's gedanken experiment with a rocket* - Consider a rocket in an empty space and subjected to an acceleration equal to g . Suppose that inside the rocket there is a single quantum object,

described by a Schrödinger equation with a confining potential $V(x, t)$, e.g. the rocket walls, in the laboratory frame. Find the Schrödinger equation in a reference frame co-moving with the rocket. Interpret this result.

8.3 Homework problems

1. *Correlation function in the model of quantum harmonic crystal* - Within the model of quantum harmonic crystal determine the correlation function $\langle 0 | (\hat{x}_n - \hat{x}_{n-l})^2 | 0 \rangle$ in the ground state. Discuss the nearest neighbor bond length at $l = 1$ and discuss the limit $l \rightarrow \infty$.
2. *Spin-wave (magnon) excitation spectrum in d -dimensions* - Find a dispersion relation of the spin-wave excitations (collective superposition of states with only one spin flipped) around the ferromagnetic ground state of a Heisenberg model with the exchange interaction between spins localized on nearest neighbor sites of a hypercubic lattice with the lattice constant a in d -dimensions. Hint: repeat steps from tutorials but the discrete Fourier transform must be d -dimensional.

9 Week IX, 25/04-01/05/2022

9.1 Lecture

... cont., relative phase of the wave functions and its dependence on the magnetic flux, unit of quantum flux.

&6 *Dirac magnetic monopole* - history of a magnetic monopole and the consideration of Dirac, theorem of non-existence of the single vector potential corresponding to the magnetic monopole, Wu and Yang consideration about the magnetic monopole, two different vector potentials covering the whole sphere in 3-dimensional space, gauge transformation between these potentials, quantization of the magnetic monopole due to single valuedness of the wave function, topological character of this quantization.

IV. Berry phases and topological states in quantum mechanics

&1. *Adiabatic approximation* - adiabatic processes and adiabatic invariants in classical mechanics, two different time scales, to be continued ...

9.2 Tutorial

Mid term exam:

Problem 1.

A particle with mass m is moving along the one dimension line in the presence of a time and space dependent potential

$$V(x, t) = V_\infty(x) |\sin(\Omega t)|$$

where

$$V_\infty(x) = \begin{cases} V_0 & \text{for } 0 < x < L \\ \infty & \text{for otherwise.} \end{cases}$$

(Assume that the hard wall potential is 'extremely hard', i.e. $\infty \cdot 0 = \infty$, and $V_0 > 0$.) In the initial moment at $t = 0$ the particle is in a quantum state given by the wave function $\Phi(x, t = 0) = \mathcal{N} \sin^3(\frac{\pi}{L}x)$. Find the normalization factor \mathcal{N} and the wave function $\Phi(x, t)$ at a given time t for: a) $0 < t < \pi/\Omega$, and b) $\pi/\Omega < t < 2\pi/\Omega$. Plot the time change of the global phase factor, i.e. that which depends on V_0 , between 0 and $2\pi/\Omega$. Discuss if this plot depends on a specific form of the initial state from the Hamiltonian domain. Hint: $\sin^3 a = (3 \sin a - \sin 3a)/4$.

Problem 2.

Two particles of mass m_1 and m_2 in one dimension are interacting with each other with the potential $V(|x_1 - x_2|)$ and the Hamiltonian of this system is

$$\hat{H} = \frac{\hat{p}_1^2}{2m_1} + \frac{\hat{p}_2^2}{2m_2} + V(|x_1 - x_2|).$$

Translation operator acting on the wave function gives

$$\hat{U}(a)\psi(x_1, x_2) = \psi(x_1 - a, x_2 - a).$$

a) Show that

$$\hat{U}(a) = e^{-\frac{i}{\hbar}a\hat{P}},$$

where $\hat{P} = \hat{p}_1 + \hat{p}_2$ is a total momentum.

b) Show that the total momentum is conserved.

10 Week X, 02-08/05/2022

10.1 Lecture

3 days May's holiday

10.2 Tutorial

no tutorial, special Faculty schedule

11 Week XI, 09-15/05/2022

11.1 Lecture

... cont., quantum mechanical examples of adiabatic processes: infinite quantum well and Born-Oppenheimer approximation, adiabatic theorem in quantum mechanics and its prove.

&2. *Berry (geometric) phases* - Anholonomy in case of periodic processes, example of pendulum on the Earth surface, Foucault pendulum and Hannay's angle, relation of this angle to solid angle, geometric phase in quantum mechanics, the approximate form of the wave function in case of adiabatic processes, definition of dynamical and geometric phases, condition on the geometric phase, expression on the geometric phase in case of the change of the Hamiltonian parameters, flux interpretation of geometric phase, Berry's potential, Berry's flux, is the geometric phase real? is the geometric phase measurable?

11.2 Tutorial

1. *Mid-term exam problems* - discussion of exam problems
2. *Gauge transformation in quantum mechanics* - Check by explicit calculation that the form of the Schrödinger equations before and after the gauge transformation are the same.
3. *Quantum particle in uniform electric field* - Solve the problem of a quantum particle moving in a static uniform electric field $\mathbf{E} = E_0\mathbf{e}_x$. Solve the problem in: a) *static gauge* with $V = -E_0x$, $\mathbf{A} = 0$, and b) *dynamic gauge* with $V = 0$ and $\mathbf{A} = -E_0t\mathbf{e}_x$. Using these solutions, obtain solutions in the other gauges by taking a gauge transformation with $\chi = -E_0xt$ scalar function.

11.3 Homework problems

1. *Quantum particle in a gravity field* - Solve a problem of a quantum particle with mass m falling in a uniform gravity field $V = mgz$ for $z > 0$, and assuming that for $z \leq 0$ the potential is of infinite height (hard wall potential). Discuss an energy quantization and a motion of a classical (gaussian) wave packet. Compare to results in an infinite space with the uniform gravity field, i.e. without the hard wall boundary. Hint: Am. J. Phys. **67**, 776 (1999), arxiv:2009.03744.
2. *Gauge invariant density, current, and continuity equation* - For a quantum particle with charge in the presence of an electromagnetic field find (guess) forms for the particle density and the current density, such that they are gauge invariant. Check if the continuity equation is gauge invariant as well.

12 Week XII, 16-22/05/2022

12.1 Lecture

&3. *Aharonov-Bohm phase as a Berry phase* - reinterpretation of the Aharonov-Bohm geometry in terms of adiabatic change of the system and interpretation of the magnetic flux as a geometric phase.

&4. *Emergent of a Berry monopole in a two level system* - generic form of a zero dimensional, two-level quantum mechanical Hamiltonian, unique eigenvalues and non-unique eigenvectors, necessity of using two gauges (e.g. north and south) determination of the Berry gauge dependent potential vector and the Berry gauge independent flux, magnetic monopole interpretation of the Berry vector potentials, a monopole at the degeneracy point in the Hamiltonian parameter space, emergence of the gauge structure in the adiabatically changed two level system.

12.2 Tutorial

1. *DC Josephson effect* - Using a simplified two-state model due to Feynman derive the Josephson equation on the tunneling current $J = J_0 \sin(\theta_2 - \theta_1)$ between two superconductors with fixed phases of their wave functions.
2. *AC Josephson effect* - Using a simplified two-state model due to Feynman derive the Josephson equation on the tunneling current between two superconductors with fixed phases of their wave functions which are biased by the gate voltage $V(t)$. Analyze in details the case where $V(t) = V_0$.

12.3 Homework problems

1. *Gauge invariant current density in superconductors* - Assuming the wave function as $\Psi(\mathbf{r}, t) = \sqrt{n(\mathbf{r}, t)} e^{i\theta(\mathbf{r}, t)}$ and show that the current density is $\mathbf{J} = qn\mathbf{v}$, where $\mathbf{v} = \frac{\hbar}{m} \nabla\theta - \frac{q}{m} \mathbf{A}$. Check gauge invariance.
2. *AC Josephson effect - Shapiro steps* - Using a simplified two-state model due to Feynman derive the Josephson equation on the tunneling current between two superconductors with fixed phases of their wave functions which are biased by the gate voltage $V(t)$. Analyze in details the case where $V(t) = V_0 + V_1 \cos(\omega t)$.

13 Week XIII, 23-29/05/2022

13.1 Lecture

V. Scattering Theory

&1. *Introduction to scattering experiments* - typical scattering geometry, conservation laws in scattering experiments, elastic and inelastic scatterings, types of scattering experiments.

&2. *Scattering cross section* - incoming current and scattered current and their units, definition of differential and total cross sections, geometrical interpretation of the total cross section.

&3. *General theory of scattering - Lippmann-Schwinger equation* - derivation of a formal solution of the Schrödinger equation for continuous energies in term of Lippmann-Schwinger equation.

&4. *Resolvents, Green's functions and Lippmann-Schwinger equation* - definition of a resolvent for a given Hamiltonian $\hat{G}(E) = (E - \hat{H})^{-1}$, application of the η trick, $E \rightarrow E + i\eta$ with $\eta \rightarrow 0^+$, formulation of the Lippmann-Schwinger equation using the resolvent, formal solution of the Schrödinger equation by using $\hat{G}(E)$, relation of the η trick with the retarded boundary condition, formulation of the problem in terms of a time dependent Schrödinger equation and its solution by retarded Green's function, prove of the relation between the retarded boundary (initial) conditions with the $\eta \rightarrow 0^+$ procedure.

&5. *Formal solution of the Lippmann-Schwinger equation* - iterative solution to Lippmann-Schwinger equation, summation of the geometric progress, definition of

the t-matrix operator (transition matrix), t-matrix as an effective scattering potential, to be continued ...

13.2 Tutorial

1. *SQUID - Superconducting QUantum Interference Device* - Consider two superconductors forming a loop with two Josephson junctions. Derive the formula for the current as a function of the magnetic flux inside the loop.
2. *Beyond the adiabatic limit* - Taking the wave function expansion

$$\Psi_n(x, t) = \psi_n(x, t) e^{i\Theta_n(t)} e^{i\gamma_n(t)} + \sum_{m \neq n} c_m(t) \psi_m(x, t)$$

check the correctness of the expression for the geometric phase up to the first order in the ϵ .

3. *Berry phase in the Born-Oppenheimer approximation* - Consider two-particle hamiltonian

$$\hat{H} = \frac{\hat{\mathbf{P}}^2}{2M} + \frac{\hat{\mathbf{p}}^2}{2m} + V(\mathbf{R}, \mathbf{r}),$$

where small and great letters correspond to light and heavy particles. In the limit $M \gg m$ one can take an ansatz on the wave function

$$\psi_n(\mathbf{R}, \mathbf{r}) = \phi_n(\mathbf{R}) \xi_n(\mathbf{R}, \mathbf{r}),$$

where the eigen-problem for fast degrees of freedom (m)

$$\hat{h}(\mathbf{R}) |n\mathbf{R}\rangle = \epsilon_n(\mathbf{R}) |n\mathbf{R}\rangle$$

or

$$\hat{h}(\mathbf{R}) \xi_n(\mathbf{R}, \mathbf{r}) = \epsilon_n(\mathbf{R}) \xi_n(\mathbf{R}, \mathbf{r}),$$

depends on \mathbf{R} only parametrically. Show that the eigen-problem for the slow degrees of freedom (M) is described by the Hamiltonian

$$\hat{H}_{\text{eff}} = \frac{1}{2M} (\hat{\mathbf{P}} + \hbar \mathbf{A}_B)^2 + \epsilon_n(\mathbf{R}),$$

where $\mathbf{A}_B = i \int d_3r \xi_n(\mathbf{R}, \mathbf{r})^* \nabla_{\mathbf{R}} \xi_n(\mathbf{R}, \mathbf{r})$ is the Berry phase. Interpret these results.

13.3 Homework problems

1. *Expanding adiabatically quantum well* - Calculate the geometric phase γ_n in the problem of infinite quantum well in one dimension if the well expands adiabatically from L_1 to L_2 . If the expansion happens at a constant rate $dL/dt = v$ determine the dynamic phase Θ_n . If the well contracts to the original size, what is the Berry's phase? Make comments on these partial results.

2. *Particle in a rotating magnetic field* - An external magnetic field changes in time as

$$\mathbf{B}(t) = \begin{pmatrix} B_0 \sin \alpha \cos \omega t \\ B_0 \sin \alpha \sin \omega t \\ B_0 \cos \alpha \end{pmatrix}.$$

Assume that at $t = 0$ the particle's spin points along the magnetic field. Check that the exact solution is in the form

$$\chi(t) = \begin{pmatrix} (\cos(\lambda t/2) + i \frac{\omega_1 + \omega}{\lambda} \sin(\lambda t/2)) \cos(\alpha/2) e^{-i\omega t/2} \\ (\cos(\lambda t/2) + i \frac{\omega_1 - \omega}{\lambda} \sin(\lambda t/2)) \sin(\alpha/2) e^{i\omega t/2} \end{pmatrix},$$

where $\omega_1 = qB_0/m$ and $\lambda = \sqrt{\omega^2 + \omega_1^2 + 2\omega\omega_1 \cos \alpha}$. Compute the probability that the particle's spin will be anti-parallel to the magnetic field. Find the adiabatic conditions under which this probability vanishes. Compute the Berry phase during a single cycle of rotation of the magnetic field.

3. *Two level system - Berry phase* - For a two level system discussed in the lecture derive all results on Berryology in details.

14 Week XIV, 30/05-05/06/2022

14.1 Lecture

... continued, Lippmann-Schwinger equation for the wave function, Born series for the wave function and for the t-matrix, multiple scattering interpretation of the Born series expansion.

&6. *Formal solution of the scattering problem in terms of the Green's function* - Resolvents and Green's functions for noninteracting and interacting problems, density of states and its relation to the Green's function, formal solution for the resolvent in terms of the t-matrix, proof of the Friedel sum rule theorem, proof of the optical theorem, definition of the s-matrix in on-shell scattering and its relation to the t-matrix, to be continued ...

14.2 Tutorial

1. *Resolvent in one-level model* - Find the resolvent and the Green's function for a system with the Hamiltonian

$$\hat{H} = \epsilon^a |a\rangle\langle a|.$$

2. *Resolvent in two-level model* - Find the resolvent and the Green's function for a system with the Hamiltonian

$$\hat{H} = \epsilon^a |a\rangle\langle a| + \epsilon^b |b\rangle\langle b| + V |a\rangle\langle b| + V^* |b\rangle\langle a|.$$

Find the corresponding perturbative series with respect to V . Find t-matrix series. Interpret results diagrammatically. This is an elementary introduction to Feynman diagrams.

3. *Green's function in $d = 3$* - Compute the Green's function $G_0(\mathbf{r}, \mathbf{r}'; E)$ for free particles in $d = 3$ dimensions.

14.3 Homework problems

- Green's function in d dimensions* - Compute the Green's function $G_0(\mathbf{r}, \mathbf{r}'; E)$ for free particles in $d = 1, d = 2$ (and arbitrary d if you are interested in) dimensions.
- Density of states in d dimensions* - The local density of states (LDOS) is defined as

$$\rho(\mathbf{r}; E) = -\frac{1}{\pi} \lim_{\mathbf{r}' \rightarrow \mathbf{r}} \text{Im} G_0(\mathbf{r}, \mathbf{r}'; E).$$

Compute this quantity in $d = 1, 2$ and 3 dimensions. Compare with results obtained on your Statistical Physics course.

15 Week XV, 06-12/06/2022

15.1 Lecture

... continued, bound states and states from the continuum, Levinson's theorem and its prove, topological character of the Levinson's theorem.

&7. *Scattering amplitude* - Expansion of the Lippmann-Schwinger equation far away from the scattering potential center, definition of the scattering amplitude, relation of the scattering amplitude and the T-matrix, relation between η -trick, retarded boundary condition and purely outgoing boundary condition, outgoing spherical wave with modified amplitude, the scattering cross-section and the scattering amplitude, different formulations of the optical theorem.

15.2 Tutorial

- Winding number* - Discussion of the winding number (index number) of a complex valued function around a closed curve in a complex plane.
- Friedel oscillations* - Find how the density of electrons oscillates far away from the localized impurity potential in three dimensional space.
- Resolvents in one- and two-level system* - From the definition of the resolvent find it for: one level system with the Hamiltonian $\hat{H} = \epsilon|0\rangle\langle 0|$, two level system with the Hamiltonian $\hat{H} = \epsilon_a |a\rangle\langle a| + \epsilon_b |b\rangle\langle b| + V |a\rangle\langle b| + V^* |b\rangle\langle a|$. Express the solution as a perturbative series in terms of V . Draw the corresponding Feynman diagrams for this problem. Find t-matrix for this problem treating V as a scattering potential. Express t-matrix as a perturbation series and draw the Feynman diagrams.

15.3 Homework problems

- Single site model with only one discrete state* - Find t-matrix, Green's function, scattering phase shift, screening charge in case of a one level ϵ model treating a shift V of the energy level as perturbation. The Hamiltonian is $\hat{H} = \epsilon|0\rangle\langle 0| + V|0\rangle\langle 0|$.

Solution:

The Green function is

$$G_0(z) = \frac{1}{z - \epsilon}, \quad (1)$$

where ϵ is the energy of this level. The DOS is $\rho_0(\omega) = \delta(\omega - \epsilon)$ and the occupation of this system is

$$n_0 = \int \rho_0(\omega) f(\omega) d\omega = \frac{1}{e^{\beta(\epsilon - \mu)} + 1}, \quad (2)$$

where μ is the chemical potential.

We perturb this system by shifting the energy by V . he the Green function is

$$G(z) = \frac{1}{z - \epsilon - V}, \quad (3)$$

the DOS is $\rho(\omega) = \delta(\omega - \epsilon - V)$, and the occupation of this system is

$$n = \int \rho_0(\omega) f(\omega) d\omega = \frac{1}{e^{\beta(\epsilon + V - \mu)} + 1}. \quad (4)$$

The change of the DOS is

$$\Delta\rho(\omega) = \delta(\omega - \epsilon - V) - \delta(\omega - \epsilon), \quad (5)$$

and the screening charge is obtained

$$Z_{sc} = n - n_0 = \frac{1}{e^{\beta(\epsilon + V - \mu)} + 1} - \frac{1}{e^{\beta(\epsilon - \mu)} + 1}. \quad (6)$$

At $T = 0$ it reduces to

$$Z_{sc} = \Theta(\mu - \epsilon - V) - \Theta(\mu - \epsilon). \quad (7)$$

Now we derive the same result within the scattering formalism. For a scalar functions we have an identity

$$G = \frac{d}{d\omega} \ln G^{-1}, \quad (8)$$

which leads to

$$\rho(\omega) = - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \text{Im} \frac{d}{d\omega} \ln G(z)^{-1}. \quad (9)$$

Hence the change of DOS is

$$\begin{aligned} \Delta\rho(\omega) &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln \frac{G(z)^{-1}}{G_0(z)^{-1}} \\ &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln \frac{z - \epsilon - V}{z - \epsilon} \\ &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln \left(1 - \frac{V}{z - \epsilon}\right) \\ &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln \frac{1}{1 - \frac{V}{z - \epsilon}} \\ &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln \frac{1}{V} T(z) \\ &= \frac{1}{\pi} \frac{d}{d\omega} \phi(\omega), \end{aligned} \quad (10)$$

where

$$T(z) = \frac{V}{1 - \frac{V}{z - \epsilon}} \quad (11)$$

is the "T-matrix"(T-scalar) here and $\phi(\omega)$ is the scattering phase shift given by the argument of $T(\omega)$.

Determination of the phase must take into account multivaluedness of the logarithmic function. Indeed from

$$\omega + i0^+ - \epsilon = |\omega - \epsilon| e^{i\pi\Theta(\epsilon - \omega)}, \quad (12)$$

$$\omega + i0^+ - \epsilon - V = |\omega - \epsilon - V| e^{i\pi\Theta(\epsilon + V - \omega)}, \quad (13)$$

we find

$$\begin{aligned} \lim_{z \rightarrow \omega + i0^+} \frac{1}{V} T(\omega) &= \lim_{z \rightarrow \omega + i0^+} \frac{z - \epsilon}{z - \epsilon - V} \\ &= \left| \frac{\omega - \epsilon}{\omega - \epsilon - V} \right| e^{i\pi(\Theta(\epsilon - \omega) - \Theta(\epsilon + V - \omega))}. \end{aligned} \quad (14)$$

Therefore,

$$\begin{aligned} \lim_{z \rightarrow \omega + i0^+} \ln \frac{1}{V} T(z) &= \ln \left| \frac{\omega - \epsilon}{\omega - \epsilon - V} \right| \\ &+ i\pi(\Theta(\epsilon - \omega) - \Theta(\epsilon + V - \omega)). \end{aligned} \quad (15)$$

The imaginary part

$$\begin{aligned} \phi(\omega) &= \pi[\Theta(\epsilon - \omega) - \Theta(\epsilon + V - \omega)] \\ &= \pi[\Theta(\omega - \epsilon - V) - \Theta(\omega - \epsilon)], \end{aligned} \quad (16)$$

since $\Theta(-x) = 1 - \Theta(x)$, is just the scattering phase shift. Although the spectrum is discrete the T-matrix has a nontrivial phase. From the Friedel sum rule we obtain

$$Z_{sc} = \frac{1}{\pi} \phi(\mu) = \Theta(\mu - \epsilon - V) - \Theta(\mu - \epsilon), \quad (17)$$

which agrees with the result before.

The change of DOS is

$$\begin{aligned} \Delta\rho(\omega) &= \frac{1}{\pi} \frac{d}{d\omega} \phi(\omega) \\ &= \frac{d}{d\omega} (\Theta(\epsilon - \omega) - \Theta(\epsilon + V - \omega)) \\ &= \delta(\omega - \epsilon - V) - \delta(\omega - \epsilon). \end{aligned} \quad (18)$$

This agrees with results from direct method given at the beginning, c.f., Eq. (5). Hence the screening charge is correctly the same. Therefore, the Friedel sum rule works for one site (discrete) non-interacting system.

Checking the consistency of choosing the phase

$$x \pm i0^+ = |x| e^{\pm i\pi\Theta(-x)}.$$

We know that

$$\frac{1}{x \pm i0} = P \frac{1}{x} \mp i\pi\delta(x).$$

On the other hand

$$\begin{aligned} \frac{1}{x \pm i0} &= \frac{d}{dx} \ln(x \pm i0^+) \\ &= \frac{d}{dx} \ln(|x| e^{\pm i\pi\Theta(-x)}) = \frac{d}{dx} (\ln|x| \pm i\pi\Theta(-x)) \\ &= P \frac{1}{x} \pm i\pi \frac{d}{dx} \Theta(-x) = P \frac{1}{x} \mp i\pi\delta(-x) \\ &= P \frac{1}{x} \mp i\pi\delta(x). \end{aligned} \quad (19)$$

□

2. *Two degenerate levels model with only one level perturbed* - Consider the model with $\hat{H} = -t|0\rangle\langle 1| - t|1\rangle\langle 0| - v|1\rangle\langle 1|$. Find t-matrix, Green's function, scattering phase shift, screening charge treating a shift v of the energy level as perturbation.

Solution:

The free Green function matrix is

$$\mathbf{G}_0(z) = \begin{pmatrix} z & t \\ t & z \end{pmatrix}^{-1} = \frac{1}{z^2 - t^2} \begin{pmatrix} z & -t \\ -t & z \end{pmatrix}, \quad (20)$$

where the on site energy $\epsilon = 0$ and $t > 0$ is the hopping amplitude between sites. In the perturbed case the second site onsite energy is changed by $-v$, so the scattering potential is

$$\mathbf{V} = \begin{pmatrix} 0 & 0 \\ 0 & -v \end{pmatrix}, \quad (21)$$

and the Green function is

$$\begin{aligned} \mathbf{G}(z) &= \begin{pmatrix} z & t \\ t & z+v \end{pmatrix}^{-1} = \\ &= \frac{1}{z(z+v) - t^2} \begin{pmatrix} z+v & -t \\ -t & z \end{pmatrix}. \end{aligned} \quad (22)$$

The perturbed Green function \mathbf{G} has singularities at

$$\omega_{1,2} = \frac{1}{2}[-v \pm \sqrt{v^2 + 4t^2}], \quad (23)$$

and the free Green function \mathbf{G}_0 is singular at

$$\omega_{1,2}^0 = \pm t. \quad (24)$$

To find DOS we need the trace of \mathbf{G} and \mathbf{G}_0 :

$$\begin{aligned} \text{Tr } \mathbf{G}(z) &= \frac{2z+v}{z(z+v) - t^2} = \\ &= \frac{2z+v}{\omega_1 - \omega_2} \left(\frac{1}{z - \omega_1} - \frac{1}{z - \omega_2} \right) \end{aligned} \quad (25)$$

Observing that $\omega_1 - \omega_2 = \sqrt{v^2 + 4t^2}$ we obtain

$$\text{Tr } \mathbf{G}(z) = \frac{2z+v}{\sqrt{v^2 + 4t^2}} \left(\frac{1}{z - \omega_1} - \frac{1}{z - \omega_2} \right), \quad (26)$$

and in the limit $z \rightarrow \omega + i0^+$

$$\begin{aligned} \text{Tr } \mathbf{G}(\omega) &= \frac{2\omega+v}{\sqrt{v^2 + 4t^2}} \left(P \frac{1}{\omega - \omega_1} - i\pi\delta(\omega - \omega_1) \right. \\ &\quad \left. - P \frac{1}{\omega - \omega_2} + i\pi\delta(\omega - \omega_2) \right). \end{aligned} \quad (27)$$

With observation that $2\omega_{1,2} + v = \pm\sqrt{v^2 + 4t^2}$ the DOS is

$$\begin{aligned} \rho(\omega) &= -\frac{1}{\pi} \text{Im Tr } \mathbf{G}(\omega) \\ &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2). \end{aligned} \quad (28)$$

The unperturbed DOS is

$$\rho(\omega) = \delta(\omega - \omega_1^0) + \delta(\omega - \omega_2^0), \quad (29)$$

and the change of DOS is

$$\begin{aligned} \Delta\rho(\omega) &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2) \\ &\quad - \delta(\omega - \omega_1^0) - \delta(\omega - \omega_2^0). \end{aligned} \quad (30)$$

The screening charge is

$$\begin{aligned} Z_{\text{sc}} &= \int d\omega \Delta\rho(\omega) f(\omega) = \frac{1}{e^{\beta(\omega_1 - \mu)} + 1} + \frac{1}{e^{\beta(\omega_2 - \mu)} + 1} \\ &\quad - \frac{1}{e^{\beta(\omega_1^0 - \mu)} + 1} - \frac{1}{e^{\beta(\omega_2^0 - \mu)} + 1} \end{aligned} \quad (31)$$

and at $T = 0$ reduces to

$$\begin{aligned} Z_{\text{sc}} &= \Theta(\mu - \omega_1) + \Theta(\mu - \omega_2) \\ &\quad - \Theta(\mu - \omega_1^0) - \Theta(\mu - \omega_2^0). \end{aligned} \quad (32)$$

Next we use scattering to determine the change of DOS and the screening charge. From the main text we have

$$\begin{aligned} \Delta\rho(\omega) &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{V}^{-1}(z)\mathbf{T}(z)] \\ &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{1} - \mathbf{V}\mathbf{G}_0]^{-1} \\ &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{1} - \mathbf{V}\mathbf{G}_0]. \end{aligned} \quad (33)$$

The matrix

$$\begin{aligned} \mathbf{1} - \mathbf{V}\mathbf{G}_0 &= \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{1}{z^2 - t^2} \begin{pmatrix} 0 & 0 \\ 0 & -v \end{pmatrix} \begin{pmatrix} z & -t \\ -t & z \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ -\frac{vt}{z^2 - t^2} & 1 + \frac{zv}{z^2 - t^2} \end{pmatrix} \end{aligned} \quad (34)$$

needs to be diagonalized. There are two eigenvalues $\lambda_1 = 1$ and $\lambda_2 = (z(z+v) - t^2)/(z^2 - t^2)$ and logarithm of diagonal matrix is a matrix of logarithm of its eigenvalues. Since the trace is base independent we have

$$\Delta\rho(\omega) = - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} (\ln\lambda_1 + \ln\lambda_2). \quad (35)$$

Now $\ln\lambda_1 = 0$ and

$$\begin{aligned} &\lim_{z \rightarrow \omega + i0^+} \ln \frac{z(z+v) - t^2}{z^2 - t^2} \\ &= \lim_{z \rightarrow \omega + i0^+} (\ln(z^2 + zv - t^2) - \ln(z^2 - t^2)) \\ &= \ln|(\omega - \omega_1)(\omega - \omega_2)| + i\pi(\Theta(\omega_1 - \omega) + \Theta(\omega_1 - \omega)) \\ &\quad - \ln|(\omega - \omega_1^0)(\omega - \omega_2^0)| - i\pi(\Theta(\omega_1^0 - \omega) + \Theta(\omega_1^0 - \omega)). \end{aligned} \quad (36)$$

Therefore we find nonzero scattering phase shift

$$\text{Tr } \phi(\omega) = \pi[\Theta(\omega - \omega_1) + \Theta(\omega - \omega_2) - \Theta(\omega - \omega_1^0) + \Theta(\omega - \omega_2^0)]. \quad (37)$$

The change of DOS

$$\begin{aligned} \Delta\rho(\omega) &= \frac{1}{\pi} \frac{d}{d\omega} \pi[\Theta(\omega - \omega_1) + \Theta(\omega - \omega_2) \\ &\quad - \Theta(\omega - \omega_1^0) + \Theta(\omega - \omega_2^0)] \\ &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2) \\ &\quad - \delta(\omega - \omega_1^0) - \delta(\omega - \omega_2^0), \end{aligned} \quad (38)$$

in agreement with result (30). From the Friedel sum rule we get

$$Z_{\text{sc}} = \frac{1}{\pi} \text{Tr } \phi(\mu) = \Theta(\mu - \omega_1) + \Theta(\mu - \omega_2) - \Theta(\mu - \omega_1^0) + \Theta(\mu - \omega_2^0), \quad (39)$$

which agrees with result (32). Therefore, the Friedel sum rule works for two site (matrix) non-interacting system as well. We also note that it is easy to generalize this two site model by perturbing asymmetrically both sites by potentials v_1 and v_2 as is presented next.

As a double check we can use Cauchy formula to determine explicitly the logarithm of a matrix function, i.e.

$$\ln(\mathbf{1} - \mathbf{V}\mathbf{G}_0) = \frac{1}{2\pi i} \int_{\Gamma} \ln\zeta [\zeta\mathbf{1} - \mathbf{1} + \mathbf{V}\mathbf{G}_0]^{-1} d\zeta, \quad (40)$$

where the contour Γ incloses all eigenvalues λ_1 and λ_2 and the function $\ln\zeta$ is analytic on and inside it. Then we compute

$$\begin{aligned} \ln(\mathbf{1} - \mathbf{V}\mathbf{G}_0) &= \\ &= \frac{1}{2\pi i} \int_{\Gamma} \ln\zeta \begin{bmatrix} \zeta - 1 & 0 \\ \frac{vt}{z^2 - t^2} & \zeta - 1 - \frac{vz}{z^2 - t^2} \end{bmatrix}^{-1} d\zeta \\ &= \frac{1}{2\pi i} \int_{\Gamma} \ln\zeta \frac{1}{(\zeta - 1)(\zeta - 1 - \frac{vz}{z^2 - t^2})} \\ &\quad \begin{bmatrix} \zeta - 1 - \frac{vz}{z^2 - t^2} & 0 \\ -\frac{vt}{z^2 - t^2} & \zeta - 1 \end{bmatrix} d\zeta. \end{aligned} \quad (41)$$

Using the theorem of residua we can compute each matrix element independently and obtain

$$\ln(\mathbf{1} - \mathbf{V}\mathbf{G}_0) = \begin{bmatrix} 0 & 0 \\ -\frac{t}{z} \ln(1 + \frac{vt}{z^2 - t^2}) & \ln(1 + \frac{vz}{z^2 - t^2}) \end{bmatrix}, \quad (42)$$

which agrees with the earlier result gives the correct trace as presented above.

3. *Two site model with general potentials* - Consider the model with $\hat{H} = -t|1\rangle\langle 2| - t|2\rangle\langle 1| + v_1|1\rangle\langle 1| + v_2|2\rangle\langle 2|$. Find t-matrix, Green'd function, scattering phase shift, screening charge treating a shift

v of the energy level as perturbation.

The free Green function matrix is

$$\begin{aligned} \mathbf{G}_0(z) &= \begin{pmatrix} z - \epsilon & t \\ t & z - \epsilon \end{pmatrix}^{-1} \\ &= \frac{1}{(z - \epsilon)^2 - t^2} \begin{pmatrix} z - \epsilon & -t \\ -t & z - \epsilon \end{pmatrix}, \end{aligned} \quad (43)$$

where ϵ is the on site energy and $t > 0$ is the hopping amplitude between sites. In the perturbed case both sites are shifted by potentials v_1 and v_2 respectively, so the scattering potential is

$$\mathbf{V} = \begin{pmatrix} v_1 & 0 \\ 0 & v_2 \end{pmatrix}, \quad (44)$$

and the Green function is

$$\begin{aligned} \mathbf{G}(z) &= \begin{pmatrix} z - \epsilon - v_1 & t \\ t & z - \epsilon - v_2 \end{pmatrix}^{-1} \\ &= \frac{1}{(z - \epsilon_1)(z - \epsilon_2) - t^2} \begin{pmatrix} z - \epsilon_2 & -t \\ -t & z - \epsilon_1 \end{pmatrix}, \end{aligned} \quad (45)$$

where we used $\epsilon_1 = \epsilon + v_1$ and $\epsilon_2 = \epsilon + v_2$.

The perturbed Green function \mathbf{G} has singularities at

$$\omega_{1,2} = \frac{1}{2}[\epsilon_1 + \epsilon_2 \pm \sqrt{(\epsilon_1 - \epsilon_2)^2 + 4t^2}], \quad (46)$$

and the free Green function \mathbf{G}_0 is singular at

$$\omega_{1,2}^0 = \epsilon \pm t. \quad (47)$$

To find DOS we need the trace of \mathbf{G} and \mathbf{G}_0 :

$$\begin{aligned} \text{Tr } \mathbf{G}(z) &= \frac{2z - \epsilon_1 - \epsilon_2}{(z - \epsilon_1)(z - \epsilon_2) - t^2} \\ &= \frac{2z - \epsilon_1 - \epsilon_2}{\omega_1 - \omega_2} \left(\frac{1}{z - \omega_1} - \frac{1}{z - \omega_2} \right) \end{aligned} \quad (48)$$

Observing that $\omega_1 - \omega_2 = \sqrt{(\epsilon_1 - \epsilon_2)^2 + 4t^2}$ we obtain

$$\text{Tr } \mathbf{G}(z) = \frac{2z - \epsilon_1 - \epsilon_2}{\sqrt{(\epsilon_1 - \epsilon_2)^2 + 4t^2}} \left(\frac{1}{z - \omega_1} - \frac{1}{z - \omega_2} \right), \quad (49)$$

and in the limit $z \rightarrow \omega + i0^+$

$$\begin{aligned} \text{Tr } \mathbf{G}(\omega) &= \frac{2\omega - \epsilon_1 - \epsilon_2}{\sqrt{(\epsilon_1 - \epsilon_2)^2 + 4t^2}} \left(P \frac{1}{\omega - \omega_1} - i\pi\delta(\omega - \omega_1) \right. \\ &\quad \left. - P \frac{1}{\omega - \omega_2} + i\pi\delta(\omega - \omega_2) \right). \end{aligned} \quad (50)$$

With observation that $2\omega_{1,2} - \epsilon_1 - \epsilon_2 = \pm\sqrt{v^2 + 4t^2}$ the DOS is

$$\begin{aligned} \rho(\omega) &= -\frac{1}{\pi} \text{Im Tr } \mathbf{G}(\omega) \\ &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2). \end{aligned} \quad (51)$$

The unperturbed DOS is

$$\rho(\omega) = \delta(\omega - \omega_1^0) + \delta(\omega - \omega_2^0), \quad (52)$$

and the change of DOS is

$$\begin{aligned} \Delta\rho(\omega) &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2) \\ &\quad - \delta(\omega - \omega_1^0) - \delta(\omega - \omega_2^0). \end{aligned} \quad (53)$$

The screening charge is

$$\begin{aligned} Z_{sc} &= \int d\omega \Delta\rho(\omega) f(\omega) = \frac{1}{e^{\beta(\omega_1 - \mu)} + 1} + \frac{1}{e^{\beta(\omega_2 - \mu)} + 1} \\ &\quad - \frac{1}{e^{\beta(\omega_1^0 - \mu)} + 1} - \frac{1}{e^{\beta(\omega_2^0 - \mu)} + 1} \end{aligned} \quad (54)$$

and at $T = 0$ reduces to

$$\begin{aligned} Z_{sc} &= \Theta(\mu - \omega_1) + \Theta(\mu - \omega_2) \\ &\quad - \Theta(\mu - \omega_1^0) - \Theta(\mu - \omega_2^0). \end{aligned} \quad (55)$$

Next we use scattering to determine the change of DOS and the screening charge. From the main text we have

$$\begin{aligned} \Delta\rho(\omega) &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{V}^{-1}(z)\mathbf{T}(z)] \\ &= \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{1} - \mathbf{V}\mathbf{G}_0]^{-1} \\ &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im Tr ln}[\mathbf{1} - \mathbf{V}\mathbf{G}_0]. \end{aligned} \quad (56)$$

The matrix

$$\begin{aligned} \mathbf{1} - \mathbf{V}\mathbf{G}_0 &= \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \\ - \frac{1}{(z - \epsilon)^2 - t^2} \begin{pmatrix} v_1 & 0 \\ 0 & v_2 \end{pmatrix} \begin{pmatrix} z - \epsilon & -t \\ -t & z - \epsilon \end{pmatrix} &= \\ = \begin{pmatrix} 1 - \frac{(z - \epsilon)v_1}{(z - \epsilon)^2 - t^2} & \frac{v_1 t}{(z - \epsilon)^2 - t^2} \\ \frac{v_2 t}{(z - \epsilon)^2 - t^2} & 1 - \frac{(z - \epsilon)v_2}{(z - \epsilon)^2 - t^2} \end{pmatrix} \end{aligned} \quad (57)$$

needs to be diagonalized. There are two eigenvalues

$$\begin{aligned} \lambda_{\pm} &= \frac{1}{2} \left[2 - \frac{(v_1 + v_2)(z - \epsilon)}{(z - \epsilon)^2 - t^2} \right. \\ &\quad \left. \pm \sqrt{\frac{(v_1 - v_2)^2(z - \epsilon)^2 + 4v_1v_2t^2}{((z - \epsilon)^2 - t^2)^2}} \right] \end{aligned} \quad (58)$$

and the logarithm of the diagonal matrix is a matrix of logarithms of its eigenvalues. Since the trace is base independent we have

$$\begin{aligned} \Delta\rho(\omega) &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} (\ln\lambda_+ + \ln\lambda_-) \\ &= - \lim_{z \rightarrow \omega + i0^+} \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \ln(\lambda_+\lambda_-). \end{aligned} \quad (59)$$

After some algebraic manipulation one can easily show that

$$\begin{aligned} \lambda_+\lambda_- &= \frac{(z - \epsilon)^2 - (v_1 + v_2)(z - \epsilon) + v_1v_2 - t^2}{(z - \epsilon)^2 - t^2} \\ &= \frac{(z - \omega_1)(z - \omega_2)}{(z - \omega_1^0)(z - \omega_2^0)}. \end{aligned} \quad (60)$$

Therefore,

$$\begin{aligned} &\lim_{z \rightarrow \omega + i0^+} \ln(\lambda_+\lambda_-) \\ &= \ln|(\omega - \omega_1)(\omega - \omega_2)| + i\pi(\Theta(\omega_1 - \omega) + \Theta(\omega_1 - \omega)) \\ &\quad - \ln|(\omega - \omega_1^0)(\omega - \omega_2^0)| - i\pi(\Theta(\omega_1^0 - \omega) + \Theta(\omega_1^0 - \omega)). \end{aligned} \quad (61)$$

Hence, we find a nonzero scattering phase shift

$$\begin{aligned} \text{Tr } \phi(\omega) &= \pi[\Theta(\omega - \omega_1) + \Theta(\omega - \omega_2) \\ &\quad - \Theta(\omega - \omega_1^0) + \Theta(\omega - \omega_2^0)]. \end{aligned} \quad (62)$$

The change of DOS

$$\begin{aligned} \Delta\rho(\omega) &= \frac{1}{\pi} \frac{d}{d\omega} \pi[\Theta(\omega - \omega_1) + \Theta(\omega - \omega_2) \\ &\quad - \Theta(\omega - \omega_1^0) + \Theta(\omega - \omega_2^0)] \\ &= \delta(\omega - \omega_1) + \delta(\omega - \omega_2) \\ &\quad - \delta(\omega - \omega_1^0) - \delta(\omega - \omega_2^0), \end{aligned} \quad (63)$$

in agreement with result (53). From the Friedel sum rule we get

$$\begin{aligned} Z_{sc} &= \frac{1}{\pi} \text{Tr } \phi(\mu) = \Theta(\mu - \omega_1) + \Theta(\mu - \omega_2) \\ &\quad - \Theta(\mu - \omega_1^0) + \Theta(\mu - \omega_2^0), \end{aligned} \quad (64)$$

which agrees with result (55). Therefore, the Friedel sum rule works for general two-site (matrix) non-interacting systems as well.

16 Week XVI, 13-14/06/2022

16.1 Lecture

&8. *Resonances, resonant states* - experimental evidence of resonance states in the cross-section, intuitive relation with quasi-bound states with a finite life-time, classification of possible solutions of the Schrödinger equation and their location on the complex energy plane and complex momentum plane: i) continuum states, ii) bound states, iii) virtual states, iv) resonant states, v) antiresonance states, two Riemann sheets of the complex energy plane and the locations of the different solutions, properties of the resonances in space and in time, Bright-Wigner formula for resonances, mathematical origin for complex eigen-energies of resonances, difference between hermitian and self-adjoint operators, purely outgoing boundary condition as a source of non-self-adjointness.

16.2 Tutorial

1. *δ-Dirac potential* - Consider particles with mass m moving in one dimension. The potential is $V(x) = -V_0\delta(x)$, where $\delta(x)$ is a Dirac distribution. Solve this problem for bound states $E < 0$ and scattered states $E > 0$ within the wave function approach with proper boundary conditions at $x = 0$. Discuss results. Next, for scattered states $E > 0$ solve this problem within: a) t-matrix formalism, b) Lippmann-Schwinger formalism.
2. *Anderson-Fano-Friedrichs-Lee resonant model* - Consider the AFFL model

$$\hat{H} = \epsilon_0|0\rangle\langle 0| + \int_R dk \epsilon_k |k\rangle\langle k| + \int dk_R V_k |0\rangle\langle k| + V_k^* |k\rangle\langle 0|.$$

Compute Green functions and find t-matrix. Discuss analytic structure of the t-matrix and its poles and a cut on both Riemann's sheets. For an explicit discussion assume that the DOS is constant $1/2D$ between $-D$ and D and zero otherwise and that $V_k = V_k^* = V$.

16.3 Homework problems

1. *Double δ-potential* - Consider a system in one dimension with the potential

$$V(x) = V_0[\delta(x+l) + \delta(x-l)],$$

with $V_0 > 0$. Assuming the purely outgoing boundary conditions, i.e. taking an ansatz for the solution

$$\psi_{\text{res}}(x) = \begin{cases} B e^{-ikx} & \text{for } x < -l \\ F e^{ikx} + G e^{-ikx} & \text{for } -l < x < l \\ C e^{ikx} & \text{for } l < x, \end{cases}$$

find equation(s) on k_n and energies $E_n = \hbar^2 k_n^2 / 2m$. By solving them numerically show that the complex solutions are admitted $E_n = \epsilon_n^r - i\Gamma_n/2$. Find them numerically, e.g. with Mathematica. These are resonances. Hint: arxiv:0705.1388.

2. *Double δ-potential - transition and reflection coefficients* - Consider again a system in one dimension with the potential

$$V(x) = V_0[\delta(x+l) + \delta(x-l)],$$

with $V_0 > 0$. Assuming the boundary conditions with incoming wave from the left, i.e. taking an ansatz for the solution

$$\psi_{\text{res}}(x) = \begin{cases} A e^{ikx} + B e^{-ikx} & \text{for } x < -l \\ F e^{ikx} + G e^{-ikx} & \text{for } -l < x < l \\ C e^{ikx} & \text{for } l < x, \end{cases}$$

derive transition $t(k) = C/A$ and reflection $r(k) = B/A$ coefficients. Show, that their divergence, i.e. when $A = 0$ (as in purely outgoing boundary condition, gives the same condition on k as in the former problem. Plot, e.g. with Mathematica, $t(E)$ for real $E = \hbar^2 k^2 / 2m > 0$ and check if positions of maxima correlate with ϵ_n^r . Hint: arxiv:0705.1388.

17 Literature

- W. Greiner, B. Müller *Quantum mechanics - symmetries*.
- L.E. Ballentine, *Quantum mechanics. A modern development*.
- A. Messiah, *Quantum mechanics*, vol. I and II.
- J.J. Sakurai, J. Napolitano, *Modern quantum mechanics*.
- L. I. Schiff, *Quantum mechanics*.
- A. Altland, http://www.thp.uni-koeln.de/Documents/altland_advqm_2012.pdf
- More to be added in the course.