

Exactly solvable Schrödinger operators related to the hypergeometric equation

Jan Dereziński*

Department of Mathematical Methods in Physics,
Faculty of Physics, University of Warsaw,
Pasteura 5, 02-093 Warszawa, Poland
email: jan.dereziński@fuw.edu.pl

Pedram Karimi

Faculty of Physics, University of Warsaw,
Pasteura 5, 02-093 Warszawa, Poland
email: pedram.karimi@fuw.edu.pl

March 6, 2026

Abstract

We study one-dimensional Schrödinger operators defined as closed operators that are exactly solvable in terms of the Gauss hypergeometric function. We allow the potentials to be complex. These operators fall into three groups. The first group can be reduced to the Gegenbauer equation, up to an affine transformation, a special case of the hypergeometric equation. The two other groups, which we call *hypergeometric of the first*, resp. *second kind*, can be reduced to the general Gauss hypergeometric equation. Each of the group is subdivided in three families, acting on the Hilbert space $L^2] - 1, 1[$, $L^2(\mathbb{R}_+)$ resp. $L^2(\mathbb{R})$. Motivated by geometric applications of these families, we call them *spherical*, *hyperbolic*, resp. *deSitterian*. All these families are known from applications in Quantum Mechanics: e.g. spherical hypergeometric Schrödinger operators of the first kind are often called *trigonometric Pöschl-Teller Hamiltonians*. For operators belonging to each family we compute their spectrum and determine their Green function (the integral kernel of their resolvent). We also describe transmutation identities that relate these Green functions. These identities interchange spectral parameters with coupling constants across different operator families. Finally, we describe how these operators arise from separation of variables of (pseudo-)Laplacians on symmetric manifolds. Our paper can be viewed as a sequel to [DL], where closed realizations of one-dimensional Schrödinger operators solvable in terms Kummer's confluent equation were studied.

Mathematics Subject Classification MSC2020: 33C05; 34L40; 47A10; 81Q80

Contents

1	Introduction	3
1.1	3×3 families of hypergeometric Hamiltonians	3
1.2	Review of 9 families	4
1.3	Transmutations of Green functions	6
1.4	Geometric applications	8
1.5	Boundary conditions	9
1.6	Comparison with literature and historic remarks	10
1.7	Plan of the paper	12

2	Preliminaries	13
2.1	Hypergeometric equation	13
2.2	Gegenbauer equation	14
2.3	Liouville transformation	14
3	Gegenbauer Hamiltonians	15
3.1	Spherical case	16
3.2	Hyperbolic case	17
3.3	DeSitterian case	19
4	Hypergeometric Hamiltonians of the first kind	21
4.1	Spherical case	22
4.2	Hyperbolic case	24
4.3	DeSitterian case	26
5	Hypergeometric Hamiltonians of the second kind	30
5.1	Spherical case	31
5.2	Hyperbolic case	34
5.3	DeSitterian case	36
6	The Laplacian on an interval, halfline and line	38
6.1	Laplacian on an interval	39
6.2	Laplacian on the halfline	40
6.3	Laplacian on the line	41
7	Geometric applications	42
7.1	Sphere	42
7.2	Hyperbolic space	43
7.3	DeSitter space	44
7.4	Sphere in double spherical coordinates	44
7.5	Hyperboloid in double spherical coordinates	45
7.6	Complex manifolds	46
7.7	Hyperboloid $\mathbb{H}^{p-1,p}$ in complex coordinates	47
A	Holomorphic families of operators	48
B	Riemann equation	49
B.1	Regular-singular points	49
B.2	Riemann equation	49
B.3	From Riemann equation to hypergeometric equation	50
C	Identities for the hypergeometric function	51
C.1	Kummer's table	51
C.2	Connection formulas	52
C.3	Degenerate case	53
C.4	Gegenbauer functions	53
C.5	Whipple transformation	54
C.6	Half integer case	54

D	Closed realizations of 1d Schrödinger operators	55
D.1	Minimal and maximal realization	55
D.2	Resolvent	56
E	Associated Legendre functions vs. Gegenbauer functions	56

1 Introduction

One-dimensional Schrödinger operators are operators of the form

$$L := -\partial_x^2 + V(x), \tag{1.1}$$

where $V(x)$ is the *potential*, which in this paper is allowed to be complex-valued. Our paper is devoted to several families of operators of the form (1.1), interpreted as *closed operators* on $L^2(]a, b[)$ for appropriate $-\infty \leq a < b \leq +\infty$, which can be reduced to the (*Gauss*) *hypergeometric equation*

$$(z(1-z)\partial_z^2 + (c - (a+b+1)z)\partial_z - ab)f(z) = 0, \tag{1.2}$$

and whose Green functions can be expressed in terms of the (*Gauss*) *hypergeometric function*.

We will also consider operators of the form (1.1) that can be reduced to the *Gegenbauer equation*.

$$\left((1-w^2)\partial_w^2 - 2(1+\alpha)w\partial_w + \lambda^2 - \left(\alpha + \frac{1}{2}\right)^2 \right) g(w) = 0. \tag{1.3}$$

The Gegenbauer equation is up to an affine transformation a special case of the hypergeometric equation. Its special property is the mirror symmetry.

Our paper can be viewed as the sequel to [DL], where one of the authors (JD) together with Jinyeop Lee studied a similar problem for the *confluent equation*. We will mostly use the same terminology and methods. We try to make the present paper reasonably self-contained, however the reader is encouraged to consult [DL], especially concerning the general theory of closed realizations of operators of the form (1.1).

In the remaining part of the introduction we give a summary of the results of our paper. In the later section these results will be discussed in detail.

1.1 3×3 families of hypergeometric Hamiltonians

Abusing the terminology, for the sake of brevity, we will use the term *Hamiltonian* for one-dimensional Schrödinger operators. We study three categories of Hamiltonians:

- (1) Those reducible to the Gegenbauer equation; they will depend on a single complex parameter, and can be viewed as a subclass of hypergeometric Hamiltonians, both of the first and second kind.
- (2) Those reducible to the hypergeometric equation by the substitution $z = \sin^2 \frac{r}{2}$ (or similar); they will be called *hypergeometric of the first kind*; they will depend on two complex parameters.
- (3) Those reducible to the hypergeometric equation by the substitution $z = \frac{1}{1+e^{2r}}$ (or similar); they will be called *hypergeometric of the second kind*; they will depend on two complex parameters.

Within each category we will consider 3 families, which differ by the choice of the interval $]a, b[$. This interval can be viewed as a subset of the complex plane. We will always assume that the endpoints are singular points of the equation. For each of $3 \times 3 = 9$ cases, for a set of parameters with a nonempty interior the operator L , defined originally on $C_c^\infty]a, b[$, possesses a unique closed realization in the sense of $L^2]a, b[$. This realization depends holomorphically on parameters, and extends to a holomorphic family of

closed operators on a larger domain. We will call it the *basic family* of closed realizations of L . For some special ranges of parameters there exist other closed realizations of L with *mixed boundary conditions*—we will not consider them in this paper.

For each operator L_\bullet from those families we will find its spectrum, denoted $\sigma(L_\bullet)$. In all cases with real potentials, these operators will be self-adjoint (so that $\sigma(L_\bullet) \subset \mathbb{R}$). More generally, the resolvent set (the complement of the spectrum) of these operators will be nonempty. For z in the resolvent set we will find the resolvent, that is $(L_\bullet - z)^{-1}$, which we will usually denote $\frac{1}{L_\bullet - z}$. The *Green function of $L_\bullet - z$* , that is integral kernel of the resolvent $\frac{1}{L_\bullet - z}$, will be denoted $\frac{1}{(L_\bullet - z)}(x, y)$, with $x, y \in]a, b[$. We will find expressions of Green functions of Hamiltonians from all 3×3 families in terms of the Gamma function and the Gauss hypergeometric function.

These 9 families were discovered in the early days of Quantum Mechanics by physicists trying to find exactly solvable models for various quantum systems. In the literature they are usually named after the researchers who discovered them. Instead of the traditional names, we will prefer to use different names: spherical, hyperbolic and deSitterian. In the spherical case $]a, b[$ is $] - 1, 1[$, in the hyperbolic case it is $]0, +\infty[=: \mathbb{R}_+$, and in the deSitterian case it is \mathbb{R} .

Our names indicate a major geometric application of these families: spherical, hyperbolic, resp. deSitterian Gegenbauer Hamiltonians appear when we separate variables for the (pseudo-)Laplacian on the sphere, on the hyperbolic space, resp. on the deSitter space.

1.2 Review of 9 families

Let us briefly review the 9 families described in the paper, referring the reader to the main text for precise statements. We will write \mathcal{A}_{un} for the domain of uniqueness, that is, the set of parameters (a subset of \mathbb{C} or $\mathbb{C} \times \mathbb{C}$) for which there exists a unique closed realization of a given differential expression. This realization in all cases depends holomorphically on its parameters, and extends to a larger domain, denoted \mathcal{A}_{hol} . \mathcal{A}_{sa} will indicate the set of parameters for which the operator is self-adjoint. Note that the operators are essentially self-adjoint on $C_c^\infty]a, b[$ if and only if the parameter belongs to $\mathcal{A}_{\text{sa}} \cap \mathcal{A}_{\text{un}}$.

Note that we always use the notion of an analytic family of closed operators in the sense of Kato [Kato, DW1]. In Appendix A we give a concise explanation of this concept.

1. Gegenbauer Hamiltonians

(a) Spherical Gegenbauer Hamiltonian, $L^2]0, \pi[$:

$$L_\alpha^s := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\sin^2 r}, \quad (1.4)$$

$$\mathcal{A}_{\text{un}} = \{\text{Re } \alpha \geq 1\}, \quad \mathcal{A}_{\text{hol}} = \{\text{Re } \alpha > -1\}, \quad \mathcal{A}_{\text{sa}} =] - 1, +\infty[.$$

(b) Hyperbolic Gegenbauer Hamiltonian, $L^2(\mathbb{R}_+)$:

$$L_\alpha^h := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\sinh^2 r}, \quad (1.5)$$

$$\mathcal{A}_{\text{un}} = \{\text{Re } \alpha \geq 1\}, \quad \mathcal{A}_{\text{hol}} = \{\text{Re } \alpha > -1\}, \quad \mathcal{A}_{\text{sa}} =] - 1, +\infty[.$$

(c) DeSitterian Gegenbauer Hamiltonian, $L^2(\mathbb{R})$:

$$L_\alpha^{\text{dS}} := -\partial_r^2 - \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r}, \quad (1.6)$$

$$\mathcal{A}_{\text{un}} = \mathcal{A}_{\text{hol}} = \mathbb{C}, \quad \mathcal{A}_{\text{sa}} = \mathbb{R}.$$

2. Hypergeometric Hamiltonians of the first kind:

(a) Spherical hypergeometric Hamiltonian of the first kind, $L^2]0, \pi[$:

$$L_{\alpha,\beta}^s := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{4 \sin^2 \frac{r}{2}} + \left(\beta^2 - \frac{1}{4}\right) \frac{1}{4 \cos^2 \frac{r}{2}}, \quad (1.7)$$

$$\mathcal{A}_{\text{un}} = \{\text{Re } \alpha, \text{Re } \beta \geq 1\}, \quad \mathcal{A}_{\text{hol}} = \{\text{Re } \alpha, \text{Re } \beta > -1\}, \quad \mathcal{A}_{\text{sa}} =]-1, +\infty[\times]-1, +\infty[.$$

(b) Hyperbolic hypergeometric Hamiltonian of the first kind, $L^2(]0, \infty[)$:

$$L_{\alpha,\beta}^h := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{4 \sinh^2 \frac{r}{2}} - \left(\beta^2 - \frac{1}{4}\right) \frac{1}{4 \cosh^2 \frac{r}{2}}, \quad (1.8)$$

$$\mathcal{A}_{\text{un}} = \{\text{Re } \alpha \geq 1\} \times \mathbb{C}, \quad \mathcal{A}_{\text{hol}} = \{\text{Re } \alpha > -1\} \times \mathbb{C}, \quad \mathcal{A}_{\text{sa}} =]-1, +\infty[\times \mathbb{R}.$$

(c) DeSitterian hypergeometric Hamiltonian of the first kind, $L^2(\mathbb{R})$,

$$L_{\alpha,\beta}^{\text{dS}} := -\partial_r^2 - \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r} \left(\frac{1}{2} + \frac{i \sinh r}{2}\right) - \left(\beta^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r} \left(\frac{1}{2} - \frac{i \sinh r}{2}\right), \quad (1.9)$$

$$\mathcal{A}_{\text{un}} = \mathcal{A}_{\text{hol}} = \mathbb{C} \times \mathbb{C}, \quad \mathcal{A}_{\text{sa}} = \{\alpha = \bar{\beta}\}.$$

3. Hypergeometric Hamiltonians of the second kind:

(a) Spherical hypergeometric Hamiltonian of the second kind, $L^2]0, \pi[$:

$$K_{\tau,\mu}^s := -\partial_u^2 + \tau \frac{\cos u}{\sin u} + \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\sin^2 u},$$

$$\mathcal{A}_{\text{un}} = \mathbb{C} \times \{\text{Re } \mu \geq 2\}, \quad \mathcal{A}_{\text{hol}} = \mathbb{C} \times \{\text{Re } \mu > -2\} \setminus \{(0, -1)\}, \quad \mathcal{A}_{\text{sa}} = \mathbb{R} \times]-2, +\infty[. \quad (1.10)$$

(b) Hyperbolic hypergeometric Hamiltonian of the second kind, $L^2(]0, \infty[)$:

$$K_{\kappa,\mu}^h := -\partial_u^2 + \kappa \frac{\cosh u}{\sinh u} + \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\sinh^2 u},$$

$$\mathcal{A}_{\text{un}} = \mathbb{C} \times \{\text{Re } \mu \geq 2\}, \quad \mathcal{A}_{\text{hol}} = \mathbb{C} \times \{\text{Re } \mu > -2\} \setminus \{(0, -1)\}, \quad \mathcal{A}_{\text{sa}} = \mathbb{R} \times]-2, +\infty[. \quad (1.11)$$

(c) DeSitterian hypergeometric Hamiltonian of the second kind, $L^2(\mathbb{R})$,

$$K_{\kappa,\mu}^{\text{dS}} := -\partial_w^2 - \kappa \frac{\sinh w}{\cosh w} - \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\cosh^2 w}, \quad (1.12)$$

$$\mathcal{A}_{\text{un}} = \mathcal{A}_{\text{hol}} = \mathbb{C} \times \mathbb{C}, \quad \mathcal{A}_{\text{sa}} = \mathbb{R} \times \mathbb{R}.$$

Gegenbauer Hamiltonians are special cases of hypergeometric Hamiltonians of both the first and second type. In fact, we have the following coincidences:

$$L_{\alpha}^s = L_{\alpha,\alpha}^s = K_{0,2\alpha}^s; \quad (1.13)$$

$$L_{\alpha}^h = L_{\alpha,\alpha}^h = K_{0,2\alpha}^h; \quad (1.14)$$

$$L_{\alpha}^{\text{dS}} = L_{\alpha,\alpha}^{\text{dS}} = K_{0,2\alpha}^{\text{dS}}. \quad (1.15)$$

Let us also list the following identities that we prove:

$$K_{\tau,-1}^s = K_{\tau,1}^s, \quad \tau \neq 0, \quad (1.16)$$

$$K_{\kappa,-1}^h = K_{\kappa,1}^h, \quad \kappa \neq 0. \quad (1.17)$$

These identities imply that $(0, -1)$ are singularities of the functions $(\tau, \mu) \mapsto K_{\tau,\mu}^s$ and $(\kappa, \mu) \mapsto K_{\kappa,\mu}^h$.

Going back to (1.13) and (1.14), note that

$$L_\alpha^s = K_{0,2\alpha}^s, \quad (1.18)$$

$$L_\alpha^h = K_{0,2\alpha}^h, \quad (1.19)$$

are the identities for holomorphic functions only for $\alpha \neq -\frac{1}{2}$, because of the above mentioned singularity. Thus the identities

$$L_{-\frac{1}{2}}^s = K_{0,-1}^s, \quad (1.20)$$

$$L_{-\frac{1}{2}}^h = K_{0,-1}^h. \quad (1.21)$$

should be used as *definitions* of $K_{0,-1}^s$ and $K_{0,-1}^h$.

1.3 Transmutations of Green functions

Green functions of distinct Hamiltonians from the above list are linked by identities, which we find quite curious. We call them *transmutation identities*, since the spectral parameter undergoes a change into a coupling constant and the other way around. They follow from various identities satisfied by the hypergeometric function and are similar to the transmutation identities for Hamiltonians related to the confluent equation described in [DL].

In the literature the name “transmutation” is sometimes used for identities of the form $A = VBV^{-1}$, e.g. in [Car, SS]. Note that our transmutation identities are not of this form, and in fact they seem to represent a novel contribution.

Here is the list of transmutation identities considered in our paper. In each case, first we indicate the change of variables involved in a given transmutation. Then we describe two versions of the identity for Green functions.

Proposition 1.1. Gegenbauer spherical — Gegenbauer deSitterian:

$$]0, \pi[\ni r \mapsto q \in \mathbb{R}, \quad \cot r = \sinh q; \quad (1.22)$$

$$\sin^{\frac{1}{2}} r \frac{1}{(L_\lambda^s - \alpha^2)}(r, r') \sin^{\frac{1}{2}} r' = \frac{1}{(L_\alpha^{\text{dS}} - \lambda^2)}(q, q'), \quad (1.23)$$

$$\frac{1}{(L_\lambda^s - \alpha^2)}(r, r') = \cosh^{\frac{1}{2}} q \frac{1}{(L_\alpha^{\text{dS}} - \lambda^2)}(q, q') \cosh^{\frac{1}{2}} q'. \quad (1.24)$$

Proposition 1.2. 1st kind spherical — 1st kind hyperbolic:

$$]0, \pi[\ni r \mapsto q \in \mathbb{R}_+, \quad \tan \frac{r}{2} = \sinh \frac{q}{2}; \quad (1.25)$$

$$\left(\cos \frac{r}{2}\right)^{-\frac{1}{2}} \frac{1}{\left(L_{\alpha,\beta}^s + \frac{\mu^2}{4}\right)}(r, r') \left(\cos \frac{r'}{2}\right)^{-\frac{1}{2}} = \frac{1}{\left(L_{\alpha,\mu}^h + \frac{\beta^2}{4}\right)}(q, q'), \quad (1.26)$$

$$\frac{1}{\left(L_{\alpha,\beta}^s + \frac{\mu^2}{4}\right)}(r, r') = \left(\cosh \frac{q}{2}\right)^{-\frac{1}{2}} \frac{1}{\left(L_{\alpha,\mu}^h + \frac{\beta^2}{4}\right)}(q, q') \left(\cosh \frac{q'}{2}\right)^{-\frac{1}{2}}. \quad (1.27)$$

Proposition 1.3. 2nd kind hyperbolic — 2nd kind deSitterian

$$\mathbb{R}_+ \ni u \mapsto w \in \mathbb{R}, \quad e^{2u} = 1 + e^{2w}; \quad (1.28)$$

$$(1 - e^{-2u})^{-\frac{1}{2}} \frac{1}{\left(K_{\nu - \frac{\beta^2}{2}, \mu}^h + \nu + \frac{\beta^2}{2}\right)}(u, u')(1 - e^{-2u'})^{-\frac{1}{2}} = \frac{1}{\left(K_{\nu + \frac{\mu^2}{2}, \beta}^{\text{dS}} + \nu - \frac{\mu^2}{2}\right)}(w, w'), \quad (1.29)$$

$$\frac{1}{\left(K_{\nu - \frac{\beta^2}{2}, \mu}^h + \nu + \frac{\beta^2}{2}\right)}(u, u') = (1 + e^{-2w})^{-\frac{1}{2}} \frac{1}{\left(K_{\nu + \frac{\mu^2}{2}, \beta}^{\text{dS}} + \nu - \frac{\mu^2}{2}\right)}(w, w') (1 + e^{-2w'})^{-\frac{1}{2}}. \quad (1.30)$$

Proposition 1.4. 1st kind spherical — 2nd kind deSitterian

$$]0, \pi[\ni r \mapsto u \in \mathbb{R}, \quad \cos r = \tanh u; \quad (1.31)$$

$$\sin^{\frac{1}{2}} r \frac{1}{\left(L_{\alpha, \beta}^s + \frac{\mu^2}{4}\right)}(r, r') \sin^{\frac{1}{2}} r' = \frac{1}{\left(K_{\kappa, \mu}^{\text{dS}} + \delta\right)}(u, u'), \quad (1.32)$$

$$\frac{1}{\left(L_{\alpha, \beta}^s + \frac{\mu^2}{4}\right)}(r, r') = \cosh^{\frac{1}{2}} u \frac{1}{\left(K_{\kappa, \mu}^{\text{dS}} + \delta\right)}(u, u') \cosh^{\frac{1}{2}} u', \quad (1.33)$$

$$\delta = \frac{\alpha^2 + \beta^2}{2}, \quad \kappa = \frac{\alpha^2 - \beta^2}{2}. \quad (1.34)$$

Proposition 1.5. 1st kind hyperbolic — 2nd kind hyperbolic

$$\mathbb{R}_+ \ni r \mapsto u \in \mathbb{R}_+, \quad \cosh r = \coth u; \quad (1.35)$$

$$\sinh^{\frac{1}{2}} r \frac{1}{\left(L_{\alpha, \beta}^h + \frac{\mu^2}{4}\right)}(r, r') \sinh^{\frac{1}{2}} r' = \frac{1}{\left(K_{\kappa, \mu}^h + \delta\right)}(u, u'), \quad (1.36)$$

$$\frac{1}{\left(L_{\alpha, \beta}^h + \frac{\mu^2}{4}\right)}(r, r') = \sinh^{\frac{1}{2}} u \frac{1}{\left(K_{\kappa, \mu}^h + \delta\right)}(u, u') \sinh^{\frac{1}{2}} u', \quad (1.37)$$

$$\delta = \frac{\alpha^2 + \beta^2}{2}, \quad \kappa = \frac{\alpha^2 - \beta^2}{2}. \quad (1.38)$$

Proposition 1.6. 1st kind deSitterian — 2nd kind spherical

$$\mathbb{R} \ni r \mapsto u \in]0, \pi[, \quad \sinh r = -\cot u; \quad (1.39)$$

$$\cosh^{\frac{1}{2}} r \frac{1}{\left(L_{\alpha, \beta}^{\text{dS}} + \frac{\mu^2}{4}\right)}(r, r') \cosh^{\frac{1}{2}} r' = \frac{1}{\left(K_{\tau, \mu}^s + \delta\right)}(u, u'), \quad (1.40)$$

$$\frac{1}{\left(L_{\alpha, \beta}^{\text{dS}} + \frac{\mu^2}{4}\right)}(r, r') = \sin^{\frac{1}{2}} u \frac{1}{\left(K_{\tau, \mu}^s + \delta\right)}(u, u') \sin^{\frac{1}{2}} u', \quad (1.41)$$

$$\delta = \frac{\alpha^2 + \beta^2}{2}, \quad \tau = \frac{i(\alpha^2 - \beta^2)}{2}. \quad (1.42)$$

1.4 Geometric applications

The main original application of hypergeometric Hamiltonians was Quantum Mechanics, as we describe in Subsection 1.6. However, probably the most important context where hypergeometric Hamiltonians appear is geometry, more precisely, the theory of symmetric spaces and Lie groups. In fact, when we separate variables for invariant differential operators, e.g. (pseudo-)Laplacians, on symmetric (pseudo-)Riemannian spaces we often obtain some forms of hypergeometric Hamiltonians.

This fact plays an important role in Quantum Field Theory on curved spacetimes, where Green functions of the d'Alembertian on deSitter and anti-deSitter spaces appear naturally; see e.g. [DeGa].

This geometric interpretation is especially striking for Gegenbauer Hamiltonians. In the following list we show how they arise after separation of variables of various d -dimensional (pseudo-)Laplacians and restriction to $d - 1$ -dimensional spherical harmonics of degree l . In all cases $\alpha = \frac{d}{2} - 1 + l$:

1. Δ_d^s , Laplacian on unit sphere \mathbb{S}^d , reduces to spherical Gegenbauer Hamiltonian L_α^s :

$$(\sin r)^{\frac{d-1}{2}} (-\Delta_d^s)(\sin r)^{-\frac{d-1}{2}} + \left(\frac{d-1}{2}\right)^2 = -\partial_r^2 + \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\sin^2 r}. \quad (1.43)$$

2. Δ_d^h , Laplacian on hyperbolic space \mathbb{H}^d , reduces to hyperbolic Gegenbauer Hamiltonian L_α^h :

$$(\sinh r)^{\frac{d-1}{2}} (-\Delta_d^h)(\sinh r)^{-\frac{d-1}{2}} - \left(\frac{d-1}{2}\right)^2 = -\partial_r^2 + \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\sinh^2 r}. \quad (1.44)$$

3. \square_d^{dS} , d'Alembertian on de Sitter space dS^d , reduces to deSitterian Gegenbauer Hamiltonian L_α^{dS} :

$$(\cosh r)^{\frac{d-1}{2}} \square_d^{\text{dS}}(\cosh r)^{-\frac{d-1}{2}} - \left(\frac{d-1}{2}\right)^2 = -\partial_r^2 - \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\cosh^2 r}. \quad (1.45)$$

All three types of hypergeometric Hamiltonians of the 1st kind have natural geometric interpretations as well. In the following list we show how they arise from separation of variables in a (pseudo-)Laplacian on a (pseudo-)sphere in “double spherical coordinates”. In all three examples the coordinates in the ambient pseudo-Euclidean space are partitioned into two groups.

In the first two cases, these groups are of dimension p and q , and then spherical coordinates are considered within each group. After restriction to products of spherical harmonics of degree l and j a hypergeometric Hamiltonian arises, expressed in the relative variable. We have $\alpha = \frac{p}{2} - 1 + l$, $\beta = \frac{q}{2} - 1 + j$.

The third case is somewhat different: $p = q$ is the dimension of holomorphic and antiholomorphic (complex) spherical coordinates. The spherical harmonics are not the usual ones: they are holomorphic and antiholomorphic harmonics on the complex $p - 1$ -dimensional sphere.

1. Δ_{p+q-1}^s , Laplacian on unit sphere \mathbb{S}^{p+q-1} , reduces to spherical hypergeometric Hamiltonian of 1st kind $L_{\alpha,\beta}^s$:

$$\begin{aligned} & \left(\sin \frac{r}{2}\right)^{\frac{p-1}{2}} \left(\cos \frac{r}{2}\right)^{\frac{q-1}{2}} (-\Delta_{p+q-1}^s) \left(\sin \frac{r}{2}\right)^{-\frac{p-1}{2}} \left(\cos \frac{r}{2}\right)^{-\frac{q-1}{2}} + \left(\frac{p+q-2}{2}\right)^2 \\ &= 4 \left(-\partial_r^2 + \frac{\left(\frac{p-2}{2}\right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{4 \sin^2 \frac{r}{2}} + \frac{\left(\frac{q-2}{2}\right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{4 \cos^2 \frac{r}{2}} \right). \end{aligned} \quad (1.46)$$

2. $\Delta_{p-1,q}$, pseudo-Laplacian on the hyperboloid $\mathbb{H}^{p-1,q}$, reduces to hyperbolic hypergeometric Hamiltonian of 1st kind $L_{\alpha,\beta}^h$:

$$\begin{aligned} & \left(\cosh \frac{r}{2} \right)^{\frac{p-1}{2}} \left(\sinh \frac{r}{2} \right)^{\frac{q-1}{2}} (-\Delta_{p-1,q}) \left(\cosh \frac{r}{2} \right)^{-\frac{p-1}{2}} \left(\sinh \frac{r}{2} \right)^{-\frac{q-1}{2}} - \left(\frac{p+q-2}{2} \right)^2 \\ & = 4 \left(-\partial_r^2 - \frac{\left(\frac{p-2}{2} \right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{4 \cosh^2 \frac{r}{2}} + \frac{\left(\frac{q-2}{2} \right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{4 \sinh^2 \frac{r}{2}} \right). \end{aligned} \quad (1.47)$$

3. $\Delta_{p-1,p}$, pseudo-Laplacian on the hyperboloid $\mathbb{H}^{p-1,p}$, reduces to deSitterian hypergeometric Hamiltonian of 1st kind $L_{\alpha,\beta}^{dS}$:

$$\begin{aligned} & (\cosh r)^{\frac{p-1}{2}} (\Delta_{p-1,p}) (\cosh r)^{-\frac{p-1}{2}} - (p-1)^2 \\ & = 4 \left(-\partial_r^2 - \frac{\left(-\Delta_{p-1}^{s,\mathbb{C}} + \left(\frac{p-1}{2} \right)^2 - \frac{1}{4} \right)}{2(1 + i \sinh r)} - \frac{\left(-\Delta_{p-1}^{s,\mathbb{C}} + \left(\frac{p-1}{2} \right)^2 - \frac{1}{4} \right)}{2(1 - i \sinh r)} \right). \end{aligned} \quad (1.48)$$

Unfortunately, we have not found a direct geometric interpretation of hypergeometric Hamiltonians of the second kind.

Remark 1.7. *We use the geometric interpretation of Gegenbauer Hamiltonians as the main justification for our names of types: spherical, hyperbolic and deSitterian. For coherence, we extend these names to hypergeometric Hamiltonians of both kinds. The names “spherical” and “hyperbolic” seem quite non-controversial, and are used in the literature in similar contexts. The name “deSitterian” is our invention. Its justification is somewhat weaker and based on the geometric interpretation of the Gegenbauer Hamiltonian, see (1.45). The natural geometric interpretation of “deSitterian hypergeometric Hamiltonians of the first kind”, which we described in (1.48) involves the hyperboloid $\mathbb{H}^{p-1,p}$, which is a deSitter space only for $p = 2$.*

1.5 Boundary conditions

For some potentials the operator L initially defined on $C_c^\infty[a, b]$ possesses many closed realizations. These realizations L_\bullet differ only by the behavior of elements of their domain near the endpoints—in other words, they differ by *boundary conditions*. The need for boundary conditions depends on the behavior of the potential V near these endpoints.

Let us consider e.g. the right endpoint b . There are two possibilities:

1. One does not need to impose boundary condition at b . This will be denoted $\nu_b(L) = 0$.
2. There is a 1-parameter family of boundary conditions at b . This is denoted $\nu_b(L) = 2$.

Analogous definitions are valid for the other endpoint a .

The 9 families considered in this paper illustrate Hamiltonians with various kinds of behaviors of the potential near endpoints. Let us list the behaviors encountered among these 9 families. We restrict ourselves to the right endpoint b , an analogous list applies to the left endpoint a . Our description is somewhat informal; for rigorous statements we refer to [DL].

1. **Short range potential.** $b = +\infty$ and $V(x)$ is integrable near $+\infty$ or constant plus integrable. Then $\nu_b(L) = 0$. Moreover, eigenfunctions in $\mathcal{D}(L_\bullet)$ with eigenvalue $-k^2 + V(+\infty)$, $\text{Re}(k) > 0$ behave as e^{-kx} with $\text{Re}(k) > 0$.

2. **Bessel type.** b is finite and

$$V(x) \sim \left(m^2 - \frac{1}{4}\right) \frac{1}{(x-b)^2}. \quad (1.49)$$

Then $\nu_b(L) = 2$ iff $|\operatorname{Re} m| < 1$, otherwise $\nu_b(L) = 0$. The behavior of elements of $\mathcal{D}(L_\bullet)$ near b is $|x-b|^{\frac{1}{2}+m}$ for $\operatorname{Re} m \geq 1$, a linear combination of $|x-b|^{\frac{1}{2}+m}$ and $|x-b|^{\frac{1}{2}-m}$ for $0 \leq \operatorname{Re} m < 1$, $m \neq 0$, and a linear combination of $|x-b|^{\frac{1}{2}}$ and $|x-b|^{\frac{1}{2}} \ln|x-b|$ for $m = 0$.

The families of operators considered in this paper will always have only “homogeneous” or “basic” boundary conditions given by $|x-b|^{\frac{1}{2}+m}$, with $\operatorname{Re} m > -1$. Thus in particular for $|\operatorname{Re} m| < 1$, $m \neq 0$, each differential expression will have two closed realizations. We will not consider mixed boundary conditions, which are discussed e.g. in. [GTV, DeGe, DeRi].

Note that $|x-b|^{\frac{1}{2}+m}$ for $m = \frac{1}{2}$ is $|x-b|$, and for $m = -\frac{1}{2}$ is $|x-b|^0$. Therefore, $m = \frac{1}{2}$ corresponds to the Dirichlet b.c. and $m = -\frac{1}{2}$ to the Neumann b.c. We discuss separately these cases in Sect. 6.

3. **Whittaker type.** b is finite and

$$V(x) \sim \left(m^2 - \frac{1}{4}\right) \frac{1}{(x-b)^2} - \frac{\beta}{|x-b|}. \quad (1.50)$$

The conditions are essentially the same as in the Bessel type, with one difference. For $\frac{1}{2} \leq \operatorname{Re} m < 1$, $(\beta, m) \neq (0, \frac{1}{2})$ the behavior of functions in the domain of closed realizations of L are linear combinations of $|x-b|^{\frac{1}{2}+m}$ and $|x-b|^{\frac{1}{2}-m} \left(1 - \frac{\beta|x-b|}{1-2m}\right)$.

In the families of operators considered in this paper we will only consider “basic” boundary conditions given by $|x-b|^{\frac{1}{2}+m} \left(1 - \frac{\beta|x-b|}{1-2m}\right)$, with $\operatorname{Re} m > -1$. See e.g. [DL, DeRi]

Table 1: Boundary behavior of the potentials.

Potential	Boundary type at a	Boundary type at b
Spherical of first kind	Bessel	Bessel
Hyperbolic of first kind	Bessel	Short range potential
DeSitterian of first kind	Short range potential	Short range potential
Spherical of second type	Whittaker	Whittaker
Hyperbolic of second type	Whittaker	Short range potential
DeSitterian of second type	Short range potential	Short range potential

1.6 Comparison with literature and historic remarks

One can roughly divide mathematical literature related to the topic of this paper into two parts: algebraic and functional-analytic.

In algebraic papers one considers differential expressions without a Hilbert space setting and without asking for self-adjointness or closedness. Functional analytic papers treat differential operators as unbounded operators on a certain Hilbert space, usually self-adjoint, sometimes only closed.

Needless to say, the algebraic literature is vast. In fact, the hypergeometric equation is one of the most classic subjects of mathematics, with history going back about three centuries. From this category one

should mention [CKS, DW] which contain an algebraic analysis of all 9 families of Schrödinger operators solvable in terms of the hypergeometric function.

1-dimensional Schrödinger operators are naturally a special case of Sturm-Liouville operators, whose history goes back to [Lio]. There exists large contemporary literature about self-adjoint or closed realizations of Sturm-Liouville operators, see e.g. [GeZin, GTV, DuSch, EE]. In our paper we use mostly [DeGe], which is summarized in Section 2 of [DL].

Each Sturm-Liouville operator can appear in many equivalent forms, often with distinct names. By a unitary transformation, called the Liouville transformation, essentially each of them can be transformed into a Schrödinger operator, often called its *Liouville form* [Lio].

For instance, in the literature one often considers the Sturm-Liouville operator called the *Jacobi operator*,

$$-(1-x)^{-\alpha}(1+x)^{-\beta}\partial_x(1-x)^{\alpha+1}(1+x)^{\beta+1}\partial_x, \quad (1.51)$$

which acts on the Hilbert space $L^2([-1, 1], (1-x)^\alpha(1+x)^\beta)$ see e.g. [GPLS, Koo]. Its eigenfunctions are the famous Jacobi polynomials. By a simple transformation the Jacobi operator is unitarily equivalent to the trigonometric Pöschl-Teller Hamiltonian (which we call spherical hypergeometric Hamiltonian of the first kind).

Another common differential operator is the Legendre operator

$$(1+x^2)\partial_x^2 - 2x\partial_x + \mu(\mu+1) - \frac{\alpha^2}{1-x^2}. \quad (1.52)$$

Acting on the Hilbert space $L^2([-1, 1], \sqrt{1-x^2})$, used in the study of spherical harmonics. It is also equivalent to the trigonometric Pöschl-Teller Hamiltonian with $\alpha = \beta$. Needless to say, spherical harmonics possess a very large literature.

The trigonometric Pöschl-Teller Hamiltonian itself is also often studied, see e.g. [FS].

The Scarf Hamiltonian with $\alpha = \beta$ (which we call the deSitterian Gegenbauer Hamiltonian) has also a large literature because of its special properties: for certain values of parameters it is reflectionless.

An interesting review of various exactly solvable Schrödinger operators is contained in [Eve]. It does not, however, contain all 9 families that we consider.

To our knowledge, a complete analysis of all 9 families interpreted as closed operators, including the formulas for the integral kernels of their resolvents and the computation of their spectra, seems to appear for the first time in the literature. The transmutation formulas described in our paper are probably new. They are analogous to the transmutation formulas for Hamiltonians related to the confluent equation [DL]. Another novelty of our paper are the identities (1.16) and (1.17). They are analogous to an identity for the Whittaker operators described in [DeRi].

Let us briefly outline the history of these Hamiltonians in physics. In the physics community, the study of Schrödinger operators exactly solvable in terms of hypergeometric functions started in the 1930's when physicists studied biatomic and polyatomic molecules dynamics and the exact solution to the Schrödinger equation. The main reason these physicists considered such Hamiltonians was often due to the limitations of the perturbation method. Therefore, various kinds of exactly solvable Hamiltonians were suggested which fit the experimental data. That was the primary motivation behind the Hamiltonians proposed by Morse [Mor], Rosen-Morse [MoR], Eckart [Eck], and Manning-Rosen [MaR].

The second motivation for this line of research seems to be the search for exact solutions to the Schrödinger equation. In the work of Eckart [Eck] and Rosen-Morse [MoR], it is clearly noted that these potentials are new exactly solvable potentials, although their main motivation was still the study of polyatomic molecules. It seems that for Pöschl and Teller [PT], the motivation leaned more towards the fact that their proposed potential was exactly solvable—they even referred to it as *exakt integrierbar* (exactly integrable).

The case of the Scarf Hamiltonian (1.9) is a bit different. In the original paper, Scarf did not consider the Hamiltonian in (4.43). We traced this naming to [CKS], where authors referred to it as the hyperbolic Scarf, or Scarf II. However, the origin of this naming is unclear to us.

The uniform study of the Schrödinger equation of hypergeometric type was started by Bose [Bos] and continued by Natanzon [Nat], Ginocchio [Gin], and Milson [Mil]. For a more systematic study and a detailed history of the topic, we refer to [DW]. In a separate line of research, the study of these potentials appeared in the factorization method of Infeld and Hull [HI], and later in the context of supersymmetric quantum mechanics and the so-called shape invariance [CKS, Cot].

Table 2 presents a comparison of the various names used in the literature and our suggested terminology.

Table 2: Names of hypergeometric Hamiltonians appearing in the literature.

Our suggestion	Name in [DW]	Name in [CKS]	Alternative names
Spherical of 1st kind	Trigonometric Pöschl–Teller	Scarf I	Pöschl–Teller of 1st kind, Trigonometric Scarf
Hyperbolic of 1st kind	Hyperbolic Pöschl–Teller	Scarf II	Pöschl–Teller of 2nd kind
DeSitterian of 1st kind	Scarf	Generalized Pöschl–Teller	Hyperbolic Scarf
Spherical of 2nd kind	Rosen–Morse	Rosen–Morse I	Trigonometric Rosen–Morse
Hyperbolic of 2nd kind	Eckart	Eckart	Generalized Morse, Hulthén
DeSitterian of 2nd kind	Manning–Rosen	Rosen–Morse II	Hyperbolic Rosen–Morse, Woods–Saxon

The geometric interpretation of hypergeometric Hamiltonians is closely related to the analysis of hypergeometric equation based on Lie groups and Lie algebras, which possess large literature [M1, V, Wa, D2]. The interpretation of the deSitterian hypergeometric Hamiltonian of the 1st kind in terms of complex spheres in 1.48 seems to be new.

1.7 Plan of the paper

In Sect. 2 we recall the definitions of hypergeometric and Gegenbauer function (the latter, following the conventions of [DGR]), and we sketch the Liouville method that allows us to transform a Sturm-Liouville operator into a 1-dimensional Schrödinger operator.

The core of the paper are the sections 3, 4 and 5, where the $3 \times 3 = 9$ families of Hamiltonians are introduced and studied. For each member of each family its spectrum and Green functions are computed. We also prove various identities that we already listed in the introduction, including the transmutation identities. These results are described in 9 theorems, which have similar proofs, based mostly on the abstract theory described in Appendix D, and more fully in [DL, DeGe]. The proof of the first theorem, Thm 3.1, is spelled out in more detail, the next ones are more concise.

Section 6 is devoted to 1-dimensional Laplacians with various boundary conditions. We explain why they are special cases of hypergeometric Hamiltonians.

In Sect. 7 we describe how Gegenbauer Hamiltonians and hypergeometric Hamiltonians of the first kind arise when we separate variables in (pseudo-)Laplacians on some symmetric spaces.

In Appendix A we recall the definition of an analytic family of closed operators following [Kato, DW1].

In Appendix B we give a concise introduction to differential equations in the complex domain and to the Riemann equation (also called the Papperitz equation). It is very useful to view the Gauss hypergeometric equation as a special case of the Riemann equation. We introduce the notation for the Riemann operator, which is directly inspired by the classic notation due to Papperitz [WW].

In Appendix C we collect identities about hypergeometric and Gegenbauer functions that we use in our paper.

In Appendix D we give a concise account of the theory of closed realizations of 1d Schrödinger operators. This account is incomplete and not fully rigorous—the reader should consult Sect. 2 of [DL] for a more detailed and rigorous exposition, or [DeGe], where a complete theory with proofs is given. Of course, the topic is classic and contained in other texts such as [DuSch, EE, DeGe].

In our paper we try to use notation and conventions that make our formulas, especially for Green functions, as simple, elegant and symmetric as possible. This often motivates us to introduce our conventions, different from the standard ones. In particular, we do not use the standard conventions for associated Legendre functions, which could be used for Gegenbauer Hamiltonians. For readers used to associated Legendre functions, in Appendix E we recall their definitions and compare them with the special functions that we use.

2 Preliminaries

2.1 Hypergeometric equation

The *hypergeometric equation* is given by the *hypergeometric operator*

$$\mathcal{F}(a, b; c; z, \partial_z) := z(1-z)\partial_z^2 + (c - (a+b+1)z)\partial_z - ab, \quad (2.1)$$

where a, b, c are arbitrary complex parameters. One of solutions of the hypergeometric equation is the famous hypergeometric function $F(a, b; c; z)$. It is usually convenient to apply to $F(a, b; c; z)$ the so-called *Olver's normalization*, which yields the function

$$\mathbf{F}(a, b; c; z) := \frac{F(a, b; c; z)}{\Gamma(c)} = \sum_{j=0}^{\infty} \frac{(a)_j (b)_j}{\Gamma(c+j)j!} z^j. \quad (2.2)$$

Note that the symmetries of the hypergeometric equation are better visible if we replace a, b, c with α, β, μ :

$$\begin{aligned} \alpha &= c - 1 & \beta &= a + b - c, & \mu &= a - b; \\ a &= \frac{1+\alpha+\beta+\mu}{2}, & b &= \frac{1+\alpha+\beta-\mu}{2}, & c &= 1 + \alpha; \end{aligned} \quad (2.3)$$

so that the hypergeometric equation has the form

$$\mathcal{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha+\beta-\mu+1}{2}; 1 + \alpha; z, \partial_z\right) F(z) = 0. \quad (2.4)$$

To understand the properties and importance of the hypergeometric equation it is useful to view it as a special case of the Riemann equation—the most general equation on the Riemann sphere possessing at most 3 singular points, all of them regular-singular. There are many books and survey articles that comprehensively treat the Riemann and hypergeometric equations, including [WW, Ince, NU, NIST1]. Throughout this paper, we follow the notation established in [DW, D1, D2]. For the convenience of the reader in Appendix B we explain how the hypergeometric equation arises from the Riemann equation.

2.2 Gegenbauer equation

The *Gegenbauer equation* is essentially the special case of the hypergeometric equation with the symmetry $w \rightarrow -w$ and the singular points put at $-1, 1, \infty$. It is given by the *Gegenbauer operator*

$$\mathcal{G}_{\alpha,\lambda}(w, \partial_w) := (1 - w^2)\partial_w^2 - 2(1 + \alpha)w\partial_w + \lambda^2 - (\alpha + \frac{1}{2})^2. \quad (2.5)$$

Here is its relationship to the Riemann operator (defined in (B.64)):

$$\mathcal{G}_{\alpha,\lambda}(w, \partial_w) = (1 - w^2)\mathcal{P} \begin{pmatrix} -1 & 1 & \infty & \\ 0 & 0 & \alpha + \lambda + \frac{1}{2} & w, \partial_w \\ -\alpha & -\alpha & \alpha - \lambda + \frac{1}{2} & \end{pmatrix}. \quad (2.6)$$

Following [DGR], we introduce two special solutions of the Gegenbauer equation

$$\mathbf{S}_{\alpha,\lambda}(w) = \frac{1}{\Gamma(\alpha + 1)} F \left(\frac{1}{2} + \alpha + \lambda, \frac{1}{2} + \alpha - \lambda; \alpha + 1; \frac{1 - w}{2} \right), \quad (2.7)$$

$$\mathbf{Z}_{\alpha,\lambda}(w) = \frac{(w \pm 1)^{-\frac{1}{2} - \alpha - \lambda}}{\Gamma(\lambda + 1)} F \left(\frac{1}{2} + \alpha + \lambda, \frac{1}{2} + \lambda; 2\lambda + 1; \frac{2}{1 \pm w} \right). \quad (2.8)$$

2.3 Liouville transformation

An operator of the form

$$-\partial_r^2 + V(r) \quad (2.9)$$

will be called a (*1-dimensional*) *Schrödinger operator*.

Let us briefly describe how to transform a 2nd order equation

$$(p(z)\partial_z^2 + q(z)\partial_z + r(z))u(z) = 0, \quad (2.10)$$

into an eigenvalue equation of a certain Schrödinger operator. Consider the operator

$$p(z)\partial_z^2 + q(z)\partial_z + r(z), \quad (2.11)$$

that defines the equation (2.10). We first multiply (2.11) from the left by a function f , from the right by a function g , obtaining

$$f(z)(p(z)\partial_z^2 + q(z)\partial_z + r(z))g(z). \quad (2.12)$$

We choose f, g in such a way, that (2.12) has the form

$$-t(z)\partial_z^2 - \frac{1}{2}t'(z)\partial_z + v(z), \quad (2.13)$$

for some function $z \mapsto t(z)$. Then we change the variable z into r , such that

$$\left(\frac{dz}{dr} \right)^2 = t(z). \quad (2.14)$$

We obtain

$$-\partial_r^2 + v(z(r)). \quad (2.15)$$

More details can be found in [DW].

Operators of the form (2.11) are often called *Sturm-Liouville operators* and the transformation that leads from (2.11) to (2.15)—a *Liouville transformation*.

For brevity, we will usually use the term *Hamiltonian* instead of (*one-dimensional*) *Schrödinger operator*. Thus e.g. the *hypergeometric operator* means the operator (2.1), whereas *hypergeometric Hamiltonians* will be various Schrödinger operators obtained by transforming the hypergeometric equation.

3 Gegenbauer Hamiltonians

Let us transform the Gegenbauer operator (2.5) as follows:

$$\begin{aligned}
 & -(1-w^2)^{\frac{\alpha}{2}+\frac{1}{4}} \mathcal{G}_{\alpha,\lambda}(w, \partial_w) (1-w^2)^{-\frac{\alpha}{2}-\frac{1}{4}} \\
 = & -(1-w^2) \mathcal{P} \begin{pmatrix} -1 & 1 & \infty \\ \frac{\alpha}{2} + \frac{1}{4} & \frac{\alpha}{2} + \frac{1}{4} & \lambda \\ -\frac{\alpha}{2} + \frac{1}{4} & -\frac{\alpha}{2} + \frac{1}{4} & -\lambda \end{pmatrix} w, \partial_w \\
 = & -(1-w^2) \partial_w^2 + w \partial_w + \left(\alpha^2 - \frac{1}{4} \right) \frac{1}{1-w^2} - \lambda^2. \tag{3.1}
 \end{aligned}$$

Thus if we set

$$L_\alpha := -(1-w^2) \partial_w^2 + w \partial_w + \left(\alpha^2 - \frac{1}{4} \right) \frac{1}{1-w^2}, \tag{3.2}$$

and $G(w)$ solves the Gegenbauer equation, then

$$(L_\alpha - \lambda^2) (1-w^2)^{\frac{\alpha}{2}+\frac{1}{4}} G(w) = 0. \tag{3.3}$$

We have interpreted the Gegenbauer equation as the eigenequation of the operator L_α with the eigenvalue λ^2 acting on functions on the complex plane. We would like to reinterpret it as an operator on functions of the real variable $r \in]a, b[$ for some $-\infty \leq a < b \leq +\infty$. To this end we need to insert the intervals $]a, b[$ into the complex plane. We will consider three intervals; in each case their endpoints are among the singular points of the Gegenbauer equation (that is, $1, -1, \infty$). For each of these intervals we perform the Liouville transformation, which yields a 1-dimensional Hamiltonian on $C_c^\infty]a, b[$. As we will see below, in each of these cases we will obtain a natural holomorphic family of closed realizations possessing elegant formulas for the Green functions.

Here are the three cases that we consider:

1. $w \in]-1, 1[$. This leads to an operator on $L^2]0, \pi[$, which we call the *spherical Gegenbauer Hamiltonian*.
2. $w \in]1, \infty[$. This leads to an operator on $L^2(\mathbb{R}_+)$, which we call the *hyperbolic Gegenbauer Hamiltonian*.
3. $w \in i\mathbb{R}$. This leads to an operator on $L^2(\mathbb{R})$, which we call the *deSitterian Gegenbauer Hamiltonian*.




Figure 1: Gegenbauer equation on the w plane.

The spherical Hamiltonian acts on the interval marked with a thick line, the hyperbolic Hamiltonian—with a dotted line, and the deSitterian Hamiltonian—with a dashed line.

3.1 Spherical case

For $r \in]0, \pi[$, in (3.1) set

$$w = \cos r, \quad \text{which solves } w' = -(1 - w^2)^{\frac{1}{2}}. \quad (3.4)$$

This leads to the Schrödinger equation

$$(L_\alpha^s - \lambda^2) \phi(r) = 0, \quad (3.5)$$

where

$$L_\alpha^s := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4} \right) \frac{1}{\sin^2 r}. \quad (3.6)$$

It has the mirror symmetry $r \rightarrow \pi - r$. It is obtained when we separate variables of the Laplacian on the sphere in any dimensions, see e.g. Subsect. 7.1. Hence our name “spherical”.

Let us define the function on $]0, \pi[$

$$\mathcal{P}_{\alpha, \lambda}^s(r) := \left(\frac{\sin r}{2} \right)^{\alpha + \frac{1}{2}} \mathbf{S}_{\alpha, \lambda}(\cos r). \quad (3.7)$$

It has the following asymptotic behaviour near 0:

$$\mathcal{P}_{\alpha, \lambda}^s(r) \sim \frac{1}{\Gamma(1 + \alpha)} \left(\frac{r}{2} \right)^{\frac{1}{2} + \alpha}. \quad (3.8)$$

The following four functions solve the eigenequation (3.5):

$$\mathcal{P}_{\alpha, \lambda}^s(r), \quad \mathcal{P}_{-\alpha, \lambda}^s(r), \quad \mathcal{P}_{\alpha, \lambda}^s(\pi - r), \quad \mathcal{P}_{-\alpha, \lambda}^s(\pi - r). \quad (3.9)$$

The following symmetries hold

$$\mathcal{P}_{\alpha, \lambda}^s(r) = \mathcal{P}_{\alpha, -\lambda}^s(r), \quad \mathcal{P}_{\alpha, \lambda}^s(\pi - r) = \mathcal{P}_{\alpha, -\lambda}^s(\pi - r). \quad (3.10)$$

The following theorem describes the basic family of closed realizations of L_α^s on $L^2]0, \pi[$.

Theorem 3.1. *For $\operatorname{Re} \alpha \geq 1$ there exists a unique closed operator L_α^s in the sense of $L^2]0, \pi[$, which on $C_c^\infty]0, \pi[$ is given by (3.6). The family $\alpha \mapsto L_\alpha^s$ is holomorphic and possesses a unique holomorphic extension to $\operatorname{Re} \alpha > -1$. It has only discrete spectrum:*

$$\sigma(L_\alpha^s) = \sigma_d(L_\alpha^s) = \left\{ (k + \alpha)^2 : k \in \mathbb{N}_0 + \frac{1}{2} \right\}. \quad (3.11)$$

Outside of the spectrum its resolvent is

$$\begin{aligned} \frac{1}{(L_\alpha^s - \lambda^2)}(x, y) &= \Gamma\left(\alpha - \lambda + \frac{1}{2}\right) \Gamma\left(\alpha + \lambda + \frac{1}{2}\right) \\ &\times \begin{cases} \mathcal{P}_{\alpha, \lambda}^s(x) \mathcal{P}_{\alpha, \lambda}^s(\pi - y), & \text{if } 0 < x < y < \pi; \\ \mathcal{P}_{\alpha, \lambda}^s(y) \mathcal{P}_{\alpha, \lambda}^s(\pi - x), & \text{if } 0 < y < x < \pi. \end{cases} \end{aligned} \quad (3.12)$$

Proof. Treating $\mathcal{P}_{\alpha,\lambda}^s(r)$ and $\mathcal{P}_{-\alpha,\lambda}^s(r)$ as a basis of solutions of (3.5), we obtain a connection formula found using (C.22):

$$\mathcal{P}_{\alpha,\lambda}^s(\pi - r) = -\frac{\cos \pi \lambda}{\sin \pi \alpha} \mathcal{P}_{\alpha,\lambda}^s(r) + \frac{\pi}{\sin \pi \alpha} \frac{1}{\Gamma(\frac{1}{2} + \alpha + \lambda) \Gamma(\frac{1}{2} + \alpha - \lambda)} \mathcal{P}_{-\alpha,\lambda}^s(r). \quad (3.13)$$

From

$$\mathcal{P}_{-\alpha,\lambda}^s(r) \sim \frac{1}{\Gamma(1 - \alpha)} \left(\frac{r}{2}\right)^{\frac{1}{2} - \alpha}, \quad (3.14)$$

and (3.8) we obtain

$$\mathcal{W}(\mathcal{P}_{-\alpha,\lambda}^s, \mathcal{P}_{\alpha,\lambda}^s) = \frac{\sin \pi \alpha}{\pi}. \quad (3.15)$$

This yields the Wronskians

$$\mathcal{W}(\mathcal{P}_{\alpha,\lambda}^s(\pi - r), \mathcal{P}_{\alpha,\lambda}^s(r)) = \frac{1}{\Gamma(\alpha - \lambda + \frac{1}{2}) \Gamma(\alpha + \lambda + \frac{1}{2})} \quad (3.16)$$

$$\mathcal{W}(\mathcal{P}_{\alpha,\lambda}^s(\pi - r), \mathcal{P}_{-\alpha,\lambda}^s(r)) = \frac{\cos(\pi \lambda)}{\pi}. \quad (3.17)$$

Thus (3.12) is an integral kernel of the form (D.7). Let us denote it by $R_\alpha^s(-\lambda^2; x, y)$.

Let us fix temporarily λ . For $\operatorname{Re} \alpha > -1$ the integral kernel $R_\alpha^s(-\lambda^2; x, y)$ is square integrable, and hence it defines a Hilbert-Schmidt operator, which we denote $R_\alpha^s(-\lambda^2)$. By Appendix D.2 there exists an operator L_α^s , which is a closed realization of the left hand side of (3.6) such that $R_\alpha^s(-\lambda^2) = (L_\alpha^s + \lambda^2)^{-1}$.

The Hilbert-Schmidt norm of $R_\alpha^s(-\lambda^2)$ is uniformly bounded for $\alpha \in K$ with $K \subset \{\operatorname{Re} \alpha > -1\}$ compact. Besides, $R_\alpha^s(-\lambda^2; x, y)$ for any x, y depends analytically on α . Hence for any $f, g \in C_c(]0, \pi[)$

$$\alpha \mapsto \int \overline{f(x)} R_\alpha^s(-\lambda^2; x, y) g(y) dx dy \quad (3.18)$$

is analytic. Using Thm A.1 we obtain that $\{\operatorname{Re} \alpha > 0\} \ni \alpha \mapsto R_\alpha^s(-\lambda^2) \in B(L^2]0, \pi[)$ is an analytic family of bounded operators. Hence, by Thm A.2, $\{\operatorname{Re} \alpha > 0\} \ni \alpha \mapsto L_\alpha^s \in C(L^2]0, \pi[)$ is an analytic family of closed operators.

For $\operatorname{Re} \alpha \geq 1$, $r^{\frac{1}{2} - \alpha}$ is not L^2 -integrable, and only $r^{\frac{1}{2} + \alpha}$ is. Hence for such α the right hand side of (3.6) possesses a unique closed realization, which has to coincide with the operator L_α^s that we have just constructed. In particular, L_α^s does not depend on λ . By the uniqueness of analytic continuation, L_α^s does not depend on the choice of λ for all $\operatorname{Re} \alpha > -1$.

The singularities of the Gamma function yield the discrete spectrum of L_α^s . \square

3.2 Hyperbolic case

In (3.1), for $r \in \mathbb{R}_+$ we set

$$w = \cosh r, \quad \text{which solves } w' = (w^2 - 1)^{\frac{1}{2}}. \quad (3.19)$$

This leads to the Schrödinger equation

$$(L_\alpha^h + \lambda^2) \phi(r) = 0, \quad (3.20)$$

where

$$L_\alpha^h := -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\sinh^2 r}. \quad (3.21)$$

It is obtained when we separate variables of the Laplacian on the hyperbolic space of any dimensions, see e.g. Subsect. 7.2. Hence our name “hyperbolic”.

Let us define functions on \mathbb{R}_+

$$\mathcal{P}_{\alpha,\lambda}^h(r) := \left(\frac{\sinh r}{2}\right)^{\alpha+\frac{1}{2}} \mathbf{S}_{\alpha,\lambda}(\cosh r), \quad (3.22)$$

$$\mathcal{Q}_{\alpha,\lambda}^h(r) := \frac{(\sinh r)^{\alpha+\frac{1}{2}}}{2^\lambda} \mathbf{Z}_{\alpha,\lambda}(\cosh r). \quad (3.23)$$

They have the following asymptotic behavior:

$$\mathcal{P}_{\alpha,\lambda}^h(r) \sim \frac{1}{\Gamma(1+\alpha)} \left(\frac{r}{2}\right)^{\frac{1}{2}+\alpha}, \quad r \sim 0; \quad (3.24)$$

$$\mathcal{Q}_{\alpha,\lambda}^h(r) \sim \frac{1}{\Gamma(1+\lambda)} e^{-\lambda r}, \quad r \rightarrow +\infty. \quad (3.25)$$

The following four functions solve the eigenequation (3.20):

$$\mathcal{P}_{\alpha,\lambda}^h(r), \quad \mathcal{P}_{-\alpha,\lambda}^h(r), \quad \mathcal{Q}_{\alpha,\lambda}^h(r), \quad \mathcal{Q}_{\alpha,-\lambda}^h(r). \quad (3.26)$$

The following symmetries are obvious:

$$\mathcal{P}_{\alpha,\lambda}^h(r) = \mathcal{P}_{\alpha,-\lambda}^h(r), \quad \mathcal{Q}_{\alpha,\lambda}^h(r) = \mathcal{Q}_{-\alpha,\lambda}^h(r). \quad (3.27)$$

The following theorem describes the basic family of closed realizations of L_α^h on $L^2[0, \infty[$.

Theorem 3.2. *For $\operatorname{Re} \alpha \geq 1$ there exists a unique closed operator L_α^h in the sense of $L^2(\mathbb{R}_+)$, which on $C_c^\infty(\mathbb{R}_+)$ is given by (3.21). The family $\alpha \mapsto L_\alpha^h$ is holomorphic and possesses a unique holomorphic extension to $\operatorname{Re} \alpha > -1$. Here is its discrete spectrum and spectrum:*

$$\sigma_d(L_\alpha^h) = \left\{ -\left(\frac{1}{2} + \alpha\right)^2 \right\}, \quad -1 < \operatorname{Re} \alpha < -\frac{1}{2}; \quad (3.28)$$

$$\sigma_d(L_\alpha^h) = \emptyset, \quad -\frac{1}{2} \leq \operatorname{Re} \alpha; \quad (3.29)$$

$$\sigma(L_\alpha^h) = [0, +\infty[\cup \sigma_d(L_\alpha^h). \quad (3.30)$$

Outside of the spectrum, for $\operatorname{Re} \lambda > 0$, its resolvent is

$$\begin{aligned} \frac{1}{(L_\alpha^h + \lambda^2)}(x, y) &= \frac{\Gamma\left(\frac{1}{2} + \alpha + \lambda\right)}{\sqrt{\pi}} \\ &\times \begin{cases} \mathcal{P}_{\alpha,\lambda}^h(x) \mathcal{Q}_{\alpha,\lambda}^h(y) & \text{if } 0 < x < y < \infty; \\ \mathcal{Q}_{\alpha,\lambda}^h(x) \mathcal{P}_{\alpha,\lambda}^h(y) & \text{if } 0 < y < x < \infty. \end{cases} \end{aligned} \quad (3.31)$$

Proof. Consider $\mathcal{P}_{\alpha,\lambda}^h(r)$ and $\mathcal{P}_{-\alpha,\lambda}^h(r)$ as a basis of solutions of (3.20). The connection formula is

$$\mathcal{Q}_{\alpha,\lambda}^h(r) = -\frac{\sqrt{\pi}}{\sin \pi \alpha \Gamma\left(\frac{1}{2} - \alpha + \lambda\right)} \mathcal{P}_{\alpha,\lambda}^h(r) + \frac{\sqrt{\pi}}{\sin \pi \alpha \Gamma\left(\frac{1}{2} + \alpha + \lambda\right)} \mathcal{P}_{-\alpha,\lambda}^h(r). \quad (3.32)$$

Similarly as in the spherical case, we obtain

$$\mathcal{W}(\mathcal{P}_{-\alpha,\lambda}^h, \mathcal{P}_{\alpha,\lambda}^h) = \frac{\sin \alpha}{\pi}. \quad (3.33)$$

This yields the Wronskians

$$\mathcal{W}(\mathcal{Q}_{\alpha,\lambda}^h(r), \mathcal{P}_{\alpha,\lambda}^h(r)) = \frac{\sqrt{\pi}}{\Gamma(\frac{1}{2} + \alpha + \lambda)}.$$

For $\operatorname{Re} \alpha > -1$ and $\operatorname{Re} \lambda > 0$ the functions $\mathcal{P}_{\alpha,\lambda}^h(r)$ resp. $\mathcal{Q}_{\alpha,\lambda}^h(r)$, are square integrable at the endpoints. Using the Schur test we check that under these conditions the integral kernel 3.31 defines a bounded operator, depending analytically on α .

For $\operatorname{Re} \alpha \geq 1$, $r^{\frac{1}{2}-\alpha}$ is not square integrable near 0. Therefore, for such α (3.21) possesses a unique closed realization.

Looking for singularities of the Gamma function we find the discrete spectrum:

$$\sigma_d(L_\alpha^h) = \left\{ -\left(n + \frac{1}{2} + \alpha\right)^2 \mid n \in \mathbb{N}_0, \operatorname{Re}\left(n + \frac{1}{2} + \alpha\right) < 0 \right\}. \quad (3.34)$$

It is easy to see that this coincides with (3.29) □

3.3 DeSitterian case

For $r \in \mathbb{R}$, in (3.1) we set

$$w = -i \sinh r, \quad \text{which solves } w' = (w^2 - 1)^{\frac{1}{2}}. \quad (3.35)$$

This leads to the Schrödinger equation

$$(L_\alpha^{\text{dS}} + \lambda^2) \phi(r) = 0, \quad (3.36)$$

where

$$L_\alpha^{\text{dS}} := -\partial_r^2 - \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r}. \quad (3.37)$$

It has the mirror symmetry $r \rightarrow -r$. It is obtained when we separate variables in the d'Alembertian on the deSitter space of any dimension, see Subsect. 7.3. Hence our name “deSitterian”.

The following theorem describes all closed realizations of L_α^{dS} on $L^2(\mathbb{R})$.

For $r \geq 0$ we introduce the following function which solves the eigenequation

$$\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(r) := e^{-i\frac{\pi}{2}(\frac{1}{2}+\alpha+\lambda)} \frac{(\cosh r)^{\alpha+\frac{1}{2}}}{2^\lambda} \mathbf{Z}_{\alpha,\lambda}(-i \sinh r). \quad (3.38)$$

We extend it to $r \leq 0$ by analytic continuation. Here is its asymptotics:

$$\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(r) \sim \frac{1}{\Gamma(\lambda+1)} e^{-\lambda r}, \quad r \rightarrow +\infty. \quad (3.39)$$

Thus the following functions solve the eigenequation (3.36):

$$\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(r), \quad \mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(-r), \quad \mathcal{Q}_{\alpha,-\lambda}^{\text{dS}}(r), \quad \mathcal{Q}_{\alpha,-\lambda}^{\text{dS}}(-r). \quad (3.40)$$

Let us describe closed realizations of L_α^{dS} on $L^2(\mathbb{R})$:

Theorem 3.3. For any $\alpha \in \mathbb{C}$ there exists a unique closed operator L_α^{dS} in the sense of $L^2(\mathbb{R})$ that on $C_c^\infty(\mathbb{R})$ is given by (3.37). The function $\mathbb{C} \ni \alpha \mapsto L_\alpha^{\text{dS}}$ is holomorphic. It satisfies $L_\alpha^{\text{dS}} = L_{-\alpha}^{\text{dS}}$. Outside of the spectrum, for $\text{Re } \lambda > 0$, its resolvent is

$$\begin{aligned} & \frac{1}{(L_\alpha^{\text{dS}} + \lambda^2)}(x, y) \\ &= \frac{\Gamma\left(\frac{1}{2} - \alpha + \lambda\right) \Gamma\left(\frac{1}{2} + \alpha + \lambda\right)}{2} \begin{cases} \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(x) \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(-y) & -\infty < x < y < \infty; \\ \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(y) \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(-x) & -\infty < y < x < \infty. \end{cases} \end{aligned} \quad (3.41)$$

To describe the discrete spectrum and spectrum of L_α^{dS} , without loss of generality we can assume that $\text{Re } \alpha \geq 0$. Then

$$\sigma_{\text{d}}(L_\alpha^{\text{dS}}) = \left\{ -(\alpha - k)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < \text{Re } \alpha \right\} \quad (3.42)$$

$$\sigma(L_\alpha^{\text{dS}}) = [0, \infty[\cup \sigma_{\text{d}}(L_\alpha^{\text{dS}}). \quad (3.43)$$

Proof. We can use the proof of the theorem 4.3. Note that

$$\mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(r) = \frac{\Gamma\left(\frac{1}{2} + \lambda\right)}{\sqrt{\pi}} \mathcal{Q}_{\alpha, \alpha, 2\lambda}^{\text{dS}}(r) = \frac{\Gamma(1 + 2\lambda)}{2^{2\lambda} \Gamma(1 + \lambda)} \mathcal{Q}_{\alpha, \alpha, 2\lambda}^{\text{dS}}(r). \quad (3.44)$$

Hence, the Wronskians are

$$\mathcal{W}\left(\mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(-r), \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(r)\right) = \frac{2}{\Gamma\left(\frac{1}{2} + \alpha + \lambda\right) \Gamma\left(\frac{1}{2} - \alpha + \lambda\right)}, \quad (3.45)$$

$$\mathcal{W}\left(\mathcal{Q}_{\alpha, -\lambda}^{\text{dS}}(-r), \mathcal{Q}_{\alpha, \lambda}^{\text{dS}}(r)\right) = \frac{2 \cos(\pi\alpha)}{\pi}. \quad (3.46)$$

Using the Schur test we check that the integral kernel (3.41) defines a bounded operator.

The singularities of the Gamma function are at $\lambda = -\frac{1}{2} + \alpha - n$ and $\lambda = -\frac{1}{2} - \alpha - n$, $n \in \mathbb{N}_0$. This gives the following discrete spectrum:

$$\sigma_{\text{d}}(L_\alpha^{\text{dS}}) = \left\{ -\left(\frac{1}{2} + \alpha + n\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(\frac{1}{2} + \alpha + n\right) < 0 \right\} \quad (3.47)$$

$$\cup \left\{ -\left(\frac{1}{2} - \alpha + n\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(\frac{1}{2} - \alpha + n\right) < 0 \right\}. \quad (3.48)$$

For $\text{Re } \alpha \geq 0$ the right hand side of (3.47) is empty. This yields (3.42). \square

Proof of Prop. 1.1. The transformation $\cot r = \sinh q$ implies

$$\cos r = \tanh q, \quad \tan r = \sinh q, \quad (3.49)$$

$$\frac{dq}{dr} = -\frac{1}{\sin r} = \cosh q. \quad (3.50)$$

The Whipple transformation (C.25), (C.26) on the interval $w \in]-1, 1[$ has two versions: above this interval and below, that is

$$\mathbf{Z}_{\alpha, \lambda}(w \pm i0) = (\pm i \sqrt{1 - w^2})^{-\frac{1}{2} - \alpha - \lambda} \mathbf{S}_{\lambda, \alpha} \left(\frac{w}{\pm i \sqrt{1 - w^2}} \right). \quad (3.51)$$

This yields

$$\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(q) = \left(\frac{2}{\sin r}\right)^{\frac{1}{2}} \mathcal{P}_{\lambda,\alpha}^{\text{s}}(r), \quad (3.52)$$

$$\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(-q) = \left(\frac{2}{\sin r}\right)^{\frac{1}{2}} \mathcal{P}_{\lambda,\alpha}^{\text{s}}(\pi - r). \quad (3.53)$$

This yields

$$(\sin r)^{\frac{1}{2}} \frac{1}{(L_{\lambda}^{\text{s}} - \alpha^2)}(r, r') (\sin r')^{\frac{1}{2}} = \frac{1}{(L_{\alpha}^{\text{dS}} - \lambda^2)}(q, q'), \quad (3.54)$$

which proves Prop. 1.1. \square

Remark 3.4. Using (3.50), (3.52) and (3.53) we obtain

$$\mathcal{W}(\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(-q), \mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(q)) = 2\mathcal{W}(\mathcal{P}_{\lambda,\alpha}^{\text{s}}(\pi - r), \mathcal{P}_{\lambda,\alpha}^{\text{s}}(r)), \quad (3.55)$$

$$\mathcal{W}(\mathcal{Q}_{\alpha,\lambda}^{\text{dS}}(-q), \mathcal{Q}_{\alpha,-\lambda}^{\text{dS}}(q)) = 2\mathcal{W}(\mathcal{P}_{\lambda,\alpha}^{\text{s}}(\pi - r), \mathcal{P}_{-\lambda,\alpha}^{\text{s}}(r)). \quad (3.56)$$

Therefore, (3.16) and (3.17) imply (3.45) and (3.46). This can be used in an alternative proof of Thm 3.3.

4 Hypergeometric Hamiltonians of the first kind

Let us transform the hypergeometric operator as follows:

$$\begin{aligned} & -z^{\frac{\alpha}{2}+\frac{1}{4}}(1-z)^{\frac{\beta}{2}+\frac{1}{4}} \mathcal{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha+\beta-\mu+1}{2}; 1+\alpha; z, \partial_z\right) z^{-\frac{\alpha}{2}-\frac{1}{4}}(1-z)^{-\frac{\beta}{2}-\frac{1}{4}} \\ &= -z(1-z) \mathcal{P}\left(\begin{array}{ccc} 0 & 1 & \infty \\ \frac{\alpha}{2}+\frac{1}{4} & \frac{\beta}{2}+\frac{1}{4} & \frac{\mu}{2} \\ -\frac{\alpha}{2}+\frac{1}{4} & -\frac{\beta}{2}+\frac{1}{4} & -\frac{\mu}{2} \end{array}; z, \partial_z\right) \\ & -z(1-z) \left(\partial_z^2 + \left(\frac{1}{2z} - \frac{1}{2(1-z)}\right) \partial_z\right) \\ & + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{4z} + \left(\beta^2 - \frac{1}{4}\right) \frac{1}{4(1-z)} - \frac{\mu^2}{4}. \end{aligned} \quad (4.1)$$

Thus if we set

$$L_{\alpha,\beta} := -z(1-z) \left(\partial_z^2 + \left(\frac{1}{2z} - \frac{1}{2(1-z)}\right) \partial_z\right) + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{4z} + \left(\beta^2 - \frac{1}{4}\right) \frac{1}{4(1-z)} \quad (4.2)$$

and $F(z)$ solves the hypergeometric equation (2.4), then

$$\left(L_{\alpha,\beta} - \frac{\mu^2}{4}\right) z^{\frac{\alpha}{2}+\frac{1}{4}}(1-z)^{\frac{\beta}{2}+\frac{1}{4}} F(z) = 0. \quad (4.3)$$

The hypergeometric equation has been interpreted as the eigenequation of the operator $L_{\alpha,\beta}$ with the eigenvalue $\frac{\mu^2}{4}$ acting on functions on the complex plane. Let us reinterpret this operator as acting on functions on three intervals whose endpoints are singularities of the hypergeometric equations. We will see again that these choices lead to simple formulas for Green functions. In each of these cases we perform the Liouville transformation, which yields a 1-dimensional Hamiltonian. We will consider three cases:

1. $z \in]0, 1[$, which leads to an operator on $L^2]0, \pi[$, which we call the *spherical hypergeometric Hamiltonian of the 1st kind*;
2. $z \in]-\infty, 0[$, which leads to an operator on $L^2(\mathbb{R}_+)$, which we call the *hyperbolic hypergeometric Hamiltonian of the 1st kind*;
3. $z \in \frac{1}{2} + i\mathbb{R}$, which leads to an operator on $L^2(\mathbb{R})$, which we call the *deSitterian hypergeometric Hamiltonian of the 1st kind*.

It will be natural to introduce the parameters

$$\delta := \frac{1}{2}(\alpha^2 + \beta^2), \quad (4.4)$$

$$\kappa := \frac{1}{2}(\alpha^2 - \beta^2), \quad \tau := \frac{i}{2}(\alpha^2 - \beta^2) = i\kappa. \quad (4.5)$$




Figure 2: Hypergeometric Hamiltonians of the first kind on the z -plane. The spherical Hamiltonian acts on the interval marked with a thick line, the hyperbolic Hamiltonian—with a dotted line, and deSitterian Hamiltonian—with a dashed line.

4.1 Spherical case

For $r \in]0, \pi[$, set in (4.2)

$$z = \sin^2 \frac{r}{2} = \frac{1 - \cos r}{2}, \quad \text{which solves } z' = z^{\frac{1}{2}}(1 - z)^{\frac{1}{2}}. \quad (4.6)$$

This leads to the Schrödinger equation

$$\left(L_{\alpha, \beta}^s - \frac{\mu^2}{4} \right) \phi(r) = 0, \quad (4.7)$$

where

$$\begin{aligned} L_{\alpha, \beta}^s &:= -\partial_r^2 + \left(\alpha^2 - \frac{1}{4} \right) \frac{1}{4 \sin^2 \frac{r}{2}} + \left(\beta^2 - \frac{1}{4} \right) \frac{1}{4 \cos^2 \frac{r}{2}} \\ &= -\partial_r^2 + \left(\delta - \frac{1}{4} \right) \frac{1}{\sin^2 r} + \kappa \frac{\cos r}{\sin^2 r}. \end{aligned} \quad (4.8)$$

In [DW] $L_{\alpha,\beta}^s$ is called the *trigonometric Pöschl-Teller Hamiltonian*

The case $\alpha = \beta$ is especially important and coincides with the spherical Gegenbauer Hamiltonian:

$$L_{\alpha}^s := L_{\alpha,\alpha}^s. \quad (4.9)$$

For $r \in]0, \pi[$, define the function

$$\begin{aligned} P_{\alpha,\beta,\mu}^s(r) &:= \left(\sin \frac{r}{2}\right)^{\alpha+\frac{1}{2}} \left(\cos \frac{r}{2}\right)^{\beta+\frac{1}{2}} \mathbf{F} \left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha+\beta-\mu+1}{2}; 1+\alpha; \sin^2 \left(\frac{r}{2}\right) \right) \\ &= \left(\sin \frac{r}{2}\right)^{\alpha+\frac{1}{2}} \left(\cos \frac{r}{2}\right)^{-\alpha-\mu-\frac{1}{2}} \mathbf{F} \left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha-\beta+\mu+1}{2}; 1+\alpha; -\tan^2 \left(\frac{r}{2}\right) \right). \end{aligned} \quad (4.10)$$

Note that

$$P_{\alpha,\beta,\mu}^s(r) = P_{\alpha,-\beta,\mu}^s(r) = P_{\alpha,\beta,-\mu}^s(r). \quad (4.11)$$

Asymptotically our function behaves like

$$P_{\alpha,\beta,\mu}^s(r) \sim \frac{1}{\Gamma(1+\alpha)} \left(\frac{r}{2}\right)^{\alpha+\frac{1}{2}}, \quad r \sim 0. \quad (4.12)$$

Now the following functions solve the eigenvalue problem (4.7):

$$P_{\alpha,\beta,\mu}^s(r), \quad P_{-\alpha,\beta,\mu}^s(r), \quad P_{\beta,\alpha,\mu}^s(\pi-r), \quad P_{-\beta,\alpha,\mu}^s(\pi-r). \quad (4.13)$$

The following theorem describes the basic holomorphic family of closed realizations of $L_{\alpha,\beta}^s$ on $L^2]0, \pi[$.

Theorem 4.1. *For $\operatorname{Re} \alpha, \operatorname{Re} \beta \geq 1$ there exists a unique closed operator $L_{\alpha,\beta}^s$ in the sense of $L^2]0, \pi[$, which on $C_c^\infty]0, \pi[$ is given by (4.8). The family $\alpha, \beta \mapsto L_{\alpha,\beta}^s$ is holomorphic and possesses a unique holomorphic extension to $\operatorname{Re} \alpha, \operatorname{Re} \beta > -1$. It has only discrete spectrum:*

$$\sigma(L_{\alpha,\beta}^s) = \sigma_d(L_{\alpha,\beta}^s) = \left\{ \left(k + \frac{\alpha+\beta}{2}\right)^2 : k \in \mathbb{N}_0 + \frac{1}{2} \right\}. \quad (4.14)$$

Outside of the spectrum its resolvent is

$$\begin{aligned} \frac{1}{\left(L_{\alpha,\beta}^s - \frac{\mu^2}{4}\right)}(x, y) &= \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right) \\ &\times \begin{cases} P_{\alpha,\beta,\mu}^s(x) P_{\beta,\alpha,\mu}^s(\pi-y) & \text{if } 0 < x < y < \pi; \\ P_{\alpha,\beta,\mu}^s(y) P_{\beta,\alpha,\mu}^s(\pi-x) & \text{if } 0 < y < x < \pi. \end{cases} \end{aligned} \quad (4.15)$$

Proof. Considering $P_{\alpha,\beta,\mu}^s(r)$, and $P_{-\alpha,\beta,\mu}^s(r)$ as a basis of solutions of (4.7), we can rewrite the connection formula (C.9) as

$$P_{\beta,\alpha,\mu}^s(\pi-r) = \frac{\pi P_{\alpha,\beta,\mu}^s(r)}{\sin(-\pi\alpha)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)} + \frac{\pi P_{-\alpha,\beta,\mu}^s(r)}{\sin(\pi\alpha)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)}. \quad (4.16)$$

Using (4.12) and arguing as in the proof of (3.15), we obtain

$$\mathcal{W}(P_{-\alpha,\beta,\mu}^s(r), P_{\alpha,\beta,\mu}^s(r)) = \frac{\sin \pi\alpha}{\pi}. \quad (4.17)$$

From the connection formula and (4.17) we obtain the Wronskian

$$\mathcal{W}(P_{\beta,\alpha,\mu}^s(\pi-r), P_{\alpha,\beta,\mu}^s(r)) = \frac{1}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)}. \quad (4.18)$$

The L^2 integrability conditions at the endpoints are $\operatorname{Re} \alpha > -1$, $\operatorname{Re} \beta > -1$, for $P_{\alpha,\beta,\mu}^s(r)$ and $P_{\beta,\alpha,\mu}^s(\pi-r)$ respectively. With these conditions, we can write the candidate for the resolvent (4.15). The L^2 norm of this integral kernel is finite. Hence it defines a bounded (even Hilbert-Schmidt) operator. For $\operatorname{Re} \alpha \geq 1$, $\operatorname{Re} \beta \geq 1$ it is a unique candidate for the resolvent. \square

4.2 Hyperbolic case

For $r \in \mathbb{R}_+$, in (4.2) we set

$$z = -\sinh^2 \frac{r}{2} = \frac{1 - \cosh r}{2}, \quad \text{which solves } z' = -(-z)^{\frac{1}{2}}(1-z)^{\frac{1}{2}}. \quad (4.19)$$

This leads to the Schrödinger equation

$$\left(L_{\alpha,\beta}^h + \frac{\mu^2}{4}\right) \phi(r) = 0, \quad (4.20)$$

where

$$\begin{aligned} L_{\alpha,\beta}^h &:= -\partial_r^2 + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{4 \sinh^2 \frac{r}{2}} - \left(\beta^2 - \frac{1}{4}\right) \frac{1}{4 \cosh^2 \frac{r}{2}} \\ &= -\partial_r^2 + \left(\delta - \frac{1}{4}\right) \frac{1}{\sinh^2 r} + \kappa \frac{\cosh r}{\sinh^2 r}. \end{aligned} \quad (4.21)$$

In [DW] $L_{\alpha,\beta}^h$ is called the *hyperbolic Pöschl-Teller Hamiltonian*.

The case $\alpha = \beta$ is especially important and coincides with the hyperbolic Gegenbauer operator:

$$L_{\alpha}^h := L_{\alpha,\alpha}^h. \quad (4.22)$$

For $r \in \mathbb{R}_+$, let us define

$$\begin{aligned} P_{\alpha,\beta,\mu}^h(r) &:= \left(\sinh \frac{r}{2}\right)^{\alpha+\frac{1}{2}} \left(\cosh \frac{r}{2}\right)^{\beta+\frac{1}{2}} \mathbf{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha+\beta-\mu+1}{2}; 1+\alpha; -\sinh^2 \frac{r}{2}\right) \\ &= \left(\sinh \frac{r}{2}\right)^{\alpha+\frac{1}{2}} \left(\cosh \frac{r}{2}\right)^{-\alpha-\mu-\frac{1}{2}} \mathbf{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha-\beta+\mu+1}{2}; 1+\alpha; \tanh^2 \frac{r}{2}\right), \end{aligned} \quad (4.23)$$

$$\begin{aligned} Q_{\alpha,\beta,\mu}^h(r) &:= \left(\sinh \frac{r}{2}\right)^{-\mu-\beta-\frac{1}{2}} \left(\cosh \frac{r}{2}\right)^{\beta+\frac{1}{2}} \mathbf{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{-\alpha+\beta+\mu+1}{2}; 1+\mu; -\sinh^{-2} \frac{r}{2}\right) \\ &= \left(\sinh \frac{r}{2}\right)^{\alpha+\frac{1}{2}} \left(\cosh \frac{r}{2}\right)^{-\alpha-\mu-\frac{1}{2}} \mathbf{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha-\beta+\mu+1}{2}; 1+\mu; \cosh^{-2} \frac{r}{2}\right). \end{aligned} \quad (4.24)$$

Note that

$$P_{\alpha,\beta,\mu}^h(r) = P_{\alpha,-\beta,\mu}^h(r) = P_{\alpha,\beta,-\mu}^h(r), \quad (4.25)$$

$$Q_{\alpha,\beta,\mu}^h(r) = Q_{\alpha,-\beta,\mu}^h(r) = Q_{-\alpha,\beta,\mu}^h(r), \quad (4.26)$$

Asymptotically,

$$P_{\alpha,\beta,\mu}^h(r) \sim \frac{1}{\Gamma(1+\alpha)} \left(\frac{r}{2}\right)^{\alpha+\frac{1}{2}}, \quad r \rightarrow 0; \quad (4.27)$$

$$Q_{\alpha,\beta,\mu}^h(r) \sim \frac{2^\mu}{\Gamma(1+\mu)} e^{-\frac{\mu}{2}r}, \quad r \rightarrow +\infty. \quad (4.28)$$

Now the following functions solve the eigenvalue problem (4.42):

$$P_{\alpha,\beta,\mu}^h(r), \quad P_{-\alpha,\beta,\mu}^h(r), \quad Q_{\alpha,\beta,\mu}^h(r), \quad Q_{\alpha,\beta,-\mu}^h(r). \quad (4.29)$$

The following theorem describes the basic holomorphic family of closed realizations of $L_{\alpha,\beta}^h$ on the Hilbert space $L^2(\mathbb{R}_+)$.

Theorem 4.2. *For $\operatorname{Re} \alpha \geq 1$ there exists a unique closed operator $L_{\alpha,\beta}^h$ in the sense of $L^2(\mathbb{R}_+)$, which on $C_c^\infty(\mathbb{R}_+)$ is given by (4.21). The family $\alpha, \beta \mapsto L_{\alpha,\beta}^h$ is holomorphic and possesses a unique holomorphic extension to $\operatorname{Re} \alpha > -1$. Its discrete spectrum and spectrum are*

$$\begin{aligned} \sigma_d(L_{\alpha,\beta}^h) = & \left\{ -\left(\frac{\alpha+\beta}{2} + k\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < -\operatorname{Re} \frac{\alpha+\beta}{2} \right\}, \\ & \cup \left\{ -\left(\frac{\alpha-\beta}{2} + k\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < -\operatorname{Re} \frac{\alpha-\beta}{2} \right\}, \end{aligned} \quad (4.30)$$

$$\sigma(L_{\alpha,\beta}^h) = [0, \infty] \cup \sigma_d(L_{\alpha,\beta}^h). \quad (4.31)$$

Outside of the spectrum, for $\operatorname{Re} \mu > 0$, its resolvent is

$$\begin{aligned} \frac{1}{(L_{\alpha,\beta}^h + \frac{\mu^2}{4})}(x, y) = & \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right) \\ & \times \begin{cases} P_{\alpha,\beta,\mu}^h(x) Q_{\alpha,\beta,\mu}^h(y) & \text{if } 0 < x < y < \infty; \\ P_{\alpha,\beta,\mu}^h(y) Q_{\alpha,\beta,\mu}^h(x) & \text{if } 0 < y < x < \infty. \end{cases} \end{aligned} \quad (4.32)$$

Proof. Considering $P_{-\alpha,\beta,\mu}^h(r)$, and $P_{\alpha,\beta,\mu}^h(r)$ as a basis of solutions of (4.42), we can rewrite connection formula (C.13) as

$$Q_{\alpha,\beta,\mu}^h(r) = -\frac{\pi P_{\alpha,\beta,\mu}^h(r)}{\sin \pi \alpha \Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)} + \frac{\pi P_{-\alpha,\beta,\mu}^h(r)}{\sin(\alpha\pi) \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}. \quad (4.33)$$

Using (4.27) and again arguing as in the proof of (3.15), we obtain we obtain

$$\mathcal{W}(P_{-\alpha,\beta,\mu}^h(r), P_{\alpha,\beta,\mu}^h(r)) = \frac{\sin \pi \alpha}{\pi}. \quad (4.34)$$

This yields the Wronskian

$$\mathcal{W}(Q_{\alpha,\beta,\mu}^h(r), P_{\alpha,\beta,\mu}^h(r)) = \frac{1}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}. \quad (4.35)$$

The L^2 integrability conditions at the endpoints are $\operatorname{Re} \alpha > -1$ and $\operatorname{Re} \mu > 0$ for $P_{\alpha,\beta,\mu}^h(r)$ and $Q_{\alpha,\beta,\mu}^h(r)$, respectively. Using the Schur Test we check that integral kernel (4.32) defines a bounded operator. We also see that for $\operatorname{Re} \alpha > -1, \operatorname{Re} \mu > 0$ it is a unique candidate for the resolvent. \square

Proof of Prop. 1.2. The transformation

$$\tan \frac{r}{2} = \sinh \frac{q}{2}. \quad (4.36)$$

implies

$$\sin \frac{r}{2} = \tanh \frac{q}{2}, \quad \frac{dq}{dr} = \frac{1}{\cos \frac{r}{2}} = \cosh \frac{q}{2}; \quad (4.37)$$

$$P_{\alpha,\beta,\mu}^s(r) = \frac{1}{\left(\cosh \frac{q}{2}\right)^{\frac{1}{2}}} P_{\alpha,\mu,\beta}^h(q), \quad (4.38)$$

$$P_{\beta,\alpha,\mu}^s(\pi - r) = \frac{1}{\left(\cosh \frac{q}{2}\right)^{\frac{1}{2}}} Q_{\alpha,\mu,\beta}^h(q). \quad (4.39)$$

We obtain the transmutation relation

$$\frac{1}{\left(L_{\alpha,\beta}^s + \frac{\mu^2}{4}\right)}(r, r') = \left(\cosh \frac{q}{2}\right)^{-\frac{1}{2}} \frac{1}{\left(L_{\alpha,\mu}^h + \frac{\beta^2}{4}\right)}(q, q') \left(\cosh \frac{q'}{2}\right)^{-\frac{1}{2}}. \quad (4.40)$$

□

4.3 DeSitterian case

For $r \in \mathbb{R}$, in (4.2) we set

$$z = \frac{1}{2} - i \cosh \frac{r}{2} \sinh \frac{r}{2} = \frac{1 - i \sinh r}{2}, \quad \text{which solves } z' = (-z)^{\frac{1}{2}}(1 - z)^{\frac{1}{2}}. \quad (4.41)$$

This leads to the Schrödinger equation

$$\left(L_{\alpha,\beta}^{\text{dS}} + \frac{\mu^2}{4}\right) \phi(r) = 0, \quad (4.42)$$

where

$$L_{\alpha,\beta}^{\text{dS}} := -\partial_r^2 - \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r} \left(\frac{1}{2} + \frac{i \sinh r}{2}\right) - \left(\beta^2 - \frac{1}{4}\right) \frac{1}{\cosh^2 r} \left(\frac{1}{2} - \frac{i \sinh r}{2}\right) \quad (4.43)$$

$$= -\partial_r^2 - \left(\delta - \frac{1}{4}\right) \frac{1}{\cosh^2 r} - \tau \frac{\sinh r}{\cosh^2 r}. \quad (4.44)$$

This Hamiltonian was proposed and solved by F. Scarf [Sca] and in [DW] it is called the *Scarf Hamiltonian*.

The case $\alpha = \beta$ is especially important and coincides with the deSitterian Gegenbauer Hamiltonian:

$$L_{\alpha}^{\text{dS}} := L_{\alpha,\alpha}^{\text{dS}}. \quad (4.45)$$

Define for $r \geq 0$

$$Q_{\alpha,\beta,\mu}^{\text{dS}}(r) := \left(\frac{i + \sinh r}{2}\right)^{-\frac{\beta}{2} - \frac{\mu}{2} - \frac{1}{4}} \left(\frac{-i + \sinh r}{2}\right)^{\frac{\beta}{2} + \frac{1}{4}} \\ \times \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{-\alpha + \beta + \mu + 1}{2}; 1 + \mu; \frac{2}{1 - i \sinh r} \right) \quad (4.46)$$

$$= \left(\frac{i + \sinh r}{2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} \left(\frac{-i + \sinh r}{2}\right)^{-\frac{\alpha}{2} - \frac{\mu}{2} - \frac{1}{4}} \\ \times \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; 1 + \mu; \frac{2}{1 + i \sinh r} \right). \quad (4.47)$$

We extend $r \rightarrow Q_{\alpha,\beta,\mu}^{\text{dS}}(r)$ to $r < 0$ by analyticity. It satisfies

$$Q_{\alpha,\beta,\mu}^{\text{dS}}(r) \sim \frac{2^\mu e^{-\frac{\mu}{2}r}}{\Gamma(1+\mu)}, \quad r \rightarrow +\infty. \quad (4.48)$$

Note that

$$Q_{\alpha,\beta,\mu}^{\text{dS}}(r) = Q_{\alpha,-\beta,\mu}^{\text{dS}}(r) = Q_{-\alpha,\beta,\mu}^{\text{dS}}(r). \quad (4.49)$$

Now the following functions solve the eigenvalue problem (4.42):

$$Q_{\alpha,\beta,\mu}^{\text{dS}}(r), \quad Q_{\alpha,\beta,-\mu}^{\text{dS}}(r), \quad Q_{\beta,\alpha,\mu}^{\text{dS}}(-r), \quad Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r). \quad (4.50)$$

The following theorem describes all closed realizations of $L_{\alpha,\beta}^{\text{dS}}$ on $L^2(\mathbb{R})$.

Theorem 4.3. *For any $\alpha, \beta \in \mathbb{C}$ there exists a unique closed operator L_α^{dS} in the sense of $L^2(\mathbb{R})$ that on $C_c^\infty(\mathbb{R})$ is given by (4.43). The function $\mathbb{C} \ni (\alpha, \beta) \mapsto L_\alpha^{\text{dS}}$ is holomorphic. We have $L_{\alpha,\beta}^{\text{dS}} = L_{-\alpha,\beta}^{\text{dS}} = L_{\alpha,-\beta}^{\text{dS}}$.*

Outside of the spectrum, for $\text{Re } \mu > 0$, its resolvent is

$$\begin{aligned} \frac{1}{(L_{\alpha,\beta}^{\text{dS}} + \frac{\mu^2}{4})}(x, y) &= \frac{\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)}{2\pi} \\ &\times \begin{cases} Q_{\alpha,\beta,\mu}^{\text{dS}}(x) Q_{\beta,\alpha,\mu}^{\text{dS}}(-y) & \text{if } -\infty < x < y < \infty; \\ Q_{\beta,\alpha,\mu}^{\text{dS}}(-y) Q_{\alpha,\beta,\mu}^{\text{dS}}(x) & \text{if } -\infty < y < x < \infty. \end{cases} \end{aligned} \quad (4.51)$$

To describe the discrete spectrum of $L_{\alpha,\beta}^{\text{dS}}$ assume without loss of generality that $\text{Re}(\alpha + \beta) \geq 0$. We also assume that $\text{Re}(\alpha - \beta) \geq 0$ (the case $\text{Re}(\alpha - \beta) \leq 0$ is analogous). Then

$$\begin{aligned} \sigma_d(L_{\alpha,\beta}^{\text{dS}}) &= \left\{ -\left(\frac{\alpha + \beta}{2} - k\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < \text{Re} \frac{\alpha + \beta}{2} \right\} \\ &\cup \left\{ -\left(\frac{\alpha - \beta}{2} - k\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < \text{Re} \frac{\alpha - \beta}{2} \right\}, \end{aligned} \quad (4.52)$$

$$\sigma(L_{\alpha,\beta}^{\text{dS}}) = [0, \infty] \cup \sigma_d(L_{\alpha,\beta}^{\text{dS}}). \quad (4.53)$$

Proof. In the connection formula (C.15) we insert $z = \frac{1-is}{2}$ and multiply it by $\left(\frac{1-is}{2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} \left(\frac{1+is}{2}\right)^{\frac{\beta}{2} + \frac{1}{4}}$, obtaining

$$\begin{aligned} \left(\frac{1-is}{2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} \left(\frac{1+is}{2}\right)^{\frac{\beta}{2} + \frac{1}{4}} \mathbf{F}_{\alpha,\beta,\mu} \left(\frac{1-is}{2}\right) &= \frac{\pi \left(\frac{1-is}{2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} \left(\frac{1+is}{2}\right)^{\frac{\beta}{2} + \frac{1}{4}} \left(\frac{is-1}{2}\right)^{-\frac{1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,\beta,\alpha} \left(\frac{2}{1-is}\right)}{\sin(-\pi\mu) \Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} \\ &+ \frac{\pi \left(\frac{1-is}{2}\right)^{\frac{\alpha}{2} + \frac{1}{4}} \left(\frac{1+is}{2}\right)^{\frac{\beta}{2} + \frac{1}{4}} \left(\frac{is-1}{2}\right)^{-\frac{1-\alpha-\beta+\mu}{2}} \mathbf{F}_{-\mu,\beta,\alpha} \left(\frac{2}{1-is}\right)}{\sin(\pi\mu) \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}. \end{aligned} \quad (4.54)$$

We transform separately (4.54) above and below the real line: obtaining resp.

$$\begin{aligned} & \frac{\pi e^{i\frac{\pi}{2}(-\alpha-\frac{1}{2}-\frac{\mu}{2})} \left(\frac{s+i}{2}\right)^{-\frac{\beta}{2}-\frac{1}{4}-\frac{\mu}{2}} \left(\frac{s-i}{2}\right)^{\frac{\beta}{2}+\frac{1}{4}} \mathbf{F}_{\mu,\beta,\alpha}\left(\frac{2}{1-is}\right)}{\sin(-\pi\mu)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} \\ & + \frac{\pi e^{i\frac{\pi}{2}(-\alpha-\frac{1}{2}+\frac{\mu}{2})} \left(\frac{s+i}{2}\right)^{-\frac{\beta}{2}-\frac{1}{4}+\frac{\mu}{2}} \left(\frac{s-i}{2}\right)^{\frac{\beta}{2}+\frac{1}{4}} \mathbf{F}_{-\mu,\beta,\alpha}\left(\frac{2}{1-is}\right)}{\sin(\pi\mu)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}, \quad s > 0 \end{aligned} \quad (4.55)$$

$$\begin{aligned} & \frac{\pi e^{i\frac{\pi}{2}(\alpha+\frac{1}{2}+\frac{\mu}{2})} \left(\frac{-s-i}{2}\right)^{-\frac{\beta}{2}-\frac{1}{4}-\frac{\mu}{2}} \left(\frac{-s+i}{2}\right)^{\frac{\beta}{2}+\frac{1}{4}} \mathbf{F}_{\mu,\beta,\alpha}\left(\frac{2}{1-is}\right)}{\sin(-\pi\mu)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} \\ & + \frac{\pi e^{i\frac{\pi}{2}(\alpha+\frac{1}{2}-\frac{\mu}{2})} \left(\frac{-s-i}{2}\right)^{-\frac{\beta}{2}-\frac{1}{4}+\frac{\mu}{2}} \left(\frac{-s+i}{2}\right)^{\frac{\beta}{2}+\frac{1}{4}} \mathbf{F}_{-\mu,\beta,\alpha}\left(\frac{2}{1-is}\right)}{\sin(\pi\mu)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}. \quad s < 0 \end{aligned} \quad (4.56)$$

Inserting $s = \sinh(-r) = -\sinh r$ into (4.56), and using uniqueness of analytic continuation we get

$$\begin{aligned} p_{\alpha,\beta,\mu}^{\text{dS}}(r) & := \left(\frac{1-i\sinh r}{2}\right)^{\frac{\alpha}{2}+\frac{1}{4}} \left(\frac{1+i\sinh r}{2}\right)^{\frac{\beta}{2}+\frac{1}{4}} \mathbf{F}_{\alpha,\beta,\mu}\left(\frac{1-i\sinh r}{2}\right) \\ & = \frac{\pi e^{i\frac{\pi}{2}(\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\sin(-\pi\mu)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ & \quad + \frac{\pi e^{i\frac{\pi}{2}(\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\sin(\pi\mu)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)} Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r). \end{aligned} \quad (4.57)$$

By replacing α to $-\alpha$ we obtain the second identity

$$\begin{aligned} p_{-\alpha,\beta,\mu}^{\text{dS}}(r) & = \frac{\pi e^{i\frac{\pi}{2}(-\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\sin(-\pi\mu)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ & \quad + \frac{\pi e^{i\frac{\pi}{2}(-\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\sin(\pi\mu)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)} Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r). \end{aligned} \quad (4.58)$$

We can rewrite them via

$$\begin{aligned} \begin{bmatrix} p_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ p_{-\alpha,\beta,\mu}^{\text{dS}}(r) \end{bmatrix} & = \frac{\pi}{\sin(\pi\mu)} \begin{bmatrix} -\frac{e^{i\frac{\pi}{2}(\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} & \frac{e^{i\frac{\pi}{2}(\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)} \\ -\frac{e^{i\frac{\pi}{2}(-\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)} & \frac{e^{i\frac{\pi}{2}(-\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)} \end{bmatrix} \\ & \quad \times \begin{bmatrix} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r) \end{bmatrix}. \end{aligned} \quad (4.59)$$

We evaluate

$$\det \begin{bmatrix} -\frac{e^{i\frac{\pi}{2}(\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} & \frac{e^{i\frac{\pi}{2}(\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)} \\ -\frac{e^{i\frac{\pi}{2}(-\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)} & \frac{e^{i\frac{\pi}{2}(-\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)} \end{bmatrix} = -i \frac{\sin(\pi\mu)\sin(\pi\alpha)}{\pi^2} \quad (4.60)$$

Therefore we have

$$\begin{aligned} \begin{bmatrix} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r) \end{bmatrix} &= \frac{-i\pi}{\sin(\pi\alpha)} \begin{bmatrix} \frac{e^{\frac{i\pi}{2}(-\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1-\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha-\beta+\mu}{2})} & -\frac{e^{\frac{i\pi}{2}(\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1+\alpha-\beta+\mu}{2})} \\ \frac{e^{\frac{i\pi}{2}(-\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1-\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha-\beta-\mu}{2})} & -\frac{e^{\frac{i\pi}{2}(\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1+\alpha-\beta-\mu}{2})} \end{bmatrix} \\ &\times \begin{bmatrix} p_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ p_{-\alpha,\beta,\mu}^{\text{dS}}(r) \end{bmatrix} \end{aligned} \quad (4.61)$$

In (4.59), we exchange α and β , and replace r with $-r$, and we obtain

$$\begin{aligned} \begin{bmatrix} p_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ p_{-\beta,\alpha,\mu}^{\text{dS}}(-r) \end{bmatrix} &= \frac{\pi}{\sin(\pi\mu)} \begin{bmatrix} -\frac{e^{\frac{i\pi}{2}(\beta+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta-\mu}{2})} & \frac{e^{\frac{i\pi}{2}(\beta-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \\ -\frac{e^{\frac{i\pi}{2}(-\beta+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta-\mu}{2})} & \frac{e^{\frac{i\pi}{2}(-\beta-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \end{bmatrix} \\ &\times \begin{bmatrix} Q_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ Q_{\alpha,\beta,-\mu}^{\text{dS}}(r) \end{bmatrix}. \end{aligned} \quad (4.62)$$

Using (C.12), we know that

$$\begin{aligned} \begin{bmatrix} p_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ p_{-\alpha,\beta,\mu}^{\text{dS}}(r) \end{bmatrix} &= \frac{\pi}{\sin(\pi\beta)} \begin{bmatrix} -\frac{1}{\Gamma(\frac{1+\alpha-\beta-\mu}{2})\Gamma(\frac{1+\alpha-\beta+\mu}{2})} & \frac{1}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1+\alpha+\beta+\mu}{2})} \\ -\frac{1}{\Gamma(\frac{1-\alpha-\beta-\mu}{2})\Gamma(\frac{1-\alpha-\beta+\mu}{2})} & \frac{1}{\Gamma(\frac{1-\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \end{bmatrix} \\ &\times \begin{bmatrix} p_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ p_{-\beta,\alpha,\mu}^{\text{dS}}(-r) \end{bmatrix}. \end{aligned} \quad (4.63)$$

Thus we obtained the connection formula

$$\begin{aligned} \begin{bmatrix} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r) \end{bmatrix} &= \frac{-i\pi^3}{\sin(\pi\alpha)\sin(\pi\beta)\sin(\pi\mu)} \begin{bmatrix} \frac{e^{\frac{i\pi}{2}(-\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1-\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha-\beta+\mu}{2})} & -\frac{e^{\frac{i\pi}{2}(\alpha-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1+\alpha-\beta+\mu}{2})} \\ \frac{e^{\frac{i\pi}{2}(-\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1-\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha-\beta-\mu}{2})} & -\frac{e^{\frac{i\pi}{2}(\alpha+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1+\alpha-\beta-\mu}{2})} \end{bmatrix} \\ &\times \begin{bmatrix} -\frac{1}{\Gamma(\frac{1+\alpha-\beta-\mu}{2})\Gamma(\frac{1+\alpha-\beta+\mu}{2})} & \frac{1}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1+\alpha+\beta+\mu}{2})} \\ -\frac{1}{\Gamma(\frac{1-\alpha-\beta-\mu}{2})\Gamma(\frac{1-\alpha-\beta+\mu}{2})} & \frac{1}{\Gamma(\frac{1-\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \end{bmatrix} \\ &\times \begin{bmatrix} \frac{e^{\frac{i\pi}{2}(\beta+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta-\mu}{2})} & \frac{e^{\frac{i\pi}{2}(\beta-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \\ \frac{e^{\frac{i\pi}{2}(-\beta+\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta-\mu}{2})\Gamma(\frac{1-\alpha+\beta-\mu}{2})} & \frac{e^{\frac{i\pi}{2}(-\beta-\frac{\mu}{2}+\frac{1}{2})}}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1-\alpha+\beta+\mu}{2})} \end{bmatrix} \begin{bmatrix} Q_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ Q_{\alpha,\beta,-\mu}^{\text{dS}}(r) \end{bmatrix}. \end{aligned} \quad (4.64)$$

The result of multiplication after simplification with the reflection formula for the gamma functions is

$$\begin{aligned} &\begin{bmatrix} \frac{-e^{-\frac{i\pi\mu}{2}}\cos(\pi\alpha)+e^{\frac{i\pi\mu}{2}}\cos(\pi\beta)}{\pi} & \frac{2\pi}{\Gamma(\frac{1+\alpha+\beta+\mu}{2})\Gamma(\frac{1+\alpha-\beta+\mu}{2})\Gamma(\frac{1-\alpha-\beta+\mu}{2})} \\ -2\pi & \frac{e^{-\frac{i\pi\mu}{2}}\cos(\pi\beta)+e^{\frac{i\pi\mu}{2}}\cos(\pi\alpha)}{\pi} \end{bmatrix} \\ &\times \begin{bmatrix} Q_{\alpha,\beta,\mu}^{\text{dS}}(r) \\ Q_{\alpha,\beta,-\mu}^{\text{dS}}(r) \end{bmatrix} \frac{-\pi}{\sin(\pi\mu)} = \begin{bmatrix} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) \\ Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r) \end{bmatrix}. \end{aligned} \quad (4.65)$$

Using

$$\mathcal{W}(Q_{\alpha,\beta,\mu}^{\text{dS}}(r), Q_{\alpha,\beta,-\mu}^{\text{dS}}(r)) = \frac{\sin \pi\mu}{\pi} \quad (4.66)$$

we obtain

$$\mathcal{W}(Q_{\beta,\alpha,\mu}^{\text{dS}}(-r), Q_{\alpha,\beta,\mu}^{\text{dS}}(r)) = \frac{2\pi}{\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)} \quad (4.67)$$

$$\mathcal{W}(Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r), Q_{\alpha,\beta,\mu}^{\text{dS}}(r)) = \frac{e^{i\pi\frac{\mu}{2}} \cos \pi\beta + e^{-i\pi\frac{\mu}{2}} \cos \pi\alpha}{\pi}. \quad (4.68)$$

The L^2 integrable condition at both endpoints is $\text{Re}(\mu) > 0$ for both $Q_{\alpha,\beta,\mu}^{\text{dS}}(r)$, and $Q_{\beta,\alpha,\mu}^{\text{dS}}(-r)$. Therefore, for $\text{Re} \mu > 0$, it is a unique candidate for the resolvent. Using the Schur Test we see that the integral kernel (4.51) defines a bounded operator.

From the singularities of the Gamma function we obtain

$$\sigma_{\text{d}}(L_{\alpha}^{\text{dS}}) = \left\{ -\left(n + \frac{1+\alpha+\beta}{2}\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(n + \frac{1+\alpha+\beta}{2}\right) < 0 \right\} \quad (4.69)$$

$$\cup \left\{ -\left(n + \frac{1+\alpha-\beta}{2}\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(n + \frac{1+\alpha-\beta}{2}\right) < 0 \right\} \quad (4.70)$$

$$\cup \left\{ -\left(n + \frac{1-\alpha+\beta}{2}\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(n + \frac{1-\alpha+\beta}{2}\right) < 0 \right\} \quad (4.71)$$

$$\cup \left\{ -\left(n + \frac{1-\alpha-\beta}{2}\right)^2 \mid n \in \mathbb{N}_0, \quad \text{Re}\left(n + \frac{1-\alpha-\beta}{2}\right) < 0 \right\}. \quad (4.72)$$

If $\text{Re}(\alpha + \beta) \geq 0$ and $\text{Re}(\alpha - \beta) \geq 0$, then (4.69) and (4.70) are empty and we obtain (4.52). \square

5 Hypergeometric Hamiltonians of the second kind

We transform the hypergeometric equation in a different way:

$$\begin{aligned} & -4z^{1+\frac{\alpha}{2}}(1-z)^{1+\frac{\beta}{2}} \mathcal{F}\left(\frac{\alpha+\beta+\mu+1}{2}, \frac{\alpha+\beta-\mu+1}{2}; 1+\alpha; z, \partial_z\right) z^{-\frac{\alpha}{2}}(1-z)^{-\frac{\beta}{2}} \\ &= -4z^2(z-1)^2 \mathcal{P} \left(\begin{array}{ccc|c} 0 & 1 & \infty & \\ \frac{\alpha}{2} & \frac{\beta}{2} & \frac{\mu}{2} + \frac{1}{2} & z, \partial_z \\ -\frac{\alpha}{2} & -\frac{\beta}{2} & -\frac{\mu}{2} + \frac{1}{2} & \end{array} \right) \\ &= -4z^2(1-z)^2 \left(\partial_z^2 + \left(\frac{1}{z} - \frac{1}{1-z}\right) \partial_z \right) + \alpha^2(1-z) + \beta^2 z - (\mu^2 - 1)z(1-z). \end{aligned} \quad (5.1)$$

We rearrange the terms in (5.1) containing α and β as follows:

$$\alpha^2(1-z) + \beta^2 z = \delta + \kappa(1-2z) \quad (5.2)$$

$$= \delta + \tau i(2z-1), \quad (5.3)$$

where δ, κ, τ are defined in (4.5). Thus if we set

$$K_{\kappa,\mu} := -4z^2(1-z)^2 \left(\partial_z^2 + \left(\frac{1}{z} - \frac{1}{1-z}\right) \partial_z \right) + \kappa(1-2z) - (\mu^2 - 1)z(1-z) \quad (5.4)$$

and if $F(z)$ solves the hypergeometric equation (2.4), then

$$\left(K_{\kappa,\mu} + \delta\right) z^{\frac{\alpha}{2}} (1-z)^{\frac{\beta}{2}} F(z) = 0. \quad (5.5)$$

We have interpreted (5.1) as the eigenequation of the operator $K_{\kappa,\mu}$ with eigenvalue $-\delta$. Again, we will reinterpret this operator as acting on functions on an interval whose endpoints are singularities of the hypergeometric equations. In each of these cases we perform the Liouville transformation, which yields a 1-dimensional Hamiltonian. We will consider three cases:

1. $z \in \frac{1}{2} + i\mathbb{R}$, which leads to an operator on $L^2]0, \pi[$, which we call the *spherical hypergeometric Hamiltonian of the 2nd kind*;
2. $z \in]-\infty, 0[$, which leads to an operator on $L^2]0, \infty[$ which we call the *hyperbolic hypergeometric Hamiltonian of the 2nd kind*;
3. $z \in]0, 1[$, which leads to an operator on $L^2(\mathbb{R})$, which we call *the deSitterian hypergeometric Hamiltonian of the 2nd kind*.




Figure 3: Hypergeometric Hamiltonian of the second kind on the z -plane.

The spherical Hamiltonian acts on the interval marked with a thick line, the hyperbolic—with a dotted line, and the DeSitterian—with a dashed line.

5.1 Spherical case

For $u \in]0, \pi[$, in (5.1) and (5.3) we set

$$z = \frac{1}{1 - e^{2iu}}, \quad \text{which solves } z' = 2iz(1-z). \quad (5.6)$$

This leads to the Schrödinger equation

$$(K_{\tau,\mu}^s - \delta) \phi(u) = 0, \quad (5.7)$$

where

$$K_{\tau,\mu}^s(u) := -\partial_u^2 + \tau \frac{\cos u}{\sin u} + \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\sin^2 u}. \quad (5.8)$$

This Hamiltonian is known as the *Rosen-Morse Hamiltonian* (see [DW]).

In the case $\tau = 0$ we have the coincidence:

$$L_\alpha^s = K_{0,2\alpha}^s. \quad (5.9)$$

We define for $u \in]0, \frac{\pi}{2}[$

$$\begin{aligned} & \mathbb{Q}_{\alpha,\beta,\mu}^s(u) \\ := & \left(\frac{i}{1 - e^{2iu}} \right)^{\frac{-1-\beta-\mu}{2}} \left(\frac{-i}{1 - e^{-2iu}} \right)^{\frac{\beta}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{-\alpha + \beta + \mu + 1}{2}; \mu + 1; 1 - e^{2iu} \right) \\ = & \left(\frac{i}{1 - e^{2iu}} \right)^{\frac{\alpha}{2}} \left(\frac{-i}{1 - e^{-2iu}} \right)^{\frac{-1-\alpha-\mu}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; \mu + 1; 1 - e^{-2iu} \right). \end{aligned} \quad (5.10)$$

For $u \in [\frac{\pi}{2}, \pi[$ it is continued analytically. It has the asymptotics

$$\mathbb{Q}_{\alpha,\beta,\mu}^s(u) \sim \frac{1}{\Gamma(\mu + 1)} (2u)^{\frac{\mu}{2} + \frac{1}{2}}, \quad u \sim 0. \quad (5.11)$$

Note that

$$\mathbb{Q}_{\alpha,\beta,\mu}^s(u) = \mathbb{Q}_{-\alpha,\beta,\mu}^s(u) = \mathbb{Q}_{\alpha,-\beta,\mu}^s(u), \quad (5.12)$$

$$\mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - u) = \mathbb{Q}_{\beta,-\alpha,\mu}^s(\pi - u) = \mathbb{Q}_{-\beta,\alpha,\mu}^s(\pi - u), \quad (5.13)$$

$$\left(\frac{2}{\cosh r} \right)^{\frac{1}{2}} Q_{\alpha,\beta,\mu}^{\text{dS}}(r) = \mathbb{Q}_{\alpha,\beta,\mu}^s(u), \quad (5.14)$$

$$\left(\frac{2}{\cosh r} \right)^{\frac{1}{2}} Q_{\beta,\alpha,\mu}^{\text{dS}}(-r) = \mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - u), \quad \sinh r = -\cot u. \quad (5.15)$$

Now the following functions solve the eigenvalue problem (5.7):

$$\mathbb{Q}_{\alpha,\beta,\mu}^s(u), \quad \mathbb{Q}_{\alpha,\beta,-\mu}^s(u), \quad \mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - u), \quad \mathbb{Q}_{\beta,\alpha,-\mu}^s(\pi - u). \quad (5.16)$$

The following theorem describes the basic closed realization of $K_{\tau,\mu}^s$:

Theorem 5.1. *For $\text{Re } \mu \geq 2$, $\tau \in \mathbb{C}$, there exists a unique closed operator $K_{\tau,\mu}^s$ in the sense of $L^2]0, \pi[$, which on $C_c^\infty]0, \pi[$ is given by (5.8). The family $\tau, \mu \mapsto K_{\tau,\mu}^s$ is holomorphic and possesses a unique holomorphic extension to $\text{Re } \mu > -2$, except for a singularity at $(\tau, \mu) = (0, -1)$. It has only discrete spectrum:*

$$\sigma(K_{\tau,\mu}^s) = \left\{ -\frac{\tau^2}{(2k + \mu)^2} + \left(k + \frac{\mu}{2}\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2} \right\}. \quad (5.17)$$

Set

$$\alpha := \sqrt{\delta - i\tau}, \quad \beta := \sqrt{\delta + i\tau}. \quad (5.18)$$

(It does not matter which sign of the square root is taken). Outside of the spectrum the resolvent of $K_{\tau,\mu}^s$ is

$$\begin{aligned} \frac{1}{(K_{\tau,\mu}^s - \delta)}(x, y) &= \frac{\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)}{4\pi} \\ &\times \begin{cases} \mathbb{Q}_{\alpha,\beta,\mu}^s(x) \mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - y), & \text{if } 0 < x < y < \pi; \\ \mathbb{Q}_{\alpha,\beta,\mu}^s(y) \mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - x), & \text{if } 0 < y < x < \pi. \end{cases} \end{aligned} \quad (5.19)$$

Proof. The relation $\sinh r = -\cot u$ implies

$$\frac{2}{1 - i \sinh r} = 1 - e^{2iu}, \quad \frac{2}{1 + i \sinh r} = 1 - e^{-2iu}, \quad (5.20)$$

$$\frac{du}{dr} = \sin u = \frac{1}{\cosh r}. \quad (5.21)$$

Therefore, using (4.67) and (4.68), we obtain

$$\begin{aligned} \mathcal{W}(\mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - u), \mathbb{Q}_{\alpha,\beta,\mu}^s(u)) &= 2\mathcal{W}(Q_{\beta,\alpha,\mu}^{\text{dS}}(-r), Q_{\alpha,\beta,\mu}^{\text{dS}}(r)) \\ &= \frac{4\pi}{\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right) \Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)}, \end{aligned} \quad (5.22)$$

$$\begin{aligned} \mathcal{W}(\mathbb{Q}_{\beta,\alpha,-\mu}^s(\pi - u), \mathbb{Q}_{\alpha,\beta,\mu}^s(u)) &= 2\mathcal{W}(Q_{\beta,\alpha,-\mu}^{\text{dS}}(-r), Q_{\alpha,\beta,\mu}^{\text{dS}}(r)) \\ &= \frac{2(e^{i\pi\frac{\mu}{2}} \cos \pi\beta + e^{-i\pi\frac{\mu}{2}} \cos \pi\alpha)}{\pi}. \end{aligned} \quad (5.23)$$

The L^2 integrability condition for $\mathbb{Q}_{\alpha,\beta,\mu}^s(u)$ at 0 and $\mathbb{Q}_{\beta,\alpha,\mu}^s(\pi - u)$ at π is $\text{Re } \mu > -2$. The L^2 norm of this kernel 5.19 is finite. For $\text{Re}(\mu) \geq 2$ it is a unique candidate for the resolvent.

As a byproduct we obtain a proof of Prop. 1.4 about the transmutation identity $L^{\text{dS}} \rightarrow K^s$.

The singularities of the Gamma functions in (5.19) are at

$$1 + \epsilon_1\alpha + \epsilon_2\beta + \mu = -2n, \quad n \in \mathbb{N}_0, \quad (5.24)$$

where $\epsilon_1, \epsilon_2 \in \{1, -1\}$. This implies

$$\alpha^2 = (2n + 1 + \mu)^2 + 2\epsilon_2\beta(2n + 1 + \mu) + \beta^2. \quad (5.25)$$

Hence,

$$\beta = \frac{\epsilon_2\kappa}{2n + 1 + \mu} - \epsilon_2\left(n + \frac{\mu}{2} + \frac{1}{2}\right), \quad (5.26)$$

$$\alpha = -\frac{\epsilon_2\kappa}{2n + 1 + \mu} - \epsilon_2\left(n + \frac{\mu}{2} + \frac{1}{2}\right). \quad (5.27)$$

This shows

$$\delta = \frac{\kappa^2}{(2n + 1 + \mu)^2} + \left(n + \frac{1}{2} + \frac{\mu}{2}\right)^2. \quad (5.28)$$

Replacing κ^2 with $-\tau^2$ we obtain (5.17). \square

For $\mu \in \mathbb{Z}$ we have an additional identity for the \mathbb{Q}^s function, which follows directly from (C.17):

$$\left(\frac{\alpha + \beta - \mu + 1}{2}\right)_\mu \left(\frac{\alpha - \beta - \mu + 1}{2}\right)_\mu \mathbb{Q}_{\alpha,\beta,\mu}^s(u) = \mathbb{Q}_{\alpha,\beta,-\mu}^s(u). \quad (5.29)$$

Using this with $\mu = 1$ we obtain the following unexpected identity:

Theorem 5.2. *For any $\tau \neq 0$, we have $K_{\tau,1}^s = K_{\tau,-1}^s$.*

Proof. Setting $\mu = 1$ in (5.56) we obtain

$$\frac{\alpha^2 - \beta^2}{4} \mathbb{Q}_{\alpha,\beta,1}^s(u) = \mathbb{Q}_{\alpha,\beta,-1}^s(u). \quad (5.30)$$

Setting $\mu = 1$ and $\mu = -1$ in the prefactor of the right hand side of (5.19) we obtain

$$\frac{\Gamma\left(1 - \frac{\alpha+\beta}{2}\right)\Gamma\left(1 + \frac{\alpha-\beta}{2}\right)\Gamma\left(1 - \frac{\alpha-\beta}{2}\right)\Gamma\left(1 + \frac{\alpha+\beta}{2}\right)}{4\pi} = \frac{(\alpha^2 - \beta^2)\pi}{16 \sin \frac{\pi}{2}(\alpha + \beta) \sin \frac{\pi}{2}(\alpha - \beta)}, \quad (5.31)$$

$$\frac{\Gamma\left(-\frac{\alpha+\beta}{2}\right)\Gamma\left(\frac{\alpha-\beta}{2}\right)\Gamma\left(-\frac{\alpha-\beta}{2}\right)\Gamma\left(\frac{\alpha+\beta}{2}\right)}{4\pi} = \frac{\pi}{(\alpha^2 - \beta^2) \sin \frac{\pi}{2}(\alpha + \beta) \sin \frac{\pi}{2}(\alpha - \beta)}. \quad (5.32)$$

We thus obtain

$$\frac{1}{(K_{\tau,1}^s + \delta)} = \frac{1}{(K_{\tau,-1}^s + \delta)}. \quad (5.33)$$

□

Note that (5.19) is ill defined for $(\tau, \mu) = (0, -1)$. Moreover, we know that $\{2 < \operatorname{Re} \mu\} \ni \mu \mapsto L_{\frac{\mu}{2}}^s$ is analytic, and for $\mu \neq -1$.

We have $K_{0,\mu}^s = L_{\frac{\mu}{2}}^s$. Therefore, it is natural to set

$$K_{0,-1}^s := L_{-\frac{1}{2}}^s. \quad (5.34)$$

We know that $L_{-\frac{1}{2}}^s \neq L_{\frac{1}{2}}^s$. Therefore, Thm 5.2 implies that the point $(\tau, \mu) = (0, -1)$ is a singularity of the function $(\tau, \mu) \mapsto K_{\tau,\mu}^s$. See [DeRi], where a similar phenomenon is described for the Whittaker operator.

5.2 Hyperbolic case

For $u \in \mathbb{R}_+$, in (5.1) and (5.2) we set

$$z = \frac{1}{1 - e^{2u}}, \quad \text{which solves } z' = 2z(z - 1). \quad (5.35)$$

This leads to the Schrödinger equation

$$(K_{\kappa,\mu}^h + \delta) \phi(u) = 0, \quad (5.36)$$

where

$$K_{\kappa,\mu}^h := -\partial_u^2 + \kappa \frac{\cosh u}{\sinh u} + \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\sinh^2 u}. \quad (5.37)$$

This Hamiltonian in [DW] is called the *Eckart Hamiltonian*.

In the case $\kappa = 0$ we have the coincidence

$$L_\alpha^h = K_{0,2\alpha}^h. \quad (5.38)$$

We define

$$\begin{aligned} \mathbb{P}_{\alpha,\beta,\mu}^h(u) &:= (e^{2u} - 1)^{-\frac{\alpha}{2}} (1 - e^{-2u})^{-\frac{\beta}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha + \beta - \mu + 1}{2}; \alpha + 1; \frac{1}{1 - e^{2u}} \right) \\ &= (e^{2u} - 1)^{-\frac{\alpha}{2}} (1 - e^{-2u})^{\frac{1+\alpha+\mu}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; \alpha + 1; e^{-2u} \right), \end{aligned} \quad (5.39)$$

$$\begin{aligned} \mathbb{Q}_{\alpha,\beta,\mu}^h(u) &:= (e^{2u} - 1)^{\frac{1+\beta+\mu}{2}} (1 - e^{-2u})^{-\frac{\beta}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{-\alpha + \beta + \mu + 1}{2}; \mu + 1; 1 - e^{2u} \right) \\ &= (e^{2u} - 1)^{-\frac{\alpha}{2}} (1 - e^{-2u})^{\frac{1+\alpha+\mu}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; \mu + 1; 1 - e^{-2u} \right). \end{aligned} \quad (5.40)$$

Note that

$$\mathbb{P}_{\alpha,\beta,\mu}^h(u) \sim \frac{1}{\Gamma(1+\alpha)} e^{-\alpha u}, \quad u \sim +\infty; \quad (5.41)$$

$$\mathbb{Q}_{\alpha,\beta,\mu}^h(u) \sim \frac{1}{\Gamma(1+\mu)} (2u)^{\frac{\mu}{2}+\frac{1}{2}}, \quad u \sim 0. \quad (5.42)$$

We have

$$\mathbb{P}_{\alpha,\beta,\mu}^h(u) = \mathbb{P}_{\alpha,-\beta,\mu}^h(u) = \mathbb{P}_{\alpha,\beta,-\mu}^h(u), \quad (5.43)$$

$$\mathbb{Q}_{\alpha,\beta,\mu}^h(u) = \mathbb{Q}_{-\alpha,\beta,\mu}^h(u) = \mathbb{Q}_{\alpha,-\beta,\mu}^h(u); \quad (5.44)$$

$$\left(\frac{2}{\sinh r}\right)^{\frac{1}{2}} P_{\alpha,\beta,\mu}^h(r) = \mathbb{P}_{\alpha,\beta,\mu}^h(u), \quad (5.45)$$

$$\left(\frac{2}{\sinh r}\right)^{\frac{1}{2}} Q_{\alpha,\beta,\mu}^h(r) = \mathbb{Q}_{\alpha,\beta,\mu}^h(u), \quad \cosh r = \coth u. \quad (5.46)$$

Now the following functions solve the eigenvalue problem (5.36):

$$\mathbb{P}_{\alpha,\beta,\mu}^h(u), \quad \mathbb{P}_{-\alpha,\beta,\mu}^h(u), \quad \mathbb{Q}_{\alpha,\beta,\mu}^h(u), \quad \mathbb{Q}_{\alpha,\beta,-\mu}^h(u). \quad (5.47)$$

Let us describe the basic closed realization of $K_{\kappa,\mu}^h$ on $L^2(\mathbb{R}_+)$.

Theorem 5.3. *For $\kappa \in \mathbb{C}, \operatorname{Re} \mu \geq 2$ there exists a unique closed operator $K_{\kappa,\mu}^h$ in the sense of $L^2(\mathbb{R}_+)$, which on $C_c^\infty(\mathbb{R}_+)$ is given by (5.37). The family $\kappa, \mu \mapsto K_{\kappa,\mu}^h$ is holomorphic and possesses a unique holomorphic extension to $\operatorname{Re} \mu > -2$, $(\kappa, \mu) \neq (0, 1)$. Its discrete spectrum and spectrum are*

$$\sigma_d(K_{\kappa,\mu}^h) = \left\{ -\frac{\kappa^2}{(2k+\mu)^2} - \left(k + \frac{\mu}{2}\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \operatorname{Re} \left(\frac{\kappa}{2k+\mu} + k + \frac{\mu}{2}\right) < 0 \right\}, \quad (5.48)$$

$$\sigma(K_{\kappa,\mu}^h) = [0, +\infty[\cup \sigma_d(K_{\kappa,\mu}^h). \quad (5.49)$$

Set

$$\alpha := \sqrt{\delta + \kappa}, \quad \operatorname{Re} \alpha > 0, \quad \beta := \sqrt{\delta - \kappa} \quad (5.50)$$

(The choice of the square root for β does not matter). Outside of its spectrum the resolvent of $K_{\kappa,\mu}^h$ is

$$\begin{aligned} \frac{1}{(K_{\kappa,\mu}^h + \delta)}(x, y) &= \frac{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}{2} \\ &\times \begin{cases} \mathbb{P}_{\alpha,\beta,\mu}^h(x) \mathbb{Q}_{\alpha,\beta,\mu}^h(y) & \text{if } 0 < y < x < \infty, \\ \mathbb{P}_{\alpha,\beta,\mu}^h(y) \mathbb{Q}_{\alpha,\beta,\mu}^h(x) & \text{if } 0 < x < y < \infty. \end{cases} \end{aligned} \quad (5.51)$$

Proof. The relation $\cosh r = \coth u$ implies

$$\sinh^2 \frac{r}{2} = \frac{1}{e^{2u} - 1}, \quad \cosh^2 \frac{r}{2} = \frac{1}{e^{-2u} - 1}; \quad (5.52)$$

$$\frac{du}{dr} = \frac{1}{\sinh r} = \sinh u, \quad (5.53)$$

Therefore,

$$\begin{aligned} \mathcal{W}(\mathbb{Q}_{\alpha,\beta,\mu}^h, \mathbb{P}_{\alpha,\beta,\mu}^h) &= 2\mathcal{W}(Q_{\alpha,\beta,\mu}^h, P_{\alpha,\beta,\mu}^h) \\ &= \frac{2}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}. \end{aligned} \quad (5.54)$$

The L^2 integrability conditions at endpoints are $\operatorname{Re} \alpha > 0$, and $\operatorname{Re} \mu > -2$ for $\mathbb{P}_{\alpha,\beta,\mu}^{\text{h}}(r)$ and $\mathbb{Q}_{\alpha,\beta,\mu}^{\text{h}}(r)$, respectively. For such parameters the integral kernel (5.51) defines a bounded operator. For $\operatorname{Re} \alpha > 0$, and $\operatorname{Re} \mu \geq 2$, it is the unique candidate for the resolvent.

As a byproduct we obtain a proof of Prop. 1.5 about the transmutation identity $L^{\text{h}} \rightarrow K^{\text{h}}$.

The determination of the discrete spectrum follows by similar arguments as for Thm 5.2. \square

For $\mu \in \mathbb{Z}$ we have an additional identity for the \mathbb{Q}^{h} function, which follows directly from (C.17):

$$\left(\frac{\alpha + \beta - \mu + 1}{2}\right)_{\mu} \left(\frac{\alpha - \beta - \mu + 1}{2}\right)_{\mu} \mathbb{Q}_{\alpha,\beta,\mu}^{\text{h}}(u) = \mathbb{Q}_{\alpha,\beta,-\mu}^{\text{h}}(u). \quad (5.55)$$

Using this with $\mu = 1$ we obtain

Theorem 5.4. *For any $\kappa \neq 0$, we have $K_{\kappa,1}^{\text{h}} = K_{\kappa,-1}^{\text{h}}$.*

Proof. Setting $\mu = 1$ in (5.56) we obtain

$$\frac{\alpha^2 - \beta^2}{4} \mathbb{Q}_{\alpha,\beta,1}^{\text{h}}(u) = \mathbb{Q}_{\alpha,\beta,-1}^{\text{h}}(u). \quad (5.56)$$

Moreover,

$$\Gamma\left(1 - \frac{\alpha + \beta}{2}\right) \Gamma\left(1 + \frac{\alpha - \beta}{2}\right) = \frac{(\alpha^2 - \beta^2)}{4} \Gamma\left(\frac{\alpha + \beta}{2}\right) \Gamma\left(\frac{\alpha - \beta}{2}\right), \quad (5.57)$$

We thus obtain

$$\frac{1}{(K_{\kappa,1}^{\text{h}} + \delta)} = \frac{1}{(K_{\kappa,-1}^{\text{h}} + \delta)}. \quad (5.58)$$

\square

Similarly as in the spherical case, (5.51) is ill defined for $(\kappa, \mu) = (0, -1)$. Moreover, we know that $\{2 < \operatorname{Re} \mu\} \ni \mu \mapsto L_{\frac{\mu}{2}}^{\text{h}}$ is analytic, and for $\mu \neq -1$ we have $K_{0,\mu}^{\text{h}} = L_{\frac{\mu}{2}}^{\text{h}}$. Therefore, it is natural to set

$$K_{0,-1}^{\text{h}} := L_{-\frac{1}{2}}^{\text{h}}. \quad (5.59)$$

We know that $L_{-\frac{1}{2}}^{\text{h}} \neq L_{\frac{1}{2}}^{\text{h}}$. Therefore, Thm 5.4 implies that the point $(\kappa, \mu) = (0, -1)$ is a singularity of the function $(\kappa, \mu) \mapsto K_{\kappa,\mu}^{\text{h}}$.

5.3 DeSitterian case

For $u \in \mathbb{R}$, in (5.1) and (5.2) we set

$$z = \frac{1}{1 + e^{2u}}, \quad \text{which solves } z' = 2z(z - 1). \quad (5.60)$$

This leads to the Schrödinger equation

$$(K_{\kappa,\mu}^{\text{dS}} + \delta) \phi(u) = 0, \quad (5.61)$$

where

$$K_{\kappa,\mu}^{\text{dS}} := -\partial_u^2 - \kappa \frac{\sinh u}{\cosh u} - \left(\frac{\mu^2}{4} - \frac{1}{4}\right) \frac{1}{\cosh^2 u}. \quad (5.62)$$

In [DW] it is called the *Manning-Rosen Hamiltonian*.

In the case $\kappa = 0$ we have the coincidence:

$$L_\alpha^{\text{dS}} = K_{0,2\alpha}^{\text{dS}}. \quad (5.63)$$

We define

$$\begin{aligned} \mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u) &:= (1 + e^{2u})^{-\frac{\alpha}{2}} (1 + e^{-2u})^{-\frac{\beta}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha + \beta - \mu + 1}{2}; \alpha + 1; \frac{1}{1 + e^{2u}} \right) \\ &= (1 + e^{2u})^{-\frac{\alpha}{2}} (1 + e^{-2u})^{\frac{1+\alpha+\mu}{2}} \mathbf{F} \left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; \alpha + 1; -e^{-2u} \right). \end{aligned} \quad (5.64)$$

We have the asymptotics

$$\mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u) \sim \frac{1}{\Gamma(1 + \alpha)} e^{-\alpha u}, \quad u \sim +\infty. \quad (5.65)$$

Note that

$$\mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u) = \mathbb{P}_{\alpha,-\beta,\mu}^{\text{dS}}(u) = \mathbb{P}_{\alpha,\beta,-\mu}^{\text{dS}}(u), \quad (5.66)$$

$$\left(\frac{2}{\sin r} \right)^{\frac{1}{2}} P_{\alpha,\beta,\mu}^{\text{dS}}(r) = \mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u), \quad \cos r = \tanh u. \quad (5.67)$$

Now the following functions solve the eigenvalue problem (5.61):

$$\mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u), \quad \mathbb{P}_{-\alpha,\beta,\mu}^{\text{dS}}(u), \quad \mathbb{P}_{\beta,\alpha,\mu}^{\text{dS}}(-u), \quad \mathbb{P}_{-\beta,\alpha,\mu}^{\text{dS}}(-u). \quad (5.68)$$

We will find all closed realization of $K_{\kappa,\mu}^{\text{dS}}$ in the sense of $L^2(\mathbb{R})$.

Theorem 5.5. *For any $\kappa, \mu \in \mathbb{C}$ there exists a unique closed operator $K_{\kappa,\mu}^{\text{dS}}$ in the sense of $L^2(\mathbb{R})$ that on $C_c^\infty(\mathbb{R})$ is given by (5.62). The function $\mathbb{C}^2 \ni (\kappa, \mu) \mapsto K_{\kappa,\mu}^{\text{dS}}$ is holomorphic. The discrete spectrum and spectrum of $K_{\kappa,\mu}^{\text{dS}}$ are*

$$\begin{aligned} \sigma_{\text{d}}(K_{\kappa,\mu}^{\text{dS}}) &= \left\{ -\frac{\kappa^2}{(2k + \mu)^2} - \left(k + \frac{\mu}{2}\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < -\left| \operatorname{Re} \frac{\kappa}{2k + \mu} \right| - \frac{\mu}{2} \right\} \\ &\cup \left\{ -\frac{\kappa^2}{(2k - \mu)^2} - \left(k - \frac{\mu}{2}\right)^2 \mid k \in \mathbb{N}_0 + \frac{1}{2}, \quad k < -\left| \operatorname{Re} \frac{\kappa}{2k - \mu} \right| + \frac{\mu}{2} \right\}, \end{aligned} \quad (5.69)$$

$$\sigma(K_{\kappa,\mu}^{\text{dS}}) = [\kappa, +\infty[\cup [-\kappa, +\infty[\cup \sigma_{\text{d}}(K_{\kappa,\mu}^{\text{dS}}). \quad (5.70)$$

Here, for $z \in \mathbb{C}$ we use the notation $[z, +\infty[:= \{z + t \mid t \in [0, +\infty[\}$.

Set

$$\alpha := \sqrt{\delta + \kappa}, \quad \operatorname{Re} \alpha > 0; \quad \beta := \sqrt{\delta - \kappa}, \quad \operatorname{Re} \beta > 0. \quad (5.71)$$

Outside of its spectrum, the resolvent of $K_{\kappa,\mu}^{\text{dS}}$ is

$$\begin{aligned} \frac{1}{(K_{\kappa,\mu}^{\text{dS}} + \delta)}(x, y) &= \frac{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right) \Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)}{2} \\ &\times \begin{cases} \mathbb{P}_{\beta,\alpha,\mu}^{\text{dS}}(x) \mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(y), & \text{if } -\infty < x < y < \infty; \\ \mathbb{P}_{\beta,\alpha,\mu}^{\text{dS}}(y) \mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(x), & \text{if } -\infty < y < x < \infty. \end{cases} \end{aligned} \quad (5.72)$$

We have $K_{\kappa,\mu}^{\text{dS}} = K_{\kappa,-\mu}^{\text{dS}}$

Proof. The relation $\cos r = \tanh u$ implies

$$\sin^2 \frac{r}{2} = \frac{1}{1 + e^{2u}}, \quad \cos^2 \frac{r}{2} = \frac{1}{1 + e^{-2u}}; \quad (5.73)$$

$$\frac{du}{dr} = \frac{1}{\sin r} = \cosh u. \quad (5.74)$$

This yields (5.67). Using (5.74) and (5.67), and then (4.18), we obtain

$$\begin{aligned} \mathcal{W}(\mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(u), \mathbb{P}_{\beta,\alpha,\mu}^{\text{dS}}(-u)) &= 2\mathcal{W}(P_{\alpha,\beta,\mu}^{\text{s}}(r), P_{\beta,\alpha,\mu}^{\text{s}}(\pi - r)) \\ &= \frac{2}{\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)}. \end{aligned} \quad (5.75)$$

The L^2 integrability condition at $+\infty$ of $\mathbb{P}_{\alpha,\beta,\mu}(u)$ is $\text{Re}(\alpha) > 0$, and at $-\infty$ of $\mathbb{P}_{\beta,\alpha,\mu}(-u)$ is $\text{Re}(\beta) > 0$. For such parameters, the integral kernel (5.72) defines a bounded operator and is a unique candidate for the resolvent.

As a byproduct we obtain a proof of Prop. 1.4 about the transmutation identity $L^{\text{s}} \rightarrow L^{\text{dS}}$.

The determination of the discrete spectrum is similar as in the hyperbolic case. \square

Proof of Prop. 1.3. The change of variables

$$1 + e^{2w} = e^{2u} \quad (5.76)$$

implies

$$-e^{2w} = 1 - e^{2u}, \quad -e^{-2w} = \frac{1}{1 - e^{2u}}; \quad (5.77)$$

$$\frac{dw}{du} = \frac{1}{1 - e^{2u}} = 1 + e^{-2w}, \quad (5.78)$$

$$\mathbb{P}_{\alpha,\beta,\mu}^{\text{dS}}(w) = (1 - e^{-2u})^{-\frac{1}{2}} \mathbb{P}_{\alpha,\mu,\beta}^{\text{h}}(u), \quad (5.79)$$

$$\mathbb{P}_{\beta,\alpha,\mu}^{\text{dS}}(-w) = (1 - e^{-2u})^{-\frac{1}{2}} \mathbb{Q}_{\alpha,\mu,\beta}^{\text{h}}(u). \quad (5.80)$$

We redefine parameters

$$\delta' := \frac{1}{2}(\alpha^2 - \mu^2), \quad \kappa' := \frac{1}{2}(\alpha^2 + \mu^2). \quad (5.81)$$

We obtain the transmutation identity

$$(1 - e^{-2u})^{-\frac{1}{2}} \frac{1}{(K_{\kappa,\mu}^{\text{h}} + \delta)}(u, u')(1 - e^{-2u'})^{-\frac{1}{2}} = \frac{1}{(K_{\kappa',\beta}^{\text{dS}} + \delta')}(w, w'). \quad (5.82)$$

\square

6 The Laplacian on an interval, halfline and line

The Laplacians on $]0, \pi[$, \mathbb{R}_+ and \mathbb{R} with the Dirichlet or Neumann boundary conditions at endpoints belong to the most widely used operators. Their Green functions can be easily computed in terms of elementary functions, without using hypergeometric functions. In this section we will check that they are special cases of hypergeometric Hamiltonians. We will see that this coincidence is related to various identities for hypergeometric functions from Appendix C.6.

6.1 Laplacian on an interval

Consider the Laplacian $-\partial_x^2$ on the interval $]0, \pi[$. The Dirichlet and Neumann boundary condition will be denoted D and N resp. Putting them at both 0 and π leads to 4 operators on $L^2]0, \pi[$. They are special cases of the spherical hypergeometric Hamiltonian of the first kind:

$$L_{\text{DD}} := L_{\frac{1}{2}, \frac{1}{2}}^{\text{s}} = L_{\frac{1}{2}}^{\text{s}}, \quad (6.1)$$

$$L_{\text{ND}} := L_{-\frac{1}{2}, \frac{1}{2}}^{\text{s}}, \quad (6.2)$$

$$L_{\text{DN}} := L_{\frac{1}{2}, -\frac{1}{2}}^{\text{s}}, \quad (6.3)$$

$$L_{\text{NN}} := L_{-\frac{1}{2}, -\frac{1}{2}}^{\text{s}} = L_{-\frac{1}{2}}^{\text{s}}. \quad (6.4)$$

Resolvents of these operators can be computed in terms of elementary functions. Indeed, the following functions solve

$$(-\partial_x^2 + k^2)\phi(x) = 0 \quad (6.5)$$

and satisfy the Dirichlet/Neumann boundary conditions at 0, resp. at π :

$$\text{Dirichlet: } \sinh kx, \quad \sinh k(\pi - x); \quad (6.6)$$

$$\text{Neumann: } \cosh kx, \quad \cosh k(\pi - x). \quad (6.7)$$

They have the Wronskians:

$$\mathcal{W}(\sinh k(\pi - x), \sinh kx) = k \sinh \pi k, \quad (6.8)$$

$$\mathcal{W}(\cosh k(\pi - x), \sinh kx) = k \cosh \pi k, \quad (6.9)$$

$$\mathcal{W}(\sinh k(\pi - x), \cosh kx) = k \cosh \pi k, \quad (6.10)$$

$$\mathcal{W}(\cosh k(\pi - x), \cosh kx) = k \sinh \pi k. \quad (6.11)$$

By the usual methods, we compute the spectra of operators (6.1)–(6.4), and for k^2 outside of the spectra their Green functions :

$$\begin{aligned} \sigma(L_{\text{DD}}) &= \{n^2 \mid n \in \mathbb{N}_0\}, \\ \frac{1}{L_{\text{DD}} + k^2}(x, y) &= \frac{1}{k \sinh \pi k} \begin{cases} \sinh kx \sinh k(\pi - y), & \text{if } 0 < x < y < \pi, \\ \sinh ky \sinh k(\pi - x), & \text{if } 0 < y < x < \pi; \end{cases} \end{aligned} \quad (6.12)$$

$$\begin{aligned} \sigma(L_{\text{ND}}) &= \{(n + \frac{1}{2})^2 \mid n \in \mathbb{N}_0\}, \\ \frac{1}{L_{\text{ND}} + k^2}(x, y) &= \frac{1}{k \cosh \pi k} \begin{cases} \cosh kx \sinh k(\pi - y), & \text{if } 0 < x < y < \pi, \\ \cosh ky \sinh k(\pi - x), & \text{if } 0 < y < x < \pi; \end{cases} \end{aligned} \quad (6.13)$$

$$\begin{aligned} \sigma(L_{\text{DN}}) &= \{(n + \frac{1}{2})^2 \mid n \in \mathbb{N}_0\}, \\ \frac{1}{L_{\text{DN}} + k^2}(x, y) &= \frac{1}{k \cosh \pi k} \begin{cases} \sinh kx \cosh k(\pi - y), & \text{if } 0 < x < y < \pi, \\ \sinh ky \cosh k(\pi - x), & \text{if } 0 < y < x < \pi; \end{cases} \end{aligned} \quad (6.14)$$

$$\begin{aligned} \sigma(L_{\text{NN}}) &= \{n^2 \mid n \in \mathbb{N}\}, \\ \frac{1}{L_{\text{NN}} + k^2}(x, y) &= \frac{1}{k \sinh \pi k} \begin{cases} \cosh kx \cosh k(\pi - y), & \text{if } 0 < x < y < \pi, \\ \cosh ky \cosh k(\pi - x), & \text{if } 0 < y < x < \pi. \end{cases} \end{aligned} \quad (6.15)$$

Let us check that (6.12)–(6.15) agree with the more general formula (4.15) involving the hypergeometric function and the Gamma function. We identify $k = \frac{i\mu}{2}$. By (C.31) and (C.32) we obtain

$$P_{\frac{1}{2}, \pm \frac{1}{2}, \mu}^s(x) = \frac{\sinh kx}{k\sqrt{\pi}}, \quad (6.16)$$

$$P_{-\frac{1}{2}, \pm \frac{1}{2}, \mu}^s(x) = \frac{\cosh kx}{\sqrt{\pi}}, \quad (6.17)$$

Finally, we use

$$\Gamma(1 + ik) \Gamma(1 - ik) = \frac{k\pi}{\sinh k\pi}, \quad (6.18)$$

$$\Gamma\left(\frac{1}{2} + ik\right) \Gamma\left(\frac{1}{2} - ik\right) = \frac{\pi}{\cosh \pi k}, \quad (6.19)$$

$$\Gamma(ik) \Gamma(-ik) = \frac{\pi}{k \sinh k\pi}. \quad (6.20)$$

6.2 Laplacian on the halfline

Consider the Laplacian $-\partial_x^2$ on the half-line \mathbb{R}_+ . Setting the Dirichlet and Neumann boundary conditions at 0 we obtain 2 operators on $L^2(\mathbb{R}_+)$, which are special cases of the hyperbolic Gegenbauer Hamiltonian:

$$L_D := L_{\frac{1}{2}, \frac{1}{2}}^h = L_{\frac{1}{2}, -\frac{1}{2}}^h = L_{\frac{1}{2}}^h, \quad (6.21)$$

$$L_N := L_{-\frac{1}{2}, \frac{1}{2}}^h = L_{-\frac{1}{2}, -\frac{1}{2}}^h = L_{-\frac{1}{2}}^h. \quad (6.22)$$

Let us compute their resolvents. The following functions solve

$$(-\partial_x^2 + k^2)\phi(x) = 0 \quad (6.23)$$

and satisfy the Dirichlet/Neumann boundary conditions at 0 and decay at $+\infty$:

$$\text{Dirichlet: } \sinh kx; \quad (6.24)$$

$$\text{Neumann: } \cosh kx; \quad (6.25)$$

$$\text{decaying at } +\infty: e^{-kx}, \quad \text{Re } k > 0. \quad (6.26)$$

They have the Wronskians:

$$\mathcal{W}(e^{-kx}, \sinh kx) = k, \quad (6.27)$$

$$\mathcal{W}(e^{-kx}, \cosh kx) = k. \quad (6.28)$$

Now for $\text{Re } k > 0$,

$$(L_D + k^2)^{-1}(x, y) = \frac{1}{k} \begin{cases} \sinh kx e^{-ky}, & \text{if } 0 < x < y, \\ \sinh ky e^{-kx}, & \text{if } 0 < y < x; \end{cases} \quad (6.29)$$

$$(L_N + k^2)^{-1}(x, y) = \frac{1}{k} \begin{cases} \cosh kx e^{-ky}, & \text{if } 0 < x < y, \\ \cosh ky e^{-kx}, & \text{if } 0 < y < x. \end{cases} \quad (6.30)$$

To check that (6.29) and (6.30) agree with (4.32), identify $k = \frac{\mu}{2}$. By (C.33), (C.34) and (C.35), we have

$$P_{\frac{1}{2}, \pm \frac{1}{2}, \pm \mu}^{\text{h}}(x) = \frac{\sinh kx}{k\sqrt{\pi}}, \quad (6.31)$$

$$P_{-\frac{1}{2}, \pm \frac{1}{2}, \pm \mu}^{\text{h}}(x) = \frac{\cosh kx}{\sqrt{\pi}}, \quad (6.32)$$

$$Q_{\pm \frac{1}{2}, \pm \frac{1}{2}, \mu}^{\text{h}}(x) = \frac{2^{2k}}{\Gamma(1+2k)} e^{-kx}. \quad (6.33)$$

Finally, use

$$\Gamma(k)\Gamma\left(k + \frac{1}{2}\right) = \frac{2^{-2k}\sqrt{\pi}}{k}\Gamma(2k+1), \quad (6.34)$$

$$\Gamma\left(k + \frac{1}{2}\right)\Gamma(k+1) = 2^{-2k}\sqrt{\pi}\Gamma(2k+1). \quad (6.35)$$

6.3 Laplacian on the line

Consider the Laplacian $-\partial_x^2$ on the line \mathbb{R} , denoted L . It is a special case of the deSitterian hypergeometric Hamiltonian of the first kind:

$$L := L_{\pm \frac{1}{2}, \pm \frac{1}{2}}^{\text{dS}} = L_{\pm \frac{1}{2}, \mp \frac{1}{2}}^{\text{dS}} = L_{\pm \frac{1}{2}}^{\text{dS}}. \quad (6.36)$$

It is well known how to compute its resolvent: The following functions solve

$$(-\partial_x^2 + k^2)\phi(x) = 0 \quad (6.37)$$

$$\text{decaying at } +\infty: e^{-kx}, \quad \text{Re}(k) > 0, \quad (6.38)$$

$$\text{decaying at } -\infty: e^{kx}, \quad \text{Re}(k) > 0. \quad (6.39)$$

They have the Wronskian:

$$\mathcal{W}(e^{-kx}, e^{kx}) = 2k. \quad (6.40)$$

Now for $\text{Re } k > 0$,

$$(L + k^2)^{-1}(x, y) = \frac{1}{2k} \begin{cases} e^{kx}e^{-ky}, & \text{if } x < y, \\ e^{ky}e^{-kx}, & \text{if } y < x. \end{cases} \quad (6.41)$$

To see that (6.41) agrees with (4.51), we identify $k = \frac{\mu}{2}$, use

$$Q_{\pm \frac{1}{2}, \pm \frac{1}{2}, \mu}^{\text{dS}}(x) = Q_{\pm \frac{1}{2}, \mp \frac{1}{2}, \mu}^{\text{dS}}(x) = \frac{2^{2k}}{\Gamma(2k+1)} e^{-kx}, \quad (6.42)$$

which follows from (C.35), and

$$\Gamma(k)\Gamma\left(k + \frac{1}{2}\right)^2\Gamma(k+1) = \frac{2^{-4k}\pi}{k}\Gamma(2k+1)^2. \quad (6.43)$$

7 Geometric applications

In this section we show major applications of hypergeometric Hamiltonians in geometry. We will obtain these Hamiltonians as the results of separation of variables of (pseudo-)Laplacians on various (pseudo-)spheres.

Recall that every (pseudo-)Riemannian manifold is equipped with a certain natural differential operator called the *(pseudo-)Laplacian*. Suppose we fix coordinates $x = x_1, \dots, x_d$, the (pseudo-)metric is given by the field of symmetric invertible matrices $[g_{ij}]$, so that the the “line element” is

$$ds^2 = \sum_{1 \leq i, j \leq d} g_{ij} dx_i dx_j. \quad (7.1)$$

Then the pseudo-Laplacian is given by

$$\Delta = \frac{1}{\sqrt{|\det g|}} \partial_{x_i} g^{ij} \sqrt{|\det g|} \partial_{x_j}, \quad (7.2)$$

where $[g^{ij}]$ is the inverse of $[g_{ij}]$. In the case of the Riemannian signature, Δ is called the *Laplacian* (or the *Laplace-Beltrami operator*). For the Lorentzian signature the usual name is the *d'Alembertian*.

(Pseudo-)spheres in a *(pseudo-)Euclidean space* $\mathbb{R}^{p,q}$ inherit a (pseudo-)Riemannian structure from the ambient space. If the ambient space is Euclidean, they are called *spheres*. Otherwise, they are various kinds of *hyperboloids*.

Below we describe a few examples of separation of variables for a (pseudo-)Laplacian on a (pseudo-)sphere in appropriate coordinate systems. We will see that after an appropriate gauging, subtraction of a constant and restriction to an invariant subspace one obtains various hypergeometric Hamiltonians. These computations motivate the names “spherical”, “hyperbolic” and “deSitterian” that we use in our paper for various types of hypergeometric Hamiltonians.

The best known among these Laplacians is Δ_d^s , the Laplacian on the d -dimensional sphere \mathbb{S}^d . As it is well-known, it has the spectrum

$$\sigma(\Delta_d^s) = \{-l(l+d-1) \mid l \in \mathbb{N}_0\}. \quad (7.3)$$

(This is, incidentally, a consequence of computations in Subsection 7.1 and the properties of the spherical Gegenbauer Hamiltonian). Eigenfunctions of Δ_d^s with eigenvalue $-l(l+d-1)$ will be called *d-dimensional spherical harmonics of order l*.

7.1 Sphere

The unit d -dimensional sphere is defined as

$$\mathbb{S}^d := \{X \in \mathbb{R}^{d+1} \mid X_0^2 + X_1^2 + \dots + X_d^2 = 1\}. \quad (7.4)$$

We will denote elements of \mathbb{S}^{d-1} by \hat{X} and the corresponding element of length by $d\hat{X}^2$. On \mathbb{S}^d we will use the coordinates (r, \hat{X})

$$X_0 = \cos r, \quad X_i = \sin r \hat{X}_i, \quad i = 1, \dots, d, \quad \hat{X} \in \mathbb{S}^{d-1}, \quad (7.5)$$

or (w, \hat{X}) with $\cos r = w$, $\sin r = \sqrt{1 - w^2}$: In these coordinates we first write the line element, then the Laplacian:

$$ds^2 = dr^2 + \sin^2 r d\hat{X}^2 \quad (7.6)$$

$$= \frac{dw^2}{1 - w^2} + (1 - w^2)d\hat{X}^2; \quad (7.7)$$

$$\Delta_d^s = \partial_r^2 + (d - 1) \cot r \partial_r + \frac{\Delta_{d-1}^s}{\sin^2 r} \quad (7.8)$$

$$= (1 - w^2)\partial_w^2 - dw\partial_w + \frac{\Delta_{d-1}^s}{1 - w^2}. \quad (7.9)$$

Finally, we perform an appropriate gauging:

$$(\sin r)^{\frac{d-1}{2}} (-\Delta_d^s)(\sin r)^{-\frac{d-1}{2}} + \left(\frac{d-1}{2}\right)^2 = -\partial_r^2 + \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\sin^2 r}. \quad (7.10)$$

Thus the Laplacian on \mathbb{S}^d on $(d - 1)$ -dimensional spherical harmonics of order l (7.10) reduces to the spherical Gegenbauer Hamiltonian L_α^s with $\alpha = (\frac{d}{2} - 1 + l)$.

7.2 Hyperbolic space

The d -dimensional hyperbolic space is defined as

$$\mathbb{H}^d := \{X \in \mathbb{R}^{d+1} \mid -X_0^2 + X_1^2 + \dots + X_d^2 = -1, \quad X_0 > 0\}. \quad (7.11)$$

On \mathbb{H}^d we will use the following coordinates: (r, \hat{X}) , where $\hat{X} \in \mathbb{S}^{d-1}$:

$$X_0 = \cosh r, \quad i = 1, \dots, d, \quad X_i = \sinh r \hat{X}_i, \quad (7.12)$$

or (w, \hat{X}) with $\cosh r = w$, $\sinh r = \sqrt{w^2 - 1}$. In these coordinates we first write the line element, then the Laplacian:

$$ds^2 = dr^2 + \sinh^2 r d\hat{X}^2 \quad (7.13)$$

$$= \frac{dw^2}{w^2 - 1} + (w^2 - 1)d\hat{X}^2; \quad (7.14)$$

$$\Delta_d^h = \partial_r^2 + (d - 1) \coth r \partial_r + \frac{\Delta_{d-1}^s}{\sinh^2 r} \quad (7.15)$$

$$= (w^2 - 1)\partial_w^2 + dw\partial_w + \frac{\Delta_{d-1}^s}{w^2 - 1}. \quad (7.16)$$

Finally, we perform an appropriate gauging:

$$(\sinh r)^{\frac{d-1}{2}} (-\Delta_d^h)(\sinh r)^{-\frac{d-1}{2}} - \left(\frac{d-1}{2}\right)^2 = -\partial_r^2 + \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\sinh^2 r}. \quad (7.17)$$

Thus the Laplacian on \mathbb{H}^d on $d - 1$ -dimensional spherical harmonics of order l reduces to the hyperbolic Gegenbauer Hamiltonian L_α^h with $\alpha = \frac{d}{2} - 1 + l$.

7.3 DeSitter space

The deSitter space is defined as

$$dS^d := \{X \in \mathbb{R}^{d+1} \mid -X_0^2 + X_1^2 + \cdots + X_d^2 = 1\}. \quad (7.18)$$

On dS^d we will use the following coordinates: (t, \hat{X}) , where $\hat{X} \in \mathbb{S}^{d-1}$,

$$X_0 = \sinh t, \quad X_i = \cosh t \hat{X}_i, \quad i = 1, \dots, d, \quad (7.19)$$

or (w, \hat{X}) with $\sinh t = w$, $\cosh t = \sqrt{1 + w^2}$. In these coordinates we first write the line element, then the d'Alembertian:

$$\begin{aligned} ds^2 &= -dt^2 + \cosh^2 t d\hat{X}^2 \\ &= -\frac{dw^2}{w^2 + 1} + (w^2 + 1)d\hat{X}^2 \end{aligned} \quad (7.20)$$

$$\begin{aligned} \square_d^{dS} &= -\partial_t^2 + (d-1) \tanh t \partial_t + \frac{\Delta_{d-1}^s}{\cosh^2 t} \\ &= -(w^2 + 1)\partial_w^2 - dw\partial_w + \frac{\Delta_{d-1}^s}{w^2 + 1}. \end{aligned} \quad (7.21)$$

Finally, we perform an appropriate gauging:

$$(\cosh t)^{\frac{d-1}{2}} \square_d^{dS} (\cosh t)^{-\frac{d-1}{2}} - \left(\frac{d-1}{2}\right)^2 = -\partial_t^2 - \frac{\left(\frac{d-2}{2}\right)^2 - \frac{1}{4} - \Delta_{d-1}^s}{\cosh^2 t}. \quad (7.22)$$

Thus the d'Alembertian on dS^d on $d-1$ -dimensional spherical harmonics of order l reduces to the deSitterian Gegenbauer Hamiltonian L_α^{dS} with $\alpha = \frac{d}{2} - 1 + l$.

7.4 Sphere in double spherical coordinates

Consider the unit sphere of dimension $p+q-1$ with coordinates partitioned in two groups:

$$\mathbb{S}^{p+q-1} := \{(X, Y) \in \mathbb{R}^{p+q} \mid X_1^2 + \cdots + X_p^2 + Y_1^2 + \cdots + Y_q^2 = 1\}. \quad (7.23)$$

We consider also two spheres of dimension $p-1$ and $q-1$:

$$\mathbb{S}^{p-1} = \{X \in \mathbb{R}^p : X_1^2 + \cdots + X_p^2 = 1\}, \quad \mathbb{S}^{q-1} = \{Y \in \mathbb{R}^q : Y_1^2 + \cdots + Y_q^2 = 1\} \quad (7.24)$$

\mathbb{S}^{p+q-1} is parametrized by (τ, \hat{X}, \hat{Y}) , with $0 \leq \tau \leq \frac{\pi}{2}$, $\hat{X} \in \mathbb{S}^{p-1}$, $\hat{Y} \in \mathbb{S}^{q-1}$:

$$X_i = \sin \tau \hat{X}_i, \quad i = 1, \dots, p; \quad Y_j = \cos \tau \hat{Y}_j, \quad i = 1, \dots, q. \quad (7.25)$$

Alternatively, one can use coordinates (w, \hat{X}) where

$$\sin^2 \tau = w, \quad \cos^2 \tau = 1 - w. \quad (7.26)$$

We compute the line element and the Laplacian:

$$\begin{aligned} ds^2 &= \frac{dw^2}{4w(1-w)} + wd\hat{X}^2 + (1-w)d\hat{Y}^2 \\ &= d\tau^2 + \sin^2 \tau d\hat{X}^2 + \cos^2 \tau d\hat{Y}^2 \end{aligned} \quad (7.27)$$

$$\begin{aligned} \Delta_{p+q-1}^s &= 4w(1-w)\partial_w^2 + 2(p(1-w) - qw)\partial_w + \frac{\Delta_{p-1}^s}{w} + \frac{\Delta_{q-1}^s}{1-w} \\ &= \partial_\tau^2 + ((p-1)\cot \tau - (q-1)\tan \tau)\partial_\tau + \frac{\Delta_{p-1}^s}{\sin^2 \tau} + \frac{\Delta_{q-1}^s}{\cos^2 \tau}. \end{aligned} \quad (7.28)$$

We perform an appropriate gauging:

$$\begin{aligned} & (\sin \tau)^{\frac{p-1}{2}} (\cos \tau)^{\frac{q-1}{2}} (-\Delta_{p+q-1}^s) (\sin \tau)^{-\frac{p-1}{2}} (\cos \tau)^{-\frac{q-1}{2}} + \left(\frac{p+q-2}{2}\right)^2 \\ &= -\partial_\tau^2 + \frac{\left(\frac{p-2}{2}\right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{\sin^2 \tau} + \frac{\left(\frac{q-2}{2}\right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{\cos^2 \tau}. \end{aligned} \quad (7.29)$$

Finally, we make a substitution $\tau = \frac{r}{2}$:

$$(7.29) = 4 \left(-\partial_r^2 + \frac{\left(\frac{p-2}{2}\right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{4 \sin^2 \frac{r}{2}} + \frac{\left(\frac{q-2}{2}\right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{4 \cos^2 \frac{r}{2}} \right). \quad (7.30)$$

Thus on products of a spherical harmonic of order j and l we obtain the spherical hypergeometric Hamiltonian of the first kind $L_{\alpha,\beta}^s$ with

$$\alpha = \frac{p}{2} - 1 + j, \quad \beta = \frac{q}{2} - 1 + l. \quad (7.31)$$

7.5 Hyperboloid in double spherical coordinates

Consider the hyperboloid of signature $p-1, q$ embedded in the pseudoEuclidean space of signature (p, q) :

$$\mathbb{H}^{p-1,q} := \{(X, Y) \in \mathbb{R}^{p+q} \mid -X_1^2 - \dots - X_p^2 + Y_1^2 + \dots + Y_q^2 = -1\}. \quad (7.32)$$

Let \mathbb{S}^{p-1} and \mathbb{S}^{q-1} be as in (7.24). $\mathbb{H}^{p-1,q}$ is parametrized by (τ, \hat{X}, \hat{Y}) , with $0 \leq \tau \leq \infty$, $\hat{X} \in \mathbb{S}^{p-1}$, $\hat{Y} \in \mathbb{S}^{q-1}$:

$$X_i = \cosh \tau \hat{X}_i, \quad i = 1, \dots, p; \quad Y_j = \sinh \tau \hat{Y}_j, \quad i = 1, \dots, q. \quad (7.33)$$

Alternatively, one can use coordinates (w, \hat{X}, \hat{Y}) where

$$\cosh^2 r = w, \quad \sinh^2 r = w - 1. \quad (7.34)$$

The line element and the pseudo-Laplacian in these coordinates:

$$\begin{aligned} ds^2 &= \frac{dw^2}{4w(w-1)} - w d\hat{X}^2 + (w-1) d\hat{Y}^2 \\ &= d\tau^2 - \cosh^2 \tau d\hat{X}^2 + \sinh^2 \tau d\hat{Y}^2 \end{aligned} \quad (7.35)$$

$$\begin{aligned} \Delta_{p-1,q} &= 4w(w-1) \partial_w^2 + 2(p(w-1) + qw) \partial_w - \frac{\Delta_{p-1}^s}{w} + \frac{\Delta_{q-1}^s}{w-1} \\ &= \partial_\tau^2 + ((p-1) \tanh \tau + (q-1) \coth \tau) \partial_\tau - \frac{\Delta_{p-1}^s}{\cosh^2 \tau} + \frac{\Delta_{q-1}^s}{\sinh^2 \tau}. \end{aligned} \quad (7.36)$$

We perform an appropriate gauging:

$$\begin{aligned} & (\cosh \tau)^{\frac{p-1}{2}} (\sinh \tau)^{\frac{q-1}{2}} (-\Delta_{p-1,q}) (\cosh \tau)^{-\frac{p-1}{2}} (\sinh \tau)^{-\frac{q-1}{2}} - \left(\frac{p+q-2}{2}\right)^2 \\ &= -\partial_\tau^2 - \frac{\left(\frac{p-2}{2}\right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{\cosh^2 \tau} + \frac{\left(\frac{q-2}{2}\right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{\sinh^2 \tau}. \end{aligned} \quad (7.37)$$

Finally, we substitute $\tau = \frac{r}{2}$:

$$(7.37) = 4 \left(-\partial_r^2 - \frac{\left(\frac{p-2}{2}\right)^2 - \frac{1}{4} - \Delta_{p-1}^s}{4 \cosh^2 \frac{r}{2}} + \frac{\left(\frac{q-2}{2}\right)^2 - \frac{1}{4} - \Delta_{q-1}^s}{4 \sinh^2 \frac{r}{2}} \right). \quad (7.38)$$

Thus on the product of a spherical harmonic of order j and l we obtain the hyperbolic hypergeometric Hamiltonian of the first kind $L_{\alpha,\beta}^h$ with

$$\alpha = \frac{q}{2} - 1 + l, \quad \beta = \frac{p}{2} - 1 + j. \quad (7.39)$$

7.6 Complex manifolds

All the manifolds that we used so far were real. In the next subsection we will need a complex (analytic) manifold. They have essentially the same formalism as real manifolds. Let us briefly sketch its elements. For more details, see [LeB].

Suppose that a complex manifold is equipped with local complex coordinates $z = (z_1, \dots, z_d)$ and the holomorphic line element

$$\sum_{1 \leq i, j \leq d} g_{ij} dz_i dz_j, \quad (7.40)$$

where g_{ij} is a complex symmetric invertible matrix. The corresponding complex Laplacian is defined by essentially the same formula as in the real case:

$$\Delta = \frac{1}{\sqrt{\det g}} \partial_{z_i} g^{ij} \sqrt{\det g} \partial_{z_j}. \quad (7.41)$$

Note that there is no absolute value, and $\det g$ is always non-zero. The definition of Δ does not depend on the choice of the branch of the (double-valued) square root.

Suppose that the manifold is equipped with a conjugation, in the coordinates given by $z_i \mapsto \bar{z}_i$. We then also have the anti-holomorphic line element

$$\sum_{1 \leq i, j \leq d} \bar{g}_{ij} d\bar{z}_i d\bar{z}_j. \quad (7.42)$$

and the corresponding conjugate Laplacian

$$\bar{\Delta} = \frac{1}{\sqrt{\det \bar{g}}} \partial_{\bar{z}_i} \bar{g}^{i\bar{j}} \sqrt{\det \bar{g}} \partial_{\bar{z}_j}. \quad (7.43)$$

As our first example consider the space \mathbb{C}^{d+1} equipped with the line element

$$dZ^2 = dZ_0^2 + dZ_1^2 + \dots + dZ_p^2. \quad (7.44)$$

The corresponding Laplacian is obviously

$$\Delta = \partial_{Z_0}^2 + \dots + \partial_{Z_p}^2. \quad (7.45)$$

Note that our standard identification of \mathbb{C}^{d+1} with $\mathbb{R}^{2(d+1)}$ will be

$$Z_i = \frac{1}{\sqrt{2}}(X_i + \mathbf{i}Y_i) \quad (7.46)$$

(and not $Z_i = X_i + \mathbf{i}Y_i$. Therefore,

$$\partial_{Z_i} = \frac{1}{\sqrt{2}}(\partial_{X_i} - \mathbf{i}\partial_{Y_i}), \quad (7.47)$$

(and not, as usual, $\partial_{Z_i} = \frac{1}{2}(\partial_{X_i} - \mathbf{i}\partial_{Y_i})$.) Clearly, with this definition $\langle dZ_i | \partial_{Z_j} \rangle = \delta_{ij}$.

Another example of a complex manifold is the unit sphere [HIU]

$$\mathbb{S}_{\mathbb{C}}^d := \{Z \in \mathbb{C}^{d+1} | Z_0^2 + Z_1^2 + \dots + Z_d^2 = 1\}. \quad (7.48)$$

Introducing the complex polar coordinanes

$$R := \sqrt{Z_0^2 + \cdots + Z_d^2}, \quad \hat{Z}_i := \frac{Z_i}{R}, \quad (7.49)$$

where we arbitrarily select the branch of the square root, we have the direct analog of the formula from the real case:

$$dZ^2 = dR^2 + R^2 d\hat{Z}^2, \quad (7.50)$$

where $d\hat{Z}^2$ is the complex line element on $\mathbb{S}_{\mathbb{C}}^d$. The corresponding complex Laplacian is given by the same expressions as in the real case:

$$\begin{aligned} \Delta_{d,\mathbb{C}}^s &= \sum_{0 \leq i < j \leq d} (Z_i \partial_{Z_j} - Z_j \partial_{Z_i})^2 \\ &= R^2 \Delta_{d+1,\mathbb{C}} - R^2 \partial_R^2 - d R \partial_R. \end{aligned} \quad (7.51)$$

7.7 Hyperboloid $\mathbb{H}^{p-1,p}$ in complex coordinates

Consider the hyperboloid $\mathbb{H}^{p-1,p}$ embedded in \mathbb{R}^{2p} , defined as in Subsection 7.5, with the coordinates $X_i, Y_i \in \mathbb{R}$, $i = 1, \dots, p$. We identify \mathbb{R}^{2p} with \mathbb{C}^p as in (7.46), so that we obtain two representations of $\mathbb{H}^{p-1,p}$, a real and a complex one:

$$\begin{aligned} \mathbb{H}^{p-1,p} &= \{(X, Y) \in \mathbb{R}^{2p} \mid -Y_1^2 - \cdots - Y_p^2 + X_1^2 + \cdots + X_p^2 = -1\} \\ &= \{Z \in \mathbb{C}^p \mid Z_1^2 + \cdots + Z_p^2 + \bar{Z}_1^2 + \cdots + \bar{Z}_p^2 = 1\} \end{aligned} \quad (7.52)$$

The (real) line element on $\mathbb{R}^{2p} = \mathbb{C}^p$ can be written as

$$\begin{aligned} & -dY_1^2 - \cdots - dY_p^2 + dX_1^2 + \cdots + dX_p^2 \\ &= dZ_1^2 + \cdots + dZ_p^2 + d\bar{Z}_1^2 + \cdots + d\bar{Z}_p^2 \\ &= dR^2 + R^2 d\hat{Z}^2 + d\bar{R}^2 + \bar{R}^2 d\bar{\hat{Z}}^2 \end{aligned} \quad (7.53)$$

Now on $\mathbb{H}^{p-1,p}$ we have $R^2 + \bar{R}^2 = 1$. Therefore $R^2 = \frac{1+i \sinh r}{2}$ for a unique $r \in \mathbb{R}$. Thus we can parametrize $\mathbb{H}^{p-1,p}$ with $r \in \mathbb{R}$, $\hat{Z} \in \mathbb{S}_{\mathbb{C}}^{p-1}$

$$Z_i = \sqrt{\frac{1+i \sinh r}{2}} \hat{Z}_i, \quad (7.54)$$

where we take the principal branch of square root. The (real) line element and the pseudo-Laplacian are

$$ds^2 = -\frac{1}{4} dr^2 + \frac{1+i \sinh r}{2} d\hat{Z}^2 + \frac{1-i \sinh r}{2} d\bar{\hat{Z}}^2, \quad (7.55)$$

$$\Delta_{p-1,p} = -4\partial_r^2 - 4(p-1) \tanh r \partial_r + \frac{2}{1+i \sinh r} \Delta_{p-1}^{s,\mathbb{C}} + \frac{2}{1-i \sinh r} \overline{\Delta_{p-1}^{s,\mathbb{C}}}. \quad (7.56)$$

We perform an appropriate gauging:

$$\begin{aligned} & (\cosh r)^{\frac{p-1}{2}} (\Delta_{p-1,p}) (\cosh r)^{-\frac{p-1}{2}} - (p-1)^2 \\ &= 4 \left(-\partial_r^2 - \frac{(-\Delta_{p-1}^{s,\mathbb{C}} + (\frac{p-1}{2})^2 - \frac{1}{4})}{2(1+i \sinh r)} - \frac{(-\overline{\Delta_{p-1}^{s,\mathbb{C}}} + (\frac{p-1}{2})^2 - \frac{1}{4})}{2(1-i \sinh r)} \right). \end{aligned} \quad (7.57)$$

Thus on joint eigenvectors of $\Delta_{p-1}^{\text{s},\mathbb{C}}$ and $\overline{\Delta_{p-1}^{\text{s},\mathbb{C}}}$ of degree l , resp. j we obtain the deSitterian hypergeometric Hamiltonian of the first kind $L_{\alpha,\beta}^{\text{dS}}$ with

$$\alpha = \frac{p}{2} - 1 + l, \quad \beta = \frac{p}{2} - 1 + j. \quad (7.58)$$

Acknowledgement. J.D. was supported by National Science Center (Poland) under the grant UMO-2019/35/B/ST1/01651. P.K supported by the OPUS grant no. 2022/47/B/ST2/03313 “Quantum geometry and BPS states” funded by the National Science Centre (Poland).

A Holomorphic families of operators

The concept of a holomorphic family of bounded operators is well-known. Less known is the concept of a holomorphic family of closed operators, which we use throughout our paper. Both are described in the monograph of Kato [Kato], Chap. 7, and also in [DW1]. For the convenience of the reader, we give a concise account of these concepts in this appendix.

In the following, Θ is an open subset of \mathbb{C} , $\mathcal{H}_1, \mathcal{H}_2$ are Hilbert spaces, $B(\mathcal{H}_1, \mathcal{H}_2)$ denotes the space of bounded operators from \mathcal{H}_1 to \mathcal{H}_2 and $C(\mathcal{H}_1, \mathcal{H}_2)$ the set of closed operators from \mathcal{H}_1 to \mathcal{H}_2 . $\mathcal{D}(H)$ will denote the domain of an operator H and $\rho(H) := \mathbb{C} \setminus \sigma(H)$ its resolvent set.

There exist several equivalent definitions of a holomorphic family of bounded operators. Here is one of them: Consider a function $\Theta \ni z \mapsto H(z) \in B(\mathcal{H}_1, \mathcal{H}_2)$. We say that this is a *holomorphic family of bounded operators* if for any $f \in \mathcal{H}_2$, $g \in \mathcal{H}_1$ the function $\Theta \ni z \mapsto (f|H(z)g) \in \mathbb{C}$ is holomorphic.

We have the following practical criterion:

Theorem A.1. *Suppose that $\Theta \ni z \mapsto H(z) \in B(\mathcal{H}_1, \mathcal{H}_2)$ is a function uniformly bounded on compact subsets of Θ . Suppose in addition that there exist dense subspaces $\mathcal{X}_1 \subset \mathcal{H}_1$, $\mathcal{X}_2 \subset \mathcal{H}_2$ such that for any $f \in \mathcal{X}_1$, $g \in \mathcal{X}_2$, the function $\Theta \ni z \mapsto (f|H(z)g) \in \mathbb{C}$ is holomorphic. Then $\Theta \ni z \mapsto H(z)$ is a holomorphic family of bounded operators.*

There exists also a natural concept of a holomorphic family of closed operators, due to Kato, it is however more subtle and less known. Let us describe its definition.

Consider a function $\Theta \ni z \mapsto H(z) \in C(\mathcal{H}_1, \mathcal{H}_2)$. We say that it is a *holomorphic family of closed operators* if for each $z_0 \in \Theta$ there exists a neighborhood Θ_0 of z_0 , a Hilbert space \mathcal{K} and a holomorphic family of bounded operators $\Theta_0 \ni z \mapsto A(z) \in B(\mathcal{K}, \mathcal{H}_1)$ that map bijectively \mathcal{K} onto $\mathcal{D}(H(z))$ and

$$\Theta_0 \ni z \mapsto H(z)A(z) \in B(\mathcal{K}, \mathcal{H})$$

is a holomorphic family of bounded operators.

We have the following practical criterion, which works for operators having non-empty resolvent sets:

Theorem A.2. *Consider a function $\Theta \ni z \mapsto H(z) \in C(\mathcal{H})$. Suppose in addition that for any $z \in \Theta$ the resolvent set of $H(z)$ is nonempty. If for any $z_0 \in \Theta$ there exists $\lambda \in \mathbb{C}$ and a neighborhood Θ_0 of z_0 such that $\lambda \in \rho(H(z))$ for $z \in \Theta_0$ and $z \mapsto (H(z) - \lambda)^{-1} \in B(\mathcal{H})$ is holomorphic on Θ_0 , then $z \mapsto H(z)$ is a holomorphic family of closed operators.*

Proof. We set $\mathcal{K} := \mathcal{H}$ and $A(z) := (H(z) - \lambda)^{-1}$, so that $H(z)A(z) = 1 + \lambda A(z)$. \square

B Riemann equation

B.1 Regular-singular points

Consider an ordinary 2nd order differential equation with meromorphic coefficients:

$$(b(z)\partial_z^2 + c(z)\partial_z + d(z))u(z) = 0. \quad (\text{B.59})$$

If $\frac{c(z)}{b(z)}$ or $\frac{d(z)}{b(z)}$ are singular at $z_0 \in \mathbb{C}$, then we say that z_0 is a singular point of (B.59). Following [WW], we distinguish a special class of singular points:

Definition B.1. *We say that (B.59) has a regular-singular point at z_0 , if $\frac{c(z)}{b(z)}$ has a pole of at most 1st order at z_0 , and $\frac{d(z)}{b(z)}$ has a pole of at most 2nd order at z_0 .*

We can rewrite (B.59) as

$$(p(z)(z - z_0)^2\partial_z^2 + q(z)(z - z_0)\partial_z + r(z))u(z) = 0. \quad (\text{B.60})$$

If $p(z_0) \neq 0$, then z_0 is regular-singular iff p, q, r are analytic at 0. Define the *indicial polynomial* of z_0

$$P_{z_0}(\lambda) := \lambda(\lambda - 1)p(z_0) + \lambda q(z_0) + r(z_0). \quad (\text{B.61})$$

The roots of P_{z_0} are called *indices* of the regular-singular point z_0 .

The importance of regular-singular points and their indices stems from the Frobenius Method, described in the following theorem [WW, Ince]:

Theorem B.2. *Assume that $p(z_0) \neq 0$ and p, q, r are holomorphic in an open connected simply connected set $\Omega \subset \mathbb{C}$ containing z_0 . Let $\rho \in \mathbb{C}$ satisfy*

$$P_{z_0}(\rho) = 0, \quad P_{z_0}(\rho + n) \neq 0, \quad n = 1, 2, \dots \quad (\text{B.62})$$

Then there exists a unique function $\tilde{u}(z)$ holomorphic in Ω , such that $u(z) := (z - z_0)^\rho \tilde{u}(z)$ solves (B.60) and $\tilde{u}(z_0) = 1$.

B.2 Riemann equation

Equation with 3 singular points on the Riemann sphere $\mathbb{C} \cup \{\infty\}$, all of them regular-singular, is classified by the positions of its singularities and the values of indices. It is called the *Riemann*, or sometimes the *Papperitz equation*, and is described in the following well-known theorem [WW, Ince]. For the Riemann operator we use the notation introduced in [DW], which is inspired by the notation due to Papperitz used in [WW].

Theorem B.3. (1) *Suppose that we are given a 2nd order differential equation on the Riemann sphere having at most 3 singular points z_1, z_2, z_3 , all of them regular-singular, with the following indices*

$$z_1 : \rho_1, \tilde{\rho}_1; \quad z_2 : \rho_2, \tilde{\rho}_2; \quad z_3 : \rho_3, \tilde{\rho}_3.$$

Then the following condition is satisfied:

$$\rho_1 + \tilde{\rho}_1 + \rho_2 + \tilde{\rho}_2 + \rho_3 + \tilde{\rho}_3 = 1. \quad (\text{B.63})$$

Such an equation, normalized to have coefficient 1 at the 2nd derivative, is for finite z_1, z_2, z_3 given by the operator

$$\mathcal{P} \begin{pmatrix} z_1 & z_2 & z_3 \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix} := \partial_z^2 - \left(\frac{\rho_1 + \tilde{\rho}_1 - 1}{z - z_1} + \frac{\rho_2 + \tilde{\rho}_2 - 1}{z - z_2} + \frac{\rho_3 + \tilde{\rho}_3 - 1}{z - z_3} \right) \partial_z$$

$$+ \frac{\rho_1 \tilde{\rho}_1 (z_1 - z_2)(z_1 - z_3)}{(z - z_1)^2 (z - z_2)(z - z_3)} + \frac{\rho_2 \tilde{\rho}_2 (z_2 - z_3)(z_2 - z_1)}{(z - z_2)^2 (z - z_3)(z - z_1)} + \frac{\rho_3 \tilde{\rho}_3 (z_3 - z_1)(z_3 - z_2)}{(z - z_3)^2 (z - z_1)(z - z_2)},$$

and for $z_3 = \infty$ by the operator

$$\mathcal{P} \begin{pmatrix} z_1 & z_2 & \infty \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix} := \partial_z^2 - \left(\frac{\rho_1 + \tilde{\rho}_1 - 1}{z - z_1} + \frac{\rho_2 + \tilde{\rho}_2 - 1}{z - z_2} \right) \partial_z$$

$$+ \frac{\rho_1 \tilde{\rho}_1 (z_1 - z_2)}{(z - z_1)^2 (z - z_2)} + \frac{\rho_2 \tilde{\rho}_2 (z_2 - z_1)}{(z - z_2)^2 (z - z_1)} + \frac{\rho_3 \tilde{\rho}_3}{(z - z_1)(z - z_2)}. \quad (\text{B.64})$$

- (2) With the help of homographies (called also Möbius transformations) we can move around singularities. More precisely, if $z \mapsto w(z) = \frac{az+b}{cz+d}$, where we can assume that $ad - bc = 1$, then

$$\mathcal{P} \begin{pmatrix} w(z_1) & w(z_2) & w(z_3) \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix} = (cz + d)^4 \mathcal{P} \begin{pmatrix} z_1 & z_2 & z_3 \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix}.$$

- (3) By gauging with powers we can shift the indices:

$$(z - z_1)^{-\lambda} (z - z_2)^\lambda \mathcal{P} \begin{pmatrix} z_1 & z_2 & z_3 \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix} (z - z_1)^\lambda (z - z_2)^{-\lambda}$$

$$= \mathcal{P} \begin{pmatrix} z_1 & z_2 & z_3 \\ \rho_1 - \lambda & \rho_2 + \lambda & \rho_3 \\ \tilde{\rho}_1 - \lambda & \tilde{\rho}_2 + \lambda & \tilde{\rho}_3 \end{pmatrix}, \quad (\text{B.65})$$

$$(z - z_1)^{-\lambda} \mathcal{P} \begin{pmatrix} z_1 & z_2 & \infty \\ \rho_1 & \rho_2 & \rho_3 \\ \tilde{\rho}_1 & \tilde{\rho}_2 & \tilde{\rho}_3 \end{pmatrix} (z - z_1)^\lambda$$

$$= \mathcal{P} \begin{pmatrix} z_1 & z_2 & z_3 \\ \rho_1 - \lambda & \rho_2 & \rho_3 + \lambda \\ \tilde{\rho}_1 - \lambda & \tilde{\rho}_2 & \tilde{\rho}_3 + \lambda \end{pmatrix}. \quad (\text{B.66})$$

B.3 From Riemann equation to hypergeometric equation

The Riemann equation can be simplified. First, by Thm B.3 (2), we can assume that the points z_1, z_2, z_3 are any triplet of distinct points on the Riemann sphere. We choose them to be $0, 1, \infty$. Then, by Thm B.3 (3), we can assume that ρ_1, ρ_2 are arbitrary numbers. We choose them to be both 0. The sum of remaining indices must be 1. Hence we can assume that 0 has indices $0, 1 - c$; 1 has indices $0, c - a - b$; and ∞ has indices a, b . The hypergeometric operator is

$$\mathcal{F}(a, b; c; z, \partial_z) := z(1 - z) \mathcal{P} \begin{pmatrix} 0 & 1 & \infty \\ 0 & 0 & a \\ 1 - c & c - a - b & b \end{pmatrix} (z, \partial_z) \quad (\text{B.67})$$

$$= z(1 - z) \partial_z^2 + (c - (a + b + 1)z) \partial_z - ab. \quad (\text{B.68})$$

The hypergeometric function $F(a, b; c; z)$ is the unique solution of

$$\mathcal{F}(a, b; c; z, \partial_z)F(z) = 0, \quad F(0) = 1. \quad (\text{B.69})$$

Thus it is the solution of the hypergeometric equation given by the Frobenius method (Thm B.2) for the singularity $z = 0$ and index 0.

C Identities for the hypergeometric function

In this section, we review identities for the hypergeometric and Gegenbauer functions with the emphasis on the connection formulas and Kummer's table. There are many reviews available including [WW, NIST1]. Our presentation is close to [D1, D2].

C.1 Kummer's table

Recall that the hypergeometric equation is given by the operator $\mathcal{F}(a, b; c; z, \partial_z)$ defined in (2.1), and the hypergeometric function with Olver's normalization $\mathbf{F}(a, b; c; z)$ is defined in (2.2). In this section, for brevity and transparency, we change the notation following [D1, D2], writing

$$\mathbf{F}_{\alpha, \beta, \mu}(z) := \mathbf{F}\left(\frac{1 + \alpha + \beta + \mu}{2}, \frac{1 + \alpha + \beta - \mu}{2}; 1 + \alpha; z\right). \quad (\text{C.1})$$

It is obvious from the definition that we have the following identity

$$\mathbf{F}_{\alpha, \beta, \mu}(z) = \mathbf{F}_{\alpha, \beta, -\mu}(z). \quad (\text{C.2})$$

The following 6 functions form a set of standard solutions of the hypergeometric equation. Each of the solutions can be expressed in 4 ways (actually, 4×2 ways if we include the trivial identity (C.2)). This yields $6 \times 4 = 24$ expressions usually called *Kummer's table*:

$$\begin{aligned} \mathbf{F}_{\alpha, \beta, \mu}(z) &= (1 - z)^{\frac{-1 - \alpha - \beta + \mu}{2}} \mathbf{F}_{\alpha, -\mu, -\beta}\left(\frac{z}{z - 1}\right) \\ &= (1 - z)^{\frac{-1 - \alpha - \beta - \mu}{2}} \mathbf{F}_{\alpha, \mu, \beta}\left(\frac{z}{z - 1}\right) \\ &= (1 - z)^{-\beta} \mathbf{F}_{\alpha, -\beta, -\mu}(z). \end{aligned} \quad (\text{C.3})$$

$$\begin{aligned} (-z)^\alpha \mathbf{F}_{-\alpha, \beta, -\mu}(z) &= (-z)^\alpha (1 - z)^{\frac{-1 + \alpha - \beta - \mu}{2}} \mathbf{F}_{-\alpha, \mu, -\beta}\left(\frac{z}{z - 1}\right) \\ &= (-z)^\alpha (1 - z)^{\frac{-1 + \alpha - \beta + \mu}{2}} \mathbf{F}_{-\alpha, -\mu, \beta}\left(\frac{z}{z - 1}\right) \\ &= (-z)^\alpha (1 - z)^{-\beta} \mathbf{F}_{-\alpha, -\beta, \mu}(z). \end{aligned} \quad (\text{C.4})$$

$$\begin{aligned} (-z)^{\frac{-1 - \alpha - \beta + \mu}{2}} \mathbf{F}_{-\mu, \beta, -\alpha}(z^{-1}) &= (1 - z)^{\frac{-1 - \alpha - \beta + \mu}{2}} \mathbf{F}_{-\mu, \alpha, -\beta}\left(\frac{1}{1 - z}\right) \\ &= (-z)^{-\alpha} (1 - z)^{\frac{-1 + \alpha - \beta - \mu}{2}} \mathbf{F}_{-\mu, -\alpha, \beta}\left(\frac{1}{1 - z}\right) \\ &= (-z)^{\frac{-1 - \alpha - \beta + \mu}{2}} (1 - z)^{-\beta} \mathbf{F}_{\alpha, -\beta, -\mu}(z^{-1}). \end{aligned} \quad (\text{C.5})$$

$$\begin{aligned}
(-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,\beta,\alpha}(z^{-1}) &= (1-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,-\alpha,-\beta} \left(\frac{1}{1-z} \right) \\
&= (-z)^{-\alpha} (1-z)^{\frac{-1+\alpha-\beta+\mu}{2}} \mathbf{F}_{\mu,\alpha,\beta} \left(\frac{1}{1-z} \right) \\
&= (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} (1-z)^{-\beta} \mathbf{F}_{-\alpha,-\beta,\mu}(z^{-1}).
\end{aligned} \tag{C.6}$$

$$\begin{aligned}
\mathbf{F}_{\beta,\alpha,\mu}(1-z) &= (-z)^{\frac{-1-\alpha-\beta+\mu}{2}} \mathbf{F}_{\beta,-\mu,-\alpha} \left(1 - \frac{1}{z} \right) \\
&= (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\beta,\mu,\alpha} \left(1 - \frac{1}{z} \right) \\
&= (-z)^{-\alpha} \mathbf{F}_{\beta,-\alpha,-\mu}(1-z).
\end{aligned} \tag{C.7}$$

$$\begin{aligned}
(1-z)^{-\beta} \mathbf{F}_{-\beta,-\mu,-\alpha}(1-z) &= (-z)^{\frac{-1-\alpha-\beta+\mu}{2}} \mathbf{F}_{\beta,-\mu,-\alpha} \left(1 - \frac{1}{z} \right) \\
&= (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\beta,\mu,\alpha} \left(1 - \frac{1}{z} \right) \\
&= (-z)^{-\alpha} \mathbf{F}_{\beta,-\alpha,-\mu}(1-z).
\end{aligned} \tag{C.8}$$

C.2 Connection formulas

Here are connection formulas. For $z \notin]-\infty, 0] \cup [1, \infty[$:

$$\begin{aligned}
&\mathbf{F}_{\beta,\alpha,\mu}(1-z) \\
&= \frac{\pi \mathbf{F}_{\alpha,\beta,\mu}(z)}{\sin(-\pi\alpha)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)} + \frac{\pi z^{-\alpha} \mathbf{F}_{-\alpha,\beta,-\mu}(z)}{\sin(\pi\alpha)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)},
\end{aligned} \tag{C.9}$$

$$\begin{aligned}
&(1-z)^{-\beta} \mathbf{F}_{-\beta,\alpha,-\mu}(1-z) \\
&= \frac{\pi \mathbf{F}_{\alpha,\beta,\mu}(z)}{\sin(-\pi\alpha)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)} + \frac{\pi z^{-\alpha} \mathbf{F}_{-\alpha,\beta,-\mu}(z)}{\sin(\pi\alpha)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)},
\end{aligned} \tag{C.10}$$

$$\begin{aligned}
&\mathbf{F}_{\alpha,\beta,\mu}(z) \\
&= \frac{\pi \mathbf{F}_{\beta,\alpha,\mu}(1-z)}{\sin(-\pi\beta)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)} + \frac{\pi (1-z)^{-\beta} \mathbf{F}_{-\beta,\alpha,-\mu}(1-z)}{\sin(\pi\beta)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)},
\end{aligned} \tag{C.11}$$

$$\begin{aligned}
&z^{-\alpha} \mathbf{F}_{-\alpha,\beta,\mu}(z) \\
&= \frac{\pi \mathbf{F}_{\beta,\alpha,\mu}(z)}{\sin(-\pi\beta)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)} + \frac{\pi (1-z)^{-\beta} \mathbf{F}_{-\beta,\alpha,-\mu}(1-z)}{\sin(\pi\beta)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)}.
\end{aligned} \tag{C.12}$$

For $z \notin [0, \infty[$:

$$\begin{aligned}
& (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,\beta,\alpha}(z^{-1}) \\
&= \frac{\pi \mathbf{F}_{\alpha,\beta,\mu}(z)}{\sin(-\pi\alpha)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)} + \frac{\pi (-z)^{-\alpha} \mathbf{F}_{-\alpha,\beta,\mu}(z)}{\sin(\pi\alpha)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)}; \tag{C.13}
\end{aligned}$$

$$\begin{aligned}
& (-z)^{\frac{-1-\alpha-\beta+\mu}{2}} \mathbf{F}_{-\mu,\beta,\alpha}(z^{-1}) \\
&= \frac{\pi \mathbf{F}_{\alpha,\beta,\mu}(z)}{\sin(-\pi\alpha)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)} + \frac{\pi (-z)^{-\alpha} \mathbf{F}_{-\alpha,\beta,\mu}(z)}{\sin(\pi\alpha)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)}. \tag{C.14}
\end{aligned}$$

$$\begin{aligned}
& \mathbf{F}_{\alpha,\beta,\mu}(z) \\
&= \frac{\pi (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,\beta,\alpha}(z^{-1})}{\sin(-\pi\mu)\Gamma\left(\frac{1+\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta-\mu}{2}\right)} + \frac{\pi (-z)^{\frac{-1-\alpha-\beta+\mu}{2}} \mathbf{F}_{-\mu,\beta,\alpha}(z^{-1})}{\sin(\pi\mu)\Gamma\left(\frac{1+\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1+\alpha-\beta+\mu}{2}\right)}, \tag{C.15}
\end{aligned}$$

$$\begin{aligned}
& (-z)^{-\alpha} \mathbf{F}_{-\alpha,\beta,\mu}(z) \\
&= \frac{\pi (-z)^{\frac{-1-\alpha-\beta-\mu}{2}} \mathbf{F}_{\mu,\beta,\alpha}(z^{-1})}{\sin(-\pi\mu)\Gamma\left(\frac{1-\alpha+\beta-\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta-\mu}{2}\right)} + \frac{\pi (-z)^{\frac{-1-\alpha-\beta+\mu}{2}} \mathbf{F}_{-\mu,\beta,\alpha}(z^{-1})}{\sin(\pi\mu)\Gamma\left(\frac{1-\alpha+\beta+\mu}{2}\right)\Gamma\left(\frac{1-\alpha-\beta+\mu}{2}\right)}. \tag{C.16}
\end{aligned}$$

C.3 Degenerate case

Let $\mu \in \mathbb{Z}$. Then we have the identity

$$\begin{aligned}
& \left(\frac{\alpha + \beta - \mu + 1}{2}\right)_{\mu} \left(\frac{\alpha - \beta - \mu + 1}{2}\right)_{\mu} F\left(\frac{\alpha + \beta + \mu + 1}{2}, \frac{\alpha - \beta + \mu + 1}{2}; 1 + \mu; z\right) \\
&= z^{-\mu} F\left(\frac{\alpha + \beta - \mu + 1}{2}, \frac{\alpha - \beta - \mu + 1}{2}; 1 - \mu; z\right). \tag{C.17}
\end{aligned}$$

C.4 Gegenbauer functions

In the remaining part of this section we review some of the relations for Gegenbauer function, following mostly [DGR].

Recall that the Gegenbauer equation is defined by the operator $\mathcal{G}_{\alpha,\lambda}(w, \partial_w)$ defined in (2.5), and the two Gegenbauer functions that we use were defined in (3.1) and (2.8). Let us rewrite their definitions using the notation introduced in (C.1).

The following function satisfies the Gegenbauer equation and has value 1 at 1:

$$S_{\alpha,\pm\lambda}(w) = F_{\alpha,\alpha,\lambda}\left(\frac{1-w}{2}\right). \tag{C.18}$$

It can be easily seen from (C.3) that the solution behaving as $\left(\frac{w+1}{2}\right)^{-\alpha}$ at 1 is

$$S_{-\alpha,\pm\lambda}(w) = \left(\frac{1+w}{2}\right)^{-\alpha} F_{-\alpha,\alpha,2\lambda}\left(\frac{1-w}{2}\right) \tag{C.19}$$

Solution behaving as $w^{-\frac{1}{2}-\lambda-\alpha}$ at $+\infty$ is

$$Z_{\alpha,\lambda}(w) = (w \pm 1)^{-\frac{1}{2}-\alpha-\lambda} F_{2\lambda,\alpha,\alpha}\left(\frac{2}{1 \pm w}\right) \tag{C.20}$$

It is useful to introduce Olver's normalization

$$\mathbf{S}_{\alpha,\lambda}(w) = \frac{S_{\alpha,\lambda}(w)}{\Gamma(1+\alpha)}, \quad \mathbf{Z}_{\alpha,\lambda}(w) = \frac{Z_{\alpha,\lambda}(w)}{\Gamma(1+\lambda)}. \quad (\text{C.21})$$

We can read their connection formula from the connection formulas of hypergeometric functions (C.13) (C.9), for $\text{Im}(w) < 0$:

$$\mathbf{Z}_{\alpha,\lambda}(w) = \frac{\sqrt{\pi} 2^{-\alpha+\lambda-\frac{1}{2}}}{\sin(-\pi\alpha) \Gamma(-\alpha+\lambda+\frac{1}{2})} \mathbf{S}_{\alpha,\lambda}(w) + \frac{\sqrt{\pi} 2^{\alpha+\lambda-\frac{1}{2}} (w^2-1)_{\bullet}^{-\alpha}}{\sin \pi\alpha \Gamma(\alpha+\lambda+\frac{1}{2})} \mathbf{S}_{-\alpha,\lambda}(w), \quad (\text{C.22})$$

$$\mathbf{S}_{\alpha,\lambda}(-w) = \frac{\cos \pi\lambda}{\sin(-\pi\alpha)} \mathbf{S}_{\alpha,\lambda}(w) + \frac{\pi (1-w^2)^{-\alpha}}{\sin \pi\alpha \Gamma(\frac{1}{2}+\alpha+\lambda) \Gamma(\frac{1}{2}+\alpha-\lambda)} \mathbf{S}_{-\alpha,\lambda}(w). \quad (\text{C.23})$$

Note that the first connection formula is valid when $w \notin]-\infty, 1]$, and the second connection formula is valid when $w \notin]-\infty, -1] \cup [1, \infty[$. Here we borrow a notation from [DGR] where

$$(w^2-1)_{\bullet}^{\alpha} := (w-1)^{\alpha} (w+1)^{\alpha}. \quad (\text{C.24})$$

The functions $(w^2-1)^{\alpha}$ and $(w^2-1)_{\bullet}^{\alpha}$ coincide only if $\text{Re}(w) > 0$. In general the function $(w^2-1)_{\bullet}^{\alpha}$ is holomorphic on $\mathbb{C} \setminus]-\infty, 1]$ while $(w^2-1)^{\alpha}$ is holomorphic on $\mathbb{C} \setminus \{[-1, 1] \cup i\mathbb{R}\}$.

C.5 Whipple transformation

Gegenbauer equation has an extra symmetry compared to hypergeometric symmetry called Whipple transformation. On the level of its standard solutions it has the following form:

$$\mathbf{Z}_{\alpha,\lambda}(w) = (w^2-1)_{\bullet}^{-\frac{1}{4}-\frac{\alpha}{2}-\frac{\lambda}{2}} \mathbf{S}_{\lambda,\alpha} \left(\frac{w}{(w^2-1)_{\bullet}^{\frac{1}{2}}} \right), \quad (\text{C.25})$$

$$\mathbf{S}_{\alpha,\lambda}(w) = (w^2-1)_{\bullet}^{-\frac{1}{4}-\frac{\alpha}{2}-\frac{\lambda}{2}} \mathbf{Z}_{\lambda,\alpha} \left(\frac{w}{(w^2-1)_{\bullet}^{\frac{1}{2}}} \right), \quad \text{Re}(w) > 0. \quad (\text{C.26})$$

(C.26) is obtain by inverting (C.25) and using the fact that $w \mapsto \frac{w}{(w^2-1)_{\bullet}^{\frac{1}{2}}}$ is an involution if and only if $\text{Re}(w) > 0$.

C.6 Half integer case

Gegenbauer functions with $\alpha = \pm \frac{1}{2}$ have simple expressions in terms of elementary functions. To see this we change variables in Gegenbauer operators. For $w \in]-1, 1[$, we substitute $w = \cos \phi$:

$$\mathcal{G}_{-\frac{1}{2},\lambda}(w, \partial_w) = \partial_{\phi}^2 + \lambda^2; \quad (\text{C.27})$$

$$\mathcal{G}_{\frac{1}{2},\lambda}(w, \partial_w) = \frac{1}{\sin \phi} (\partial_{\phi}^2 + \lambda^2) \sin \phi. \quad (\text{C.28})$$

For $w \in]1, +\infty[$, we substitute $w = \cosh \theta$:

$$\mathcal{G}_{-\frac{1}{2},\lambda}(w, \partial_w) = -\partial_{\theta}^2 + \lambda^2; \quad (\text{C.29})$$

$$\mathcal{G}_{\frac{1}{2},\lambda}(w, \partial_w) = \frac{1}{\sinh \theta} (-\partial_{\theta}^2 + \lambda^2) \sinh \theta. \quad (\text{C.30})$$

(C.27)–(C.30) easily imply the first column of the following identities for Gegenbauer functions. (The second column simply follows from the definitions of Gegenbauer functions).

$$\cos \lambda \phi = S_{-\frac{1}{2}, \lambda}(\cos \phi) = F\left(\lambda, -\lambda; \frac{1}{2}; \sin^2 \frac{\phi}{2}\right), \quad (\text{C.31})$$

$$\frac{\sin \lambda \phi}{\lambda \sin \phi} = S_{\frac{1}{2}, \lambda}(\cos \phi) = F\left(1 + \lambda, 1 - \lambda; \frac{3}{2}; \sin^2 \frac{\phi}{2}\right), \quad (\text{C.32})$$

$$\cosh \lambda \theta = S_{-\frac{1}{2}, \lambda}(\cosh \theta) = F\left(\lambda, -\lambda; \frac{1}{2}; \sinh^2 \frac{\theta}{2}\right), \quad (\text{C.33})$$

$$\frac{\sinh \lambda \theta}{\lambda \sinh \theta} = S_{\frac{1}{2}, \lambda}(\cosh \theta) = F\left(1 + \lambda, 1 - \lambda; \frac{3}{2}; \sinh^2 \frac{\theta}{2}\right), \quad (\text{C.34})$$

$$2^\lambda e^{-\lambda \theta} = Z_{-\frac{1}{2}, \lambda}(\cosh \theta) = \left(2 \sinh^2 \frac{\theta}{2}\right)^{-\lambda} F\left(\lambda, \lambda + \frac{1}{2}; 1 + 2\lambda; -\frac{1}{\sinh^2 \frac{\theta}{2}}\right), \quad (\text{C.35})$$

$$\frac{2^\lambda e^{-\lambda \theta}}{\sinh \theta} = Z_{\frac{1}{2}, \lambda}(\cosh \theta) = \left(2 \sinh^2 \frac{\theta}{2}\right)^{-\lambda-1} F\left(\lambda + \frac{1}{2}, \lambda + 1; 1 + 2\lambda; -\frac{1}{\sinh^2 \frac{\theta}{2}}\right). \quad (\text{C.36})$$

The above formulas can be found e.g. in equations (4.25)–(4.28) of [DGR] (in a slightly different form). They are straightforward generalizations of well-known formulas for Chebyshev polynomials. In fact, for $n \in \mathbb{N}_0$ the usual Chebyshev polynomials are special cases of Gegenbauer functions with $\alpha = \pm \frac{1}{2}$:

$$T_n(w) = S_{-\frac{1}{2}, n}(w), \quad U_n(w) = (n+1)S_{\frac{1}{2}, n+1}(w). \quad (\text{C.37})$$

D Closed realizations of 1d Schrödinger operators

D.1 Minimal and maximal realization

The theory of self-adjoint realizations of 1d Schrödinger operators with real potentials is well-known and discussed in various sources [GeZin, GTV]. Somewhat less known is the theory of their closed realizations, which allows for complex potentials—however it is also a classic subject covered in various texts [DuSch, EE, DeGe]. We will treat [DeGe] as the basic source for this topic. It is concisely repeated in Sect. 2 of [DL].

For the convenience of the reader let us summarize some points from [DeGe, DL].

We consider an interval $]a, b[$ and an operator L acting on $f \in C_c^\infty]a, b[$ given by (1.1), that is

$$Lf := (-\partial_x^2 + V(x))f. \quad (\text{D.1})$$

A *closed realization* of L is a closed operator L_\bullet in the sense of $L^2]a, b[$ that restricted to $C_c^\infty]a, b[$ coincides with L .

There always exists the maximal closed realization, denoted L^{\max} and the minimal closed realization, denoted L^{\min} . L^{\max} restricted to $\mathcal{D}(L^{\min})$ coincides with L^{\min} .

In many cases $\mathcal{D}(L^{\min}) = \mathcal{D}(L^{\max})$, and then there exists a unique closed realization of L .

Sometimes $\mathcal{D}(L^{\max})$ is larger than $\mathcal{D}(L^{\min})$, and then there exist also closed realizations of L , denote them L_\bullet , which satisfy

$$\mathcal{D}(L^{\min}) \subset \mathcal{D}(L_\bullet) \subset \mathcal{D}(L^{\max}) \quad (\text{D.2})$$

and $L_\bullet = L^{\max} \Big|_{\mathcal{D}(L_\bullet)}$.

D.2 Resolvent

Let L_\bullet be a realization of L with separated boundary conditions. Following [DL, DeGe] we will now sketch how to find the spectrum of L_\bullet , denoted $\sigma(L_\bullet)$, and how to compute the integral kernel of $\frac{1}{L_\bullet - z}$ for $z \in \mathbb{C}$ outside of $\sigma(L_\bullet)$.

First let us recall the definition of the Wronskian of two complex functions Φ_1, Φ_2 on $]a, b[$:

$$\mathcal{W}(\Phi_1, \Phi_2)(x) := \Phi_1(x)\Phi_2'(x) - \Phi_1'(x)\Phi_2(x). \quad (\text{D.3})$$

It is easily checked that if both Φ_1 and Φ_2 are eigenfunctions of $-\partial_x^2 + V(x)$ with the same eigenvalue, then $\mathcal{W}(\Phi_1, \Phi_2)(x)$ does not depend on x , so that we can write $\mathcal{W}(\Phi_1, \Phi_2)$.

Consider a closed realization of L , denoted L_\bullet . Let $\mathcal{D}(L_\bullet) \subset L^2]a, b[$ denote the domain of L_\bullet . Suppose $z \in \mathbb{C}$ and let $\Psi_a(z, \cdot), \Psi_b(z, \cdot)$ be functions in $AC^1]a, b[$ solving the eigenvalue equation

$$(-\partial_x^2 + V(x) - z)\Psi_a(z, x) = 0, \quad (\text{D.4})$$

$$(-\partial_x^2 + V(x) - z)\Psi_b(z, x) = 0, \quad (\text{D.5})$$

$\Psi_a(z, \cdot)$ is in $\mathcal{D}(L_\bullet)$ near a and $\Psi_b(z, \cdot)$ is in $\mathcal{D}(L_\bullet)$ near b . Set

$$\mathcal{W}(x) := \mathcal{W}(\Psi_b(z, \cdot), \Psi_a(z, \cdot)). \quad (\text{D.6})$$

Define the integral kernel

$$R_\bullet(z; x, y) := \frac{1}{\mathcal{W}(z)} \begin{cases} \Psi_a(z, x)\Psi_b(z, y) & \text{if } a < x < y < b, \\ \Psi_a(z, y)\Psi_b(z, x) & \text{if } a < y < x < b. \end{cases} \quad (\text{D.7})$$

Note that (D.7) does not depend on the choice of $\Psi_a(z, \cdot)$ and $\Psi_b(z, \cdot)$.

Suppose that (D.7) defines a bounded operator, which we denote $R_\bullet(z)$. Then

$$z \notin \sigma(L_\bullet) \quad \text{and} \quad \frac{1}{L_\bullet - z} = R_\bullet(z). \quad (\text{D.8})$$

Conversely, if $z \notin \sigma(L_\bullet)$, then the functions Ψ_a, Ψ_b with the above properties exist and the operator $R_\bullet(z)$ is bounded.

Above we described the path from L_\bullet to $R_\bullet(z)$. In our constructions we will use the reverse direction, from $R_\bullet(z)$ to L_\bullet , which is described in the following theorem.

Theorem D.1. *Suppose that $z \in \mathbb{C}$ and let $\Psi_a(z, \cdot), \Psi_b(z, \cdot)$ be functions in $AC^1]a, b[$ solving the eigenvalue equations (D.4), resp. (D.5), such that $\Psi_a(z, \cdot)$ is in L^2 near a and $\Psi_b(z, \cdot)$ is in L^2 near b . Define $\mathcal{W}(x)$ by (D.6) and $R_\bullet(z; x, y)$ by (D.7). Suppose that (D.7) defines a bounded operator on $L^2]a, b[$, which we denote $R_\bullet(z)$. Then there exists a unique operator closed L_\bullet that satisfies (D.8).*

E Associated Legendre functions vs. Gegenbauer functions

In the literature, many authors use a Legendre function instead of Gegenbauer functions. Here, we briefly discuss the relations between these functions and Gegenbauer functions. For Legendre function we use [NIST2] as our reference and [D1] as our reference for Gegenbauer function.

The Legendre differential operator is

$$\mathcal{L}_\mu^\alpha := (1 + z^2)\partial_z^2 - 2z\partial_z + \mu(\mu + 1) - \frac{\alpha^2}{1 - z^2}. \quad (\text{E.1})$$

It is equivalent to the Gegenbauer operator, in fact

$$\begin{aligned} & (1-w^2)^{\mp\frac{\alpha}{2}} \mathcal{L}_\mu^\alpha (1-w^2)^{\pm\frac{\alpha}{2}} \\ & = (1-w^2) \partial_w^2 - 2(\pm\alpha+1)w\partial_w + (\mu \mp \alpha)(\mu \pm \alpha + 1) = \mathcal{G}_{\pm\alpha, \mu + \frac{1}{2}}. \end{aligned} \quad (\text{E.2})$$

Certain distinguished functions annihilated by this operator are called associated Legendre functions. There various choices for these functions, which we quote following [NIST2]:

The associated Legendre function of the first kind is

$$\begin{aligned} \mathbf{P}_\mu^\alpha(z) &= \left(\frac{z+1}{z-1}\right)^{\frac{\alpha}{2}} \mathbf{F}\left(\mu+1, -\mu; 1-\alpha; \frac{1-z}{2}\right) \\ &= \frac{2^\alpha}{(z^2-1)^\alpha} \mathbf{S}_{-\alpha, \mu + \frac{1}{2}}(z) \end{aligned} \quad (\text{E.3})$$

The Ferrers function of the first kind is

$$\begin{aligned} \mathcal{P}_\mu^\alpha(z) &= \left(\frac{z+1}{1-z}\right)^{\frac{\alpha}{2}} \mathbf{F}\left(\mu+1, -\mu; 1-\alpha; \frac{1-z}{2}\right) \\ &= \frac{2^\alpha}{(1-z^2)^\alpha} \mathbf{S}_{-\alpha, \mu + \frac{1}{2}}(z) \end{aligned} \quad (\text{E.4})$$

And **the associated Legendre function of the second kind** is

$$\begin{aligned} \mathbf{Q}_\mu^\alpha(z) &= e^{i\pi\alpha} \frac{\pi^{\frac{1}{2}} \Gamma(\alpha + \mu + 1) (z^2 - 1)^{\frac{\alpha}{2}}}{2^{\mu+1} z^{\alpha+\mu+1}} \mathbf{F}\left(\frac{\alpha + \mu}{2} + 1, \frac{\alpha + \mu}{2} + \frac{1}{2}; \frac{3}{2} + \mu; z^{-2}\right) \\ &= e^{i\pi\alpha} \frac{\pi^{\frac{1}{2}} \Gamma(\alpha + \mu + 1) (z^2 - 1)^{\frac{\alpha}{2}}}{2^{\mu+1}} \mathbf{Z}_{\alpha, \mu + \frac{1}{2}}(z). \end{aligned} \quad (\text{E.5})$$

Note that Legendre functions are represented with upper and lower indices. One should not confuse them with other, quite analogous functions defined in the text, whose parameters are lower indices.

The Legendre functions are closely related to the functions \mathcal{P}^s , \mathcal{P}^h , and \mathcal{Q}^h that we introduced in the section on Gegenbauer Hamiltonians, which you can see on the right of the following comparison:

$$\mathcal{P}_\mu^\alpha(\cos r) = \left(\frac{2}{\sin r}\right)^{\frac{1}{2}} \mathcal{P}_{-\alpha, \mu + \frac{1}{2}}^s(r), \quad (\text{E.6})$$

$$\mathbf{P}_\mu^\alpha(\cosh r) = \left(\frac{2}{\sinh r}\right)^{\frac{1}{2}} \mathcal{P}_{-\alpha, \mu + \frac{1}{2}}^h(r), \quad (\text{E.7})$$

$$\mathbf{Q}_\mu^\alpha(\cosh r) = \frac{1}{(2 \sinh r)^{\frac{1}{2}}} e^{i\pi\alpha} \sqrt{\pi} \Gamma(\alpha + \mu + 1) \mathcal{Q}_{-\alpha, \mu + \frac{1}{2}}^h(r). \quad (\text{E.8})$$

Conflict of interest. This manuscript has no conflict of interest.

References

- [Bos] Bose A.K.: *A class of solvable potentials*, Nuov. Cim., 32:679 (1964)
- [Cay] Cayley A.: *On the Schwarzian derivative and the polyhedral functions*, Trans. Camb. Phil. Soc., 13 (1880)
- [Car] R.W. Carroll, *Transmutation and Operator Differential Equations*, Mathematics Studies, Vol. **37**, North Holland (1979).

- [CKS] Cooper F., Khare A., Sukhatme U.: *Supersymmetry in Quantum Mechanics*, World Scientific, Singapore (2001)
- [Cot] Cotfas N.: *Shape-invariant hypergeometric type operators with application to quantum mechanics*, Cent. Eur. J. Phys. 4, 318 (2006)
- [D1] Dereziński J.: *Hypergeometric type functions and their symmetries*, Annales Henri Poincaré 15 1569-1653(2014) DOI: 10.1007/s00023-013-0282-4
- [D2] Dereziński J.: *Group-theoretical origin of symmetries of hypergeometric class equations and functions*, Complex differential and difference equations. DOI: 10.1515/9783110611427-001
- [DGR] Dereziński J., Gaß C., Ruba B.: *Generalized integrals of Macdonald and Gegenbauer functions.*, in "Applications and q-Extensions of Hypergeometric Functions", Contemporary Mathematics, vol. 819, Amer. Math. Soc., Providence, RI, 2025, pp. 3-41
- [DeGa] Dereziński J., Gaß C.: *Propagators in curved spacetimes from operator theory*, arXiv:2409.03279
- [DeRi] Dereziński, J., Richard, S.: *On radial Schrödinger operators with a Coulomb potential*, Annales Henri Poincaré 19 (2018), 2869-2917,
- [DeGe] Dereziński, J., Georgescu, V.: *One-dimensional Schrödinger operators with complex potentials*, Ann. Henri Poincaré 21, 1947–2008 (2020).
- [DL] Dereziński, J., Lee, J.: *Exactly Solvable Schrödinger Operators Related to the Confluent Equation*, Milan Journal of Mathematics 2025
- [DW] Dereziński J., Wrochna M.: *Exactly solvable Schrodinger operators*, Annales Henri Poincaré 12 (2011) 397-418
- [DW1] J. Dereziński and M. Wrochna, *Continuous and holomorphic functions with values in closed operators*, Journ. Math. Phys. **55**, 083512 (2014).
- [DuSch] Dunford, N., Schwartz, J. T.: *Linear operators, vol. III, Spectral operators*, Chap. XX.1.1 Spectral differential operators of second order. Wiley Interscience (1971).
- [DK] Duru I.H., Kleinert H.: *Solution of the path integral for the H-atom*, Physics Letters B 84 (2): 185-188, (1979)
- [Eck] Eckart C.: *The Penetration of a Potential Barrier by Electrons*, Phys. Rev., 35:1303 (1930)
- [EE] Edmunds, D. E., Evans, W. D.: *Spectral theory and differential operators*, 2nd ed. Oxford University Press (2018).
- [Eve] Everitt, W. N.: *A Catalogue of Sturm-Liouville Differential Equations*, in: *Sturm-Liouville Theory Past and Present*, Amrein, W. O., Hinz, A. M., Pearson, D. B. (Eds.), Birkhäuser (2000).
- [Flü] Flüge S.: *Practical Quantum Mechanics*, Springer-Verlag, Berlin (1971)
- [FS] Fucci, G., Stanfill, J.: *The exotic structure of the spectral -function for the Schrödinger operator with Pöschl-Teller potential*, arXiv:2411.17860
- [GPLS] Gesztesy, F., Littlejohn, L.L., Piorkowski, M., Stanfill, J.: *The Jacobi operator on $(-1, 1)$ and its various m -functions*, arXiv:2307.12164
- [GeZin] Gesztesy, F., Nichols, R., Zinchenko, M.: *Sturm-Liouville operators, their spectral theory, and some applications*, AMS, Colloquium Publications, vol. 67 (2024).

- [GTV] Gitman, D. M., Tyutin, I. V., Voronov, B. L.: *Self-adjoint extensions in quantum mechanics. General theory and applications to Schrödinger and Dirac equations with singular potentials*, Prog. Math. Phys. 62, Birkhäuser/Springer, New York (2012).
- [Gin] Ginocchio J.N.: *A class of exactly solvable potentials. I. One-dimensional Schrödinger equation*, Ann. Phys., 152:203 (1984)
- [HIU] Honda K., Ikawa T. AND Udagawa S.: *On Complex Spheres*, Mem. Fac. Sci. Eng. Shimane Univ. Series B: Mathematical Science 36 (2003)
- [HI] Hull T.E, Infeld L.: *The factorization method*, Rev. Mod. Phys. 23 2168 (1951)
- [Hul] Hulthén L.: *Über die Eigenlösungen der Schrödinger-Gleichung des Deuterons*, Ark. Mat. Astron. Fys. 28 A (1942)
- [Ince] Ince, E.L. *Ordinary Differential Equations*, Longmans, Green and Co, 1926
- [Kato] T. Kato, *Perturbation Theory for Linear Operators*, Springer (1966).
- [Koo] Koorwinder T.H., *Jacobi functions and analysis on noncompact semisimple Lie groups*, In: Askey, R.A., Koornwinder, T.H., Schempp, W. (eds) *Special Functions: Group Theoretical Aspects and Applications. Mathematics and Its Applications*, vol 18. Springer, Dordrecht. DOI: 10.1007/978-94-010-9787-1.1
- [Kum] Kummer E.: *Über die hypergeometrische Reihe*, Crelle 15:39-83,127-172 (1836)
- [KS] Khare A., Sukhatme U.: *Is the lowest order supersymmetric WKB approximation exact for all shape invariant potentials?* Jour. Phys. A26 (1993)
- [LeB] LeBrun C.: *Spaces of Complex Null Geodesics in Complex-Riemannian Geometry*, Transactions of the American Mathematical Society, 278(1983).
- [Lio] Liouville J.: *Solution d'un probleme d'analyse*, J. Math. Pures Appl., 2, 16-35 (1837)
- [M1] Miller, W.: *Lie Theory and Special Functions*, 1968., Academic Press, New York, London
- [Mil] Milson R.: *On the Liouville transformation and exactly-solvable Schrödinger equations*, Int. J. Theor. Phys. 37, 1735 (1998)
- [Mor] Morse P.M.: *Diatomic Molecules According to the Wave Mechanics. II. Vibrational Levels*, Phys. Rev. (1929)
- [MaR] Manning M.F., Rosen N.: *A Potential Function for the Vibrations of Diatomic Molecules*, Phys. Rev., 44:953 (1933)
- [MoR] Morse, P. M., Rosen, N.: *On the Vibrations of Polyatomic Molecule*, PhysRev.42.210,(1932)
- [Nat] Natanzon G.A.: *Study of the one-dimensional Schroedinger equation generated from the hypergeometric equation*, Vestnik Leningrad. Univ., 10:22 (1971)
- [NIST1] Olde Daalhuis A. B., National Institute of Standards and Technology, *Hypergeometric Function*, Federal Information Processing Standards Publications (FIPS) 140-2, DOI = 10.6028/nist.fips.140-2 (2001).
- [NIST2] Dunster T. M., National Institute of Standards and Technology, *Legendre and Related Functions*, Federal Information Processing Standards Publications (FIPS) 140-2, DOI = 10.6028/nist.fips.140-2 (2001).

- [NU] Nikiforov A.F., Uvarov V.B.: *Special Functions of Mathematical Physics*, Birkhäuser, Basel (1988)
- [Osg] Osgood B., *Old and new on the Schwarzian derivative, Quasiconformal Mappings and Analysis*, Springer, New York, pp.275-308 (1998)
- [PT] Pöschl G., Teller E.: *Exact Quantized Momentum Eigenvalues and Eigenstates of a General Potential Model*, Z. Physik (1933)
- [R] Rainville, E. D.: *Special Functions*, The Macmillan Co., New York (1960)
- [Sca] Scarf F.: *New Soluble Energy Band Problem*, Phys. Rev. 112, 1137 (1958)
- [Sch] Schwarz H.: *Ueber diejenigen Faelle, in welchen die Gaussische hypergeometrische Reihe eine algebraische Function ihres vierten Elementes darstellt*, J. Math. 75:292-335 (1873)
- [S] Schrödinger E.: *A method of determining quantum-mechanical eigenvalues and eigenfunctions*, Proc. Roy. Irish Acad. A46 (1940)
- [SS] Elina Shishkina, Sergei Sitnik. *Transmutations, Singular and Fractional Differential Equations with Applications to Mathematical Physics*, Series: Mathematics in Science and Engineering. 1st Edition. Elsevier. Academic Press. 592 P.
- [SW] Saxon D.S., Woods R.D.: *Diffuse surface optical model for nucleon-nuclei scattering*, Physical Review 95, 577-578 (1954)
- [V] Vilenkin, N. Ya.: *Special Functions and the Theory of Group Representations*, Translations of Mathematical Monographs, AMS, Providence 1968
- [Wa] Wawrzyńczyk, A., *Modern Theory of Special Functions*, PWN, Warszawa 1978 (Polish)
- [WW] Whittaker E.T., Watson G.N.: *A course of Modern Analysis, vol I, II*, 4th edition (reprint of the 1927 edition), Cambridge Univ. Press, New York (1962)