

## Problem Set 7 – Statistical Physics B

### Problem 1: Derivation of the virial expansion

- (a) Derive the virial expansion for the pressure up until  $\mathcal{O}(\rho^4)$ . You might want to consider the strategy from the lecture notes. Give explicit expressions for  $B_2(T)$  and  $B_3(T)$  in terms of the Mayer function.
- (b) Suppose one would add a three-body potential to the microscopic classical Hamiltonian. Will the second virial coefficient depend on this three-body potential? Prove your answer.
- (c) Derive the virial expansion of the radial distribution function for a pair-wise additive interacting classical system,

$$g(r; \rho, T) = g^{(0)}(r; T) + \rho g^{(1)}(r; T) + \dots \quad (1)$$

and give an explicit expression for the expansion coefficients in terms of the Mayer function. What is the physical interpretation of these expansion coefficients?

### Problem 2: The second virial coefficient for model potentials

We recall the definition of the second virial coefficient,  $B_2(T) = -(1/2) \int d\mathbf{r} f_M(r)$ , with  $f_M$  the Mayer function.

- (a) Compute  $B_2(T)$  for a hard-sphere system and for a square-well fluid. Under which conditions does the square-well fluid reduce to the hard-sphere limit? Check that this is also reflected in the expression of  $B_2(T)$ . What is the Boyle temperature in both cases? Give a physical explanation.
- (b) Show that under certain conditions

$$B_2(T) = -\frac{1}{6k_B T} \int_0^\infty dr 4\pi r^3 v'(r) \exp[-\beta v(r)]. \quad (2)$$

What are these conditions? Compute  $B_2(T)$  for potentials of the form  $v(r) = \alpha/r^n$  with  $n > 3$  in terms of the Euler Gamma function.

### Problem 3: Third virial coefficient for hard spheres

The third virial coefficient is given by

$$B_3(T) = -\frac{1}{3V} \int d\mathbf{r}_1 \int d\mathbf{r}_2 \int d\mathbf{r}_3 f_M(r_{12}) f_M(r_{13}) f_M(r_{23}). \quad (3)$$

- (a) Give a geometric interpretation of this formula in the case of hard-sphere interactions. Using these geometric arguments, compute  $B_3$  for hard spheres.
- (b) Introduce the Fourier transform  $\tilde{f}_M(k) = \int d\mathbf{r} f_M(r) e^{-i\mathbf{k}\cdot\mathbf{r}}$  of the Mayer function and show that

$$B_3(T) = -\frac{1}{3} \int \frac{d\mathbf{k}}{(2\pi)^3} \tilde{f}_M(k)^3. \quad (4)$$

Derive an expression for  $\tilde{f}_M(k)$  for hard spheres, and then compute  $B_3(T)$  from it. The following integral might be useful,

$$\int_0^\infty dx x^{-5/2} J_{3/2}(x)^3 = \frac{5}{48\sqrt{2\pi}}, \quad (5)$$

where  $J_\nu(x)$  is  $\nu$ -th order Bessel function of the first kind.

#### Problem 4: Lennard-Jonesium

- (a) Consider a Lennard-Jones fluid. Explain the terms in this potential. What is the minimum of the potential? Give a physical interpretation. Compute the force acting on a particle. What is the direction of the force? When is the force maximal/minimal?
- (b) Give an expression of the ratio  $B^* = B_2^{\text{LJ}}/B_2^{\text{HS}}$  in terms of dimensionless temperature  $T^* = k_B T/\epsilon$ . By a suitable variable substitution, show that

$$B^*(T^*) = \frac{8}{\sqrt{2}T^*} e^{1/T^*} \frac{1}{2} \int_0^\infty du \frac{u-1}{\sqrt{u}} e^{-(1/T^*)(u-1)^2}. \quad (6)$$

For which temperature does the Lennard-Jones fluid reduce to the hard-sphere fluid based on the second virial coefficient?

- (c) Compute the integral numerically or analytically. For an analytical calculation, you might want to consider the following steps:
- Use the coordinate transformations  $u = 1 + \cosh(t/2)$  for  $u > 2$  and  $u = 1 + \cos(t/2)$  for  $0 < u < 2$ .
  - Express your result in terms of modified Bessel functions of the first kind. The following integral representation might be useful:

$$I_\nu(x) = \frac{1}{\pi} \int_0^\pi dt e^{x \cos t} \cos(\nu t) - \frac{\sin(\pi\nu)}{\pi} \int_0^\infty dt e^{-x \cosh t - \nu t}. \quad (7)$$

- Obtain the final result

$$B^*(T^*) = \frac{\sqrt{2}\pi}{2T^*} e^{1/(2T^*)} \sum_{n=0,1} (-1)^{n+1} \left[ I_{(2n+1)/4} \left( \frac{1}{2T^*} \right) + I_{-(2n+1)/4} \left( \frac{1}{2T^*} \right) \right]. \quad (8)$$

- (d) Determine numerically the Boyle temperature as a function of  $\epsilon$  for a Lennard-Jones fluid.