

## Problem Set 9 – Statistical Physics B

### Problem 1: An ideal gas in an external potential

- (a) Write down the grand potential  $\Omega$  for  $N$  identical particles interacting via a pairwise additive potential  $v(r)$  in an external field  $V_{\text{ext}}(\mathbf{r})$ . Viewing  $\Omega$  as a functional of the intrinsic chemical potential  $u(\mathbf{r})$  or  $v(\mathbf{r}, \mathbf{r}')$ , show by explicit functional differentiation that  $\delta\Omega/\delta u(\mathbf{r}) = -\rho(\mathbf{r})$  and  $\rho^{(2)}(\mathbf{r}, \mathbf{r}') = 2\delta\Omega/\delta v(\mathbf{r}, \mathbf{r}')$ .
- (b) Prove that
- $$\langle \delta\hat{\rho}(\mathbf{r}_1) \dots \delta\hat{\rho}(\mathbf{r}_n) \rangle = -\frac{\delta^n \beta\Omega[u]}{\delta\beta u(\mathbf{r}_1) \dots \delta\beta u(\mathbf{r}_n)}, \quad n \geq 2. \quad (1)$$
- (c) Consider an ideal gas,  $v = 0$ . Derive from  $\Omega[u]$  an expression for  $\rho(\mathbf{r})$  in an external potential.
- (d) Write using DFT a formal expression for constancy of the chemical potential. From it, determine  $\mathcal{F}[\rho]$  for an ideal gas. Show that it is of the local form.
- (e) Show that  $\langle \beta^{-1} \ln f_N \rangle = -TS$ , with  $T$  temperature,  $S$  entropy, and  $f_N$  the grand-canonical probability distribution. Argue that  $\mathcal{F}[\rho]$  is indeed the intrinsic Helmholtz free energy. Does this interpretation depend on the type of interaction potential?
- (f) Compute  $\langle \delta\hat{\rho}(\mathbf{r}_1) \dots \delta\hat{\rho}(\mathbf{r}_n) \rangle$  and  $c^{(n)}(\mathbf{r}_1, \dots, \mathbf{r}_n)$  for an ideal gas in an external field.

### Problem 2: Sedimentation in the local density approximation

In the local density approximation (LDA) we set  $\mathcal{F}[\rho] = \int d\mathbf{r} f(\rho(\mathbf{r}))$ , with  $f(\rho_b)$  the Helmholtz free energy density of a homogeneous bulk system at density  $\rho_b$ . The LDA thus assumes that the *functional* form of the free energy density of a bulk system is also valid for an inhomogeneous system.

- (a) Show that the equilibrium density profiles satisfy the Euler-Lagrange equation  $f'(\rho(\mathbf{r})) + V_{\text{ext}}(\mathbf{r}) = \mu$ , with prime denoting differentiation to the argument. Give a physical interpretation for  $f'(\rho_b)$ .
- (b) Rewrite your answer from (a) as  $\nabla p(\rho(\mathbf{r})) = -\rho(\mathbf{r})\nabla V_{\text{ext}}(\mathbf{r})$ . Prove that this is equivalent to the condition for hydrostatic equilibrium.
- (c) Calculate within the local-density approximation  $c(\mathbf{r}, \mathbf{r}')$  and  $h(\mathbf{r}, \mathbf{r}')$  and show that both are proportional to  $\delta(\mathbf{r} - \mathbf{r}')$  reflecting the local nature of the approximation.
- (d) Take  $V_{\text{ext}}(\mathbf{r}) = mgz$ , with  $m$  the mass,  $g$  the gravitational acceleration and  $z$  the altitude. This describes the situation where particles sediment in an external gravitational field. Under what conditions do you expect that the local-density approximation describes this situation accurately?
- (e) Prove that

$$\frac{d \ln[\rho(z)\mathcal{V}]}{dz} = -\frac{\bar{\kappa}_T(\rho(z))}{\ell_g}, \quad \bar{\kappa}_T(\rho_b) = \left[ \beta\rho \left( \frac{\partial\mu}{\partial\rho} \right) \right]^{-1} \bigg|_{\rho=\rho_b}, \quad (2)$$

for some constant  $\mathcal{V}$  with dimensions of volume. Give an expression for  $\ell_g$ . What is the physical interpretation of  $\bar{\kappa}_T$ ?

- (f) Compute the density profile for an ideal gas. How is the integration constant determined?
- (g) Suppose we have measured the density profile of some system. Argue that from this information we can obtain the equation of state for the system.
- (h) Let us describe the situation where colloidal spheres are sedimenting in some simple fluid with density profile  $\rho(z)$ . How do the above considerations change? Hint: Because the colloidal particles are much larger than the particles in the "solvent", you can assume the solvent to be a structureless medium with given mass density  $\rho_s$ . Derive an expression for the external potential on a single particle using the equation for hydrostatics.