

Super Cartan geometry, LQG and applications

Konstantin Eder

FAU Erlangen-Nürnberg

(joint collaboration with Hanno Sahlmann)

April 22, 2022

Section 1

Introduction

Asthekar's discovery:

- GR can be formulated such that (non-reduced) symplectic phase space of canonical theory is that of $SU(2)$ Yang-Mills!
- Also exist in higher dimensions [\[Bodendorfer+Thiemann+Thurn\]](#)

In 4D: **Holst action** S_{Holst}^β : 1-parameter family of *deformed* actions of first-order Einstein gravity

Holst action

$$S_{\text{Holst}}^\beta = \frac{1}{4\kappa} \int_M \epsilon_{IJKL} e^I \wedge e^J \wedge F[\omega]^{KL} + \frac{2}{\beta} e^I \wedge e^J \wedge F[\omega]_{IJ}, \quad \beta : \text{Immirzi}$$

$I, J, \dots : \text{SO}(1, 3)$

- \rightarrow yield **same** EOM as Einstein gravity! (metric $g_{\mu\nu} = \eta_{IJ} e_\mu^I e_\nu^J$)

Canonical decomposition

- Assume spacetime is globally hyperbolic: $M = \mathbb{R} \times \Sigma$ (Σ : 3D Cauchy slice)

3+1-decomposition

$$S_{\text{Holst}}^{\beta} = \frac{1}{\kappa\beta} \int_{\mathbb{R}} dt \int_{\Sigma_t} d^3x \left(E_i^a \dot{A}_a^i - \omega_0^i G_i + N^a C_a + NC \right)$$

i,j,\dots : SO(3) (or SU(2))


- Canonically conjugate variables [\[Ashtekar+Barbero+Immirzi\]](#)

$$A_a^i = \Gamma_a^i + \beta K_a^i, \quad E_i^a = \sqrt{|\det q|} e_i^a$$

- Constraints: Gauss G_i , Diffeo C_a and Hamilton C
- \rightarrow Constraints+Hamilton EOM \Leftrightarrow Einstein equations

Quantum theory


Basic building blocks: **Spin networks** [Penrose 70's] → Spin nets associated to gauge-invariant functionals.



→ Form basis of Hilbert space [Ashtekar+Lewandowski]

$$\mathcal{H}_{\text{LQG}} = \lim_{\rightarrow} \mathcal{H}_{\gamma} = L^2(\overline{\mathcal{A}}/\overline{\mathcal{G}}, d\mu_{\text{AL}})$$

Example:



$$\Psi[A] = \pi_{j_1}(h_{e_1}[A])^{m_1}_{n_1} \pi_{j_2}(h_{e_2}[A])^{m_2}_{n_2} \pi_{j_3}(h_{e_3}[A])^{m_3}_{n_3} \iota_{m_1 m_2 m_3} \iota^{n_1 n_2 n_3}$$

Why SUSY in LQG?

Motivation:

- 1 If a quantum theory of gravity really exists, it has to be unique!
- 2 Apply ideas + techniques from one approach to the other

How to achieve this?

- Essential ingredient: **Supersymmetry**
- **Supergravity theories** as low-energy limits of certain superstring theories
- → Apply LQG methods to supergravity

Coleman-Mandula: most general Lie algebra of symmetries of the S-matrix has form

$$\mathfrak{iso}(\mathbb{R}^{1,3}) \oplus \text{internal sym.}$$

→ Only way around this through new form of symmetry:

Haag-Łopuszański-Sohnius theorem

Have to consider **super Lie algebras**, i.e. \mathbb{Z}_2 -graded algebras $(\mathfrak{g}, [\cdot, \cdot])$ of the form

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \quad \text{with} \quad [\cdot, \cdot] : (\text{anti}) \text{ commutator on } \mathfrak{g}_0 (\mathfrak{g}_1)$$

such that $[\mathfrak{g}_i, \mathfrak{g}_j] \subseteq \mathfrak{g}_{i+j}$ (+ graded Jacobi identity)

Supersymmetry

super Poincaré/super anti-de Sitter $\mathfrak{osp}(1|4)$

$$\mathfrak{iso}(\mathbb{R}^{1,3|4}) = \underbrace{\mathbb{R}^{1,3} \ltimes \mathfrak{so}(1,3)}_{\mathfrak{g}_0} \oplus \underbrace{S_{\mathbb{R}}}_{\mathfrak{g}_1}$$

Generators: P_I , M_{IJ} , Q_α (Majorana spinor)

$$[P_I, Q_\alpha] = 0 - \frac{1}{2L} Q_\beta (\gamma_I)^\beta{}_\alpha$$

$$[M_{IJ}, Q_\alpha] = \frac{1}{2} Q_\beta (\gamma_{IJ})^\beta{}_\alpha$$

$$\{Q_\alpha, Q_\beta\} = \frac{1}{2} (C\gamma_I)_{\alpha\beta} P^I + \frac{1}{4L} (C\gamma^{IJ})_{\alpha\beta} M_{IJ}$$

Further generators $Q_\alpha^r \rightarrow \mathcal{N}$ -extended **SUSY**!

$$AdS_4 := \{x \in \mathbb{R}^5 \mid -(x^0)^2 + (x^1)^2 + \dots + (x^3)^2 - (x^4)^2 = -L^2\}$$

Supersymmetry




Figure: ATLAS Silicon IFIC

Application to gravity

Most ambitious use of this kind of symmetry:

local supersymmetry \Rightarrow **supergravity** (SUGRA)

For $\mathcal{N} = 1$: contains gravitational field and a spin- $\frac{3}{2}$ fermion (*gravitino*)

What has been done?

- Canonical SUGRA adapted to LQG [Jacobson '88]
- Hidden $\mathfrak{osp}(1|2)$ -symmetry in constraint algebra, construction of $\mathfrak{osp}(1|2)$ -valued connection [Fülöp '94, Gambini+Obregon+Pullin '96]
- Quantization [Gambini+Obregon+Pullin '96, Ling+Smolin '99]

But:

- Considerations remain formal
- Origin of hidden symmetry unclear
- Generalizations?

- Canonical theory for higher D and quantization [Bodendorfer +Thiemann+Thurn '11]

But:

- SUSY not manifest
- Formalism/constraints quite complicated

My goals:

Follow [L+S]: Keep SUSY manifest as much as possible!

- Origin of $\mathfrak{osp}(1|2)$ -symmetry? \rightarrow Understand the geometric origins with a view towards generalizations:
 - Immirzi parameters
 - Higher $\mathcal{N} > 1$
 - Boundary theory (\rightarrow **black holes**)
- Mathematically rigorous formulation, both classically and in quantum theory:
 - Construction of super spin-nets (\rightarrow *super parallel transport*)
 - Structure of Hilbert spaces \leftrightarrow relation to standard quantization of fermions in LQG

Classical theory:

- Supergravity via Cartan geometry
- Holst-MacDowell-Mansouri SUGRA action for any β
- Extension to $\mathcal{N} > 1$!
- Boundary theory (\rightarrow black holes!)
- Special properties of self-dual theory

Quantum theory:

- Super spin networks, super area operator
- Applications: black holes, cosmology

Section 2

Gravity as Cartan geometry

F. Klein (Erlanger Programm): "Classify geometry of space via group symmetries".


Example: **Minkowski spacetime** $\mathbb{M} = (\mathbb{R}^{1,3}, \eta)$

- Isometry group $\text{ISO}(\mathbb{R}^{1,3}) = \mathbb{R}^{1,3} \ltimes \text{SO}_0(1, 3)$
- Fix event $p \in \mathbb{M}$: $G_p = \text{SO}_0(1, 3)$ (isotropy subgroup)

Klein geometry

$$\text{ISO}(\mathbb{R}^{1,3})/\text{SO}_0(1, 3) \cong \mathbb{M}$$

Cartan geometry



Gauge field with values in Poincaré algebra

$$A_\mu = \text{pr}_{\mathbb{R}^{1,3}} \circ A_\mu + \text{pr}_{\mathfrak{so}(1,3)} \circ A_\mu =: e_\mu + \omega_\mu$$

- e : co-frame, ω : Lorentz-connection

Cartan geometry


Gauge field with values in Poincaré algebra

$$A_\mu = \text{pr}_{\mathbb{R}^{1,3}} \circ A_\mu + \text{pr}_{\mathfrak{so}(1,3)} \circ A_\mu =: e_\mu + \omega_\mu$$

- e : co-frame, ω : Lorentz-connection

Cartan condition

$\Rightarrow e$ induces **metric** on M : $g_{\mu\nu} = \eta_{IJ} e_\mu^I e_\nu^J$



Gravity as Cartan geometry

Can use Cartan connection A to formulate **YM-type action**:

- Need non-trivial cosmological constant Λ
- \rightarrow Klein geometry: (Anti-)de Sitter space $(A)dS_4$

MacDowell-Mansouri action [M+M '77]

$$S_{\text{MM}}[A] = \frac{3}{\kappa\Lambda} \int_M \langle F[A] \wedge F[A] \rangle$$

- $F[A]$: field strength (curvature) induced by A
- $\langle \cdot, \cdot \rangle$: inner product on (anti-)de Sitter algebra ("generalized trace")
- \rightarrow yields 1st-order gravity + *boundary terms*

Gravity as Cartan geometry

Can use Cartan connection A to formulate **YM-type action**:

- need non-trivial cosmological constant Λ
- \rightarrow Klein geometry: (anti-)de Sitter space $(A)dS_4$

Holst-MacDowell-Mansouri action [Wise '09, KE '21]

$$S_{\text{H-MM}}[A] = \frac{3}{\kappa\Lambda} \int_M \langle F[A] \wedge F[A] \rangle_\beta$$

- $F[A]$: field strength (curvature) induced by A
- $\langle \cdot, \cdot \rangle_\beta$: β -deformed inner product on (anti-)de Sitter algebra induced by

$$\mathcal{P}_\beta := \frac{\mathbb{1} + i\beta\gamma_5}{2\beta}, \quad \beta : \text{Immirzi}$$

Section 3

Supergravity & boundary theory

Supergravity

AdS Supergravity as **super Cartan geometry** modeled on super Klein geometry ($\mathrm{OSp}(1|4)$, $\mathrm{Spin}(1,3)$) [D'Auria-Fré-Regge '80, D'Auria-Fré '80, Castellani-D'Auria-Fré '91, KE '21]

Super Cartan connection

$$\mathcal{A}_\mu : M \rightarrow \mathfrak{osp}(1|4)$$

Decomposition

$$\mathcal{A} = \underbrace{\mathrm{pr}_{\mathfrak{g}_1} \circ \mathcal{A}}_{\psi} + \underbrace{\mathrm{pr}_{\mathbb{R}^{1,3}} \circ \mathcal{A}}_e + \underbrace{\mathrm{pr}_{\mathfrak{spin}(1,3)} \circ \mathcal{A}}_\omega$$

- $\psi = \psi^\alpha Q_\alpha$: (spin-3/2) *Rarita-Schwinger field*

Supergravity and LQG

Holst action for (extended) Poincaré SUGRA [Tsuda '00, Kaul '07]
via MM ($\mathcal{N} = 1$, β as θ -ambiguity) [Obregon+Ortega-Cruz+Sabido '12]
as constrained BF ($\mathcal{N} = 1$) [Kowalski-Glikman+Durka+Szczachor '10]

Here: via Holst projection [KE '21]

- $\mathfrak{osp}(1|4) \cong \mathfrak{sp}(4) \oplus S_{\mathbb{R}}$
- $S_{\mathbb{R}}$ Clifford module \rightarrow can naturally extend \mathcal{P}_{β}

Definition [KE '21]

$$\mathbf{P}_{\beta} := \mathbf{0} \oplus \mathcal{P}_{\beta} \oplus \mathcal{P}_{\beta} : \mathfrak{osp}(1|4) \rightarrow \mathfrak{osp}(1|4)$$

with

$$\mathcal{P}_{\beta} := \frac{\mathbf{1} + i\beta\gamma_5}{2\beta}$$

\rightarrow induces inner product on $\mathfrak{osp}(1|4)$:

$$\langle \cdot, \cdot \rangle_{\beta} := \text{str}(\cdot \mathbf{P}_{\beta} \cdot)$$

Supergravity and LQG

super Holst-MacDowell-Mansouri action [KE '21]

$$S_{\text{SH-MM}}[\mathcal{A}] = \int_M \langle F[\mathcal{A}] \wedge F[\mathcal{A}] \rangle_\beta$$

Curvature

- $F[\mathcal{A}]^I = \Theta^{(\omega)I} - \frac{1}{4} \bar{\psi} \wedge \gamma^I \psi$
 - $F[\mathcal{A}]^{IJ} = R[\omega]^{IJ} + \frac{1}{L^2} \Sigma^{IJ} - \frac{1}{4L} \bar{\psi} \wedge \gamma^{IJ} \psi$
 - $F[\mathcal{A}]^\alpha = D^{(\omega)} \psi^\alpha - \frac{1}{2L} e^I (\gamma_I)^\alpha{}_\beta \wedge \psi^\beta$
-
- \rightarrow yields Holst action of $D = 4$, $\mathcal{N} = 1$ AdS-SUGRA + *bdy terms*
 - **Can show:** Boundary theory **unique** by SUSY-invariance!
[Andrianopoli+D'Auria '14, KE '21]

Extended SUGRA and boundary theory

What about **extended** SUSY? \rightarrow Consider $\mathcal{N} = 2$ [KE '21]

Super Cartan connection

$$\mathcal{A} = e^I P_I + \frac{1}{2} \omega^{IJ} M_{IJ} + \Psi_r^\alpha Q_\alpha^r + \hat{A} T$$

\hat{A} : U(1) gauge field (*graviphoton*)

super Holst-MacDowell-Mansouri action ($\mathcal{N} = 2$) [KE '21]

$$S_{\text{SH-MM}}^{\mathcal{N}=2}[\mathcal{A}] = \int_M \text{str}(F[\mathcal{A}] \wedge \mathbf{P}_\beta F[\mathcal{A}])$$

$$\mathbf{P}_\beta := \mathbf{0} \oplus \mathcal{P}_\beta \oplus \mathcal{P}_\beta \oplus \mathcal{P}_\beta \oplus \frac{1}{2\beta} (1 + \beta \star)$$

$\Rightarrow \beta$ literally has interpretation as θ -**ambiguity**!

Section 4

The chiral theory

In general: \mathbf{P}_β destroys manifest SUSY-invariance!

→ This changes in **chiral theory** ($\beta = \mp i$)

Holst projection [KE '21]

$$\mathbf{P}_{-i} : \mathfrak{osp}(1|4)_{\mathbb{C}} \rightarrow \mathfrak{osp}(1|2)_{\mathbb{C}}$$

$$M_{IJ} \mapsto T_i^+ = \frac{1}{2}(J_i + iK_i)$$

$$Q_\alpha \mapsto Q_A$$

Q_A : left-handed Weyl-fermion

Chiral Holst-MM action [KE '20, KE+HS '21]

$$S_{\text{SH-MM}}[\mathcal{A}] = \int_M \text{str}(F[\mathcal{A}] \wedge \mathbf{P}_{-i} F[\mathcal{A}])$$

- Manifestly $\text{OSp}(1|2)_{\mathbb{C}}$ -gauge invariant (similar for $\mathcal{N} = 2$!)
- SUSY partially becomes true gauge symmetry!

→ Geometric explanation of [Fülöp '94, Gambini+Obregon+Pullin '96, Ling+Smolin '99]

Chiral Theory

Super Ashtekar connection [KE+HS '21]

$$\mathcal{A}^+ := \mathbf{P}_{-i} \mathcal{A} = A^{+i} T_i^+ + \psi_r^A Q_A^r + \frac{1}{2} \hat{A}_{rs} T^{rs}$$

Chiral action

$$S_{\text{sH-MM}}^{\beta=-i} = \frac{i}{\kappa} \int_M \langle \mathcal{E} \wedge F(\mathcal{A}^+) \rangle + \frac{1}{4L^2} \langle \mathcal{E} \wedge \mathcal{E} \rangle + \underbrace{S_{\text{CS}}^{\text{OSp}(\mathcal{N}|2)}(\mathcal{A}^+)}_{\text{boundary term}}$$

\mathcal{E} : *super electric field*

Coupling bulk \leftrightarrow boundary

$$F(\underleftarrow{\mathcal{A}^+}) = -\frac{1}{2L^2} \underleftarrow{\mathcal{E}}$$


\mathcal{A}^+ natural candidate to quantize SUGRA à la LQG

- Contains both gravity and matter d.o.f. \rightarrow unified description, more fundamental way of quantizing fermions
- Substantially simplifies constraints
- **Boundary theory** described by **super Chern-Simons theory**
 \rightarrow natural candidate to study inner boundaries in LQG (\rightarrow BPS states, black holes)
- \leftrightarrow Boundary theories in string theory [Mikhailov + Witten '14]
- **Issues:** Non-compactness gauge group, reality conditions

Quantum theory: Overview

What do we need for quantum theory?


- Holonomies (parallel transport map) ✓
 - Hilbert spaces (✓)
 - Spin network states ✓
-
- Hilbert spaces gain **Krein structure** \leftrightarrow standard LQG with fermions
[Thiemann '98, Lewandowski+Zhang '21]
 - \rightarrow Gauge-invariant states: **super spin-networks**



Quantum theory: Overview

Super area operator:

$$\text{gAr}_S = \int_S \|\mathcal{E}\|$$




Action super area operator ($\mathcal{N} = 1$) [Ling+Smolin '00, KE+HS '22]

$$\widehat{\text{gAr}}_S |\Psi\rangle = -8\pi i l_{\text{Pl}}^2 \sqrt{j \left(j + \frac{1}{2} \right)} |\Psi\rangle$$

$j \in \mathbb{C}$: Principal series of $\text{OSp}(1|2)_{\mathbb{C}}$

SUSY black holes in LQG

Holst-MM in the presence of inner boundary: [KE '21, KE+HS '22]



- Geometric theory induces super-CS on inner boundary
- For $\mathcal{N} = 1$: $G = (\text{U})\text{OSp}(1|2)$

UOSp(1|2) state counting

$$N = \frac{1}{2\pi} \int_0^\pi d\theta \sin^2(2\theta) \left[4 - n + \sum_{i=1}^p n_i d_{j_i} \frac{\tan(d_{j_i} \theta)}{\tan \theta} \right] \prod_{l=1}^p \left(\frac{\cos(d_{j_l} \theta)}{\cos \theta} \right)^{n_l}$$

→ *analytic continuation*

Entropy

$$S = \ln N = \frac{a}{4} + \dots \rightarrow [\text{Bekenstein '73, Hawking '75}]$$

Section 5

Summary & Outlook

Classical theory:

- Gauge theoretic description of $\mathcal{N} = 1, 2$, $D = 4$ Holst-SUGRA + **unique** boundary terms compatible with SUSY
- \Rightarrow Chiral SUGRA has structure of **super YM** \rightarrow *super Ashtekar connection*
- Boundary theory: **Super Chern-Simons theory**

Quantum theory:

- Rigorous construction: **super spin-nets, super area operator** . . .
- Intriguing structure of state space \leftrightarrow strong similarities with standard quantization of fermions in LQG
- Application to quantum SUSY black holes [KE '21, KE+HS '22]
- Application to symmetry reduced model (*supercosmology*) [KE '21]

Difficulties:


- Implementation continuum limit (functional measure)
- Solution reality conditions?

\rightarrow *Resolvable in symmetry reduced model!*

Summary & Outlook

Outlook:

- Generalization to $\mathcal{N} \geq 3$ (*in particular*: $\mathcal{N} = 4, 8$)
- Limit $L \rightarrow \infty$ (vanishing cosmological constant)
[Concha+Ravera+Rodríguez '19]
- (Charged) supersymmetric BHs \leftrightarrow entropy calculations in string theory [Strominger+Vafa '96, Cardoso et al. '96, KE+HS '22]
- \leftrightarrow Boundaries in string theory and **OSp**-super Chern-Simons theory [Mikhaylov+Witten '14]



- [1] K. Eder, J. Math. Phys. **62** (2021), 063506 doi:10.1063/5.0044343 [arXiv:2101.00924 [math.DG]].
- [2] K. Eder and H. Sahlmann, Phys. Rev. D **103** (2021) no.4, 046010 doi:10.1103/PhysRevD.103.046010 [arXiv:2011.00108 [gr-qc]].
- [3] K. Eder and H. Sahlmann, JHEP **07** (2021), 071 doi:10.1007/JHEP07(2021)071 [arXiv:2104.02011 [gr-qc]].
- [4] K. Eder and H. Sahlmann, JHEP **21** (2020), 064 doi:10.1007/JHEP03(2021)064 [arXiv:2010.15629 [gr-qc]].
- [5] K. Eder, “Super Cartan geometry and the super Ashtekar connection,” arXiv:2010.09630 [gr-qc] (submitted).
- [6] K. Eder and H. Sahlmann, “Towards black hole entropy in chiral loop quantum supergravity,” arXiv:2204.01661 [gr-qc] (submitted).

Quantum theory: Overview

- Quantization: study \mathcal{A}^+ and associated holonomies
- turns out \rightarrow need additional **parametrizing supermanifold** \mathcal{S} to incorporate **anticommutative** nature of fermionic fields

Super holonomies [KE '20+'21]

$$h_e[\mathcal{A}^+] = h_e[A^+] \cdot \mathcal{P}\exp\left(-\oint_e \text{Ad}_{h_e[A^+]} \psi^{(\tilde{s})}\right) : \mathcal{S} \rightarrow \mathcal{G}$$

- $\mathcal{A}^+ = A^+ + \psi$ and $e : [0, 1] \rightarrow M \subset \mathcal{M}$
- $h_e[A^+]$: bosonic holonomy associated to A^+

Quantum theory: Overview

- $\mathcal{A}_{\mathcal{S},\gamma}$: set of generalized super connections

$$\begin{array}{ccc} \mathcal{A}_{\mathcal{S}',\gamma'} & \longrightarrow & \mathcal{A}_{\mathcal{S}',\gamma} \\ \downarrow & & \downarrow \\ \mathcal{A}_{\mathcal{S},\gamma'} & \longrightarrow & \mathcal{A}_{\mathcal{S},\gamma} \end{array}$$

- for fixed graph γ induces functor $\mathcal{A}_\gamma : \mathcal{S} \rightarrow \mathcal{A}_{\mathcal{S},\gamma}$
→ **Molotkov-Sachse-type smf**.
- → can study cylindrical functions and invariant measures on \mathcal{A}_γ
- covariance under reparametrization requires **Berezin-type integral** for fermionic d.o.f.
- → induces **Krein structure** on state space \leftrightarrow strong similarities to quantization of fermions in standard LQG

Symmetry reduction

- supersymmetric minisuperspace models [D'Eath + Hughes '88+'92, D'Eath + Hawking + Obregon '93]
- hybrid homogeneous isotropic ansatz for bosonic and fermionic d.o.f.
- in general: fermions not compatible with isotropy
- **But:** in (chiral) LQC can exploit enlarged gauge symmetry!
- \rightarrow natural interpretation in terms of homogeneous isotropic super connection [KE '20, KE+HS '20]

Symmetry reduced connection

$$\mathcal{A}^+ = c \, \check{e}^i T_i^+ + \psi_A \check{e}^{AA'} Q_{A'}$$

\check{e}^i : fiducial co-triad

- **Also:** can show that this is the most general ansatz consistent with reality conditions (contorsion tensor isotropic)

Loop quantum cosmology

- Construction of state space via super holonomies $h_e[\mathcal{A}^+]$ along straight edges of a fiducial cell
- \Rightarrow motivates state space of quantum theory

Hilbert space

$$\mathcal{H} = \overline{H_{\text{AP}}(\mathbb{C})}^{\|\cdot\|} \otimes \Lambda[\psi_{A'}]$$

- reality condition in quantum theory can be solved exactly! [Wilson-Ewing '15, KE+HS '20]

Quantum right SUSY constraint

$$\widehat{S}_{A'}^R = \frac{3g^{\frac{1}{2}}}{2\lambda_m} |v|^{\frac{1}{4}} \left((\mathcal{N}_- - \mathcal{N}_+) - \frac{\kappa\lambda_m}{6g|v|} \widehat{\Theta} \right) |v|^{\frac{1}{4}} \widehat{\phi}_{A'}$$

- λ_m : quantum area gap (full theory)

Loop quantum cosmology

- Quantum algebra between left and right SUSY constraint closes and reproduces Poisson algebra exactly!

Quantum algebra

$$[\hat{S}^{LA'}, \hat{S}_{A'}^R] = 2i\hbar\kappa\hat{H} - \frac{\hbar\kappa}{6g^{\frac{1}{2}}|v|^{\frac{1}{2}}} \hat{\pi}_{\psi}^{A'} \hat{S}_{A'}^R$$

- \rightarrow fixes some of the quantization ambiguities
- semiclassical limit: $\lambda_m \rightarrow 0$ i.e. corrections from quantum geometry negligible
- \rightarrow **Hartle-Hawking state** as solution of constraints \Leftrightarrow [D'Eath '98]

Hartle-Hawking state

$$\Psi(a) = \exp\left(\frac{3a^2}{\hbar}\right)$$